


J - I Adams
(Cadey)



ISSN - 2006 0718

NIGERIA POULTRY SCIENCE JOURNAL

(OFFICIAL JOURNAL OF THE WORLD'S POULTRY SCIENCE
ASSOCIATION-NIGERIA BRANCH)

2016 VOLUME 12

Editorial Office: Department of Animal Science, Ahmadu Bello University, Zaria, Nigeria.



EDITORIAL BOARD

Dr. A.V.Jegede Editor-in-Chief
Department of Animal Nutrition,
Federal University of Agriculture, Abeokuta, Nigeria

Dr. G.Adeleye Deputy Editor-in-Chief
Department of Animal Production and Health,
Federal University of Agriculture, Abeokuta, Nigeria

Dr. A.O.Osofowora Editorial Assistant
Department of Animal Nutrition,
Federal University of Agriculture, Abeokuta, Nigeria

Associate Editors

Dr. A.O.Makinde
(Poultry product and value addition sub-editor)
Department of Animal Science,
Obafemi Awolowo University Ile-Ife, Nigeria.

Prof. J.O.Aghede
(Poultry Nutrition and Feed sub-editor)
Department of Animal Production Technology,
Federal University of Technology, Akure, Nigeria.

Dr. R.Kassali
(Poultry Economics and French abstract sub-editor)
Department of Agricultural Economics,
Obafemi Awolowo University, Ile-Ife, Nigeria.

Dr. O.Adejoro
(Industry and Commentary sub-editor)
Zartech Farms Limited, Ibadan, Nigeria

Dr. A.Ladokun
(Reproduction and Physiology sub-editor)
Department of Animal Physiology,
Federal University of Agriculture, Abeokuta, Nigeria

Prof. M.O.Ozoje
(Animal Breeding and Genetics sub-editor)
Department of Animal Breeding,
Federal University of Agriculture, Abeokuta, Nigeria

Prof. M.A.Dipeolu
(Animal Health sub-editor)
College of Veterinary Medicine,
Federal University of Agriculture, Abeokuta, Nigeria

International Advisory Board

Prof. L.B.Sonaiya
Department of Animal Science,
Obafemi Awolowo University Ile-Ife, Nigeria.

Prof. M.O.Onagbesan
Department of Animal Physiology,
Federal University of Agriculture, Abeokuta, Nigeria

Prof. A.O.Adebambo
Department of Animal Breeding,
Federal University of Agriculture, Abeokuta, Nigeria

Prof. S.Yalcin
Department of Animal Science, Ege University, Turkey

Dr. H.M.Yakout
Department of Poultry Science,
Alexandria University, Egypt

Prof. E.A.Iyayi
Department of Animal Science,
University of Ibadan, Nigeria

Prof. O.Adeola
Department Poultry Science, Purdue University, Indiana,
USA

Dr V.Olori Aviagen Inc., Scotland, UK

Prof. Hui Li
North East Agricultural University, Harbin, China

Dr K.M.Shouq
Secretary, WPSA, Pakistan branch

REVIEWERS FOR VOLUME 12

Ayo-Ajasa
Matthew Whetton
Hello Lima Neto
A. V. Jegede
Daisy Enyebetine
O.M. Sogunle
O.M.O. Idowu
Oluseyi O. Odugwu
Abimbola O. Ladokun

Akinola A. S
Sobayo, R.A.
Adedire, M.A.
Jegede, A.V.
A.O. Talabi
Adebayo S. Bamire
Okanlawon M. Onagbesan
O.A. Yesufu
Rabirou Kassali
Oluwatunminnu A. Adebambo

Johnson O. Agbede
A.A. Odunsi
A. O. Oso
Akinbode
Egbeyale
C.O.Betiku

NIGERIA POULTRY SCIENCE JOURNAL

ISSN – 2006 0718

Typeset and formatted by
TKBV Business Centre
64, Obafemi Awolowo Road,
Opposite DSTV Office, Isale-Igbehin, Abeokuta
mobile: 08064255063

Printed by
TOLUKOYA PRINT HOUSE,
64, Obafemi Awolowo Road,
Opposite DSTV Office, Isale-Igbehin, Abeokuta
mobile: 08064255063; 08170629325

Published by
THE WORLD'S POULTRY SCIENCE ASSOCIATION - NIGERIA BRANCH

ABOUT NPSJ

Introduction

Nigerian Poultry Science Journal (NPSJ) is an international journal published by the World Poultry Science Association-Nigeria Branch (WPSA-NB). It reports original work, review and commentaries on a broad range of poultry topics including nutrition, breeding and genetics, health, welfare, physiology, reproduction, processing, economics, education, farming and any other relevant topic. The journal is published in English of the abstract, in both online and hard copy media.

Medium and Frequency of publication

Four issues per year in March, June, September and December. Issues will be available online at www.wpsanb.org.ng/npsj and as print version. Online edition will appear prior to the print edition.

Scope of journal

The journal accepts and publishes articles in all fields of poultry production as original, short/rapid communication, commentary and review articles.

Disclaimer

- a. All statements, conclusions and opinions expressed in individual articles in NPSJ are those of the authors and do not in any way represent the position and endorsement of the publisher and thus the journal accept no responsibility or liability on them.
- b. The journal will use its best endeavours to ensure that work published is that of the named authors and that the published results and conclusions are consistent with the primary data. However, it takes no responsibility for fraud or inaccuracy on the part of the authors. Papers reporting work with animals must include a reference to the code of practice adopted for the experimentation. It is expected that reported experiments have been performed according to appropriate ethical and legal standards. The journal reserve the right not to publish such experiments that violate ethical and welfare issues and take no responsibility for the breach of such right by any reported experiment.

Paper charges and subscription NPSJ charges for publication of their article. The publication charge and subscription rate are subject to review at anytime depending on the prevailing cost of production. The current charge is given below:

	Nigerian Naira	US dollar
Processing fee	N1,000	Free
Article	N800 per text page and N900 per table page	\$ 5 per page
Coloured image	N1,000 per page	\$10 per page
Subscription rate	N1,500 per issue	\$ 25 per issue

Editorial Office

Dr. A. V. Jegede

Department of Animal Nutrition,
Federal University of Agriculture,
P.M.B. 2240, Abeokuta, Nigeria.

vimbjegede@yahoo.com
jegedeav@funab.edu.ng

NIMAL NUTRITION

rowth performance, carcass yield and gut microbial status of broiler chickens
d diets containing *Morinda lucida*, *Xylopia aethiopica* and *Syzygium aromaticum*
phytobiotics

56 - 62

LAYEMI W.A., GAFAR, I.I., AWE, O.M., AKAPO O.A., BAMGBOSE A.M.

rowth performance and nutrient digestibility of cockerels fed diets containing
varying levels of water hyacinth [*Eichhornia crassipes* (Martius) Solms-Laubach]
meal supplemented with exogenous enzymes

63 - 70

ALIK, A. A., KUDU, Y. S., MUSA, S. J. AND OLAGUNJU, K. T.
Microbial Evaluation of Earthworm Meal as a Protein source in diets for
broiler Chickens

71 - 76

DOWU K.R., AGUNBIADE J.A., ADEYEMI O.A., AND AJAYI O.L., FAFIOLU
O., OLADIPUPO O.O.

NIMAL PHYSIOLOGY

thermotolerance acquisition in broiler chickens through early feed restriction:
response to road transportation

77 - 83

MONSURU OLADIMEJI ABIOJA, OLUBUNMI GRACE SODIPE, JOHN
ADESANYA ABIONA, OLUWAGBEMIGA OLANREWAJU ADELEYE,
OYEGUNLE EMMANUEL OKE, ABIMBOLA OLUWASEUN LADOKUN, JAMES
OLAMITIBO DARAMOLA, TOLULOPE JULIUS WILLIAMS AND OLUSEGUN
AYODEJI OSINOWO

Assessment of different garlic extracts on fertilizing potentials of extended rooster
semen used for artificial insemination

84 - 91

BALOGUN, ADEDEJI SULEIMON, JIMOH, OLATUJI ABUBAKAR, MUSTAPHA
ABDUL-RAHMAN, MOHAMMED ABDULRAHMAN, AKINOSUN, AKIN
AKINBOBOLA, AND OJO, J.A.

Variation in some blood biochemical indices in laying and non
muscovy duck

92 - 96

OGAH D. M.

Effect of drinking grain steep water on performance, nutrient utilization and
carcass characteristics of broiler chickens

97 - 102

OPOWOYE, I.O., ATTELL, J.O., KOLAWOLE, O.M. ADELEYE, O.O., OKUKPE,
K.M., SOLA-OJO, E.E., ADERIBIGBE, T.A. AND ALLI, O.I.

Effect of fluted pumpkin (*Telfaria occidentalis*) leaf extract on growth
performance serum chemistry and organ weights of cockerel chickens

103 - 108

LADOKUN, A.O., ADEWALE, N.O., ALABI, O.M. AND ABIONA, J.A.

Growth performance and nutrient digestibility of cockerels fed diets containing varying levels of water hyacinth [*Eichhornia crassipes* (Martius) Solms-Laubach] meal supplemented with exogenous enzymes

*MALIK, A. A., KUDU, Y. S., MUSA, S. J. AND OLAGUNJU, K. T.

Department of Animal Production, Federal University of Technology, Minna, Niger State

*Corresponding Author's e-mail: delemalik@gmail.com

Phone: 08030637763

Abstract

A six weeks' feeding trial was conducted on 288 two-weeks'-old Black Harco cockerel chicks to evaluate their growth performance when fed diets containing water hyacinth (*Eichhornia crassipes*) meal (WHM) at 0, 100, and 200 g/kg dietary inclusion levels, with no exogenous enzymes supplementation (Diet 1, Diet 2 and Diet 3 respectively) and then with exogenous enzymes (Maxigrain®) supplementation (Diet 4, Diet 5 and Diet 6 respectively); as a replacement for wheat offal (WO). The birds were randomly allocated to the six dietary treatments, with three replicates per diet and 16 chicks per replicate, and fed ad-libitum for six weeks. At the end of the feeding trial, three birds per replicate were randomly selected, removed from the brooder and placed in metabolism cages for digestibility studies. They were kept in the cages for three days to enable them adjust adequately to the conditions in the cages, followed by faecal collection that lasted for four days. Results show that there were no significant ($P > 0.05$) differences in initial body weight, final body weight, average body weight gain, total body weight gain, total feed intake, feed conversion ratio and protein efficiency ratio between birds fed 0, 10 and 20 % dietary inclusion levels of WHM. There were no significant ($P > 0.05$) enzyme and water hyacinth level interactions for these parameters, indicating that exogenous enzymes addition did not significantly ($P > 0.05$) affect feed intake, body weight gain and feed conversion ratio in cockerels fed WHM diets. Crude fibre digestibility decreased significantly ($P < 0.05$) as the dietary WHM increased, indicating decreased utilization; while the exogenous enzymes addition increased the digestibility of the nutrients. It is therefore concluded that based on growth performance parameters, WHM can effectively replace WO 100 % in the diets of cockerels without exogenous enzymes addition; however, better nutrient digestibility are obtained when the WHM-based diets are treated with exogenous enzymes.

Key words: Water hyacinth meal, cockerels, exogenous enzymes, growth performance, nutrient digestibility.

Introduction

Poultry farmers in some parts of the world usually prefer cockerel farming due to low cost of day-old chicks; besides, cockerels require less floor space and feed and there is higher price for cockerel meat than that of broiler meat. It is also believed that cockerels are less susceptible to diseases in comparison to broilers (Sil *et al.*, 2002).

Water hyacinth is a free floating perennial aquatic plant of *Eichhornia* genus in the *Pontederiaceae* family. It is native to tropical and sub-tropical South America, and is one of the fastest growing plants known; and are produced primarily by way of runners or stolons that form daughter plants. It grows profusely as a noxious aquatic weed, forming

dense mats that can spread across water surface eventually choking an entire water body, if allowed to proliferate. It is the most invasive aquatic weed species known and have become a terror and a menace in many tropical and subtropical countries of the world, including Nigeria. Many international conferences and symposia have been held on its complete eradication and control, but not much success have been achieved practically; the weed continue to spread at an alarming rate. In fact, the presence of water hyacinth disrupts all life in the water (Center *et al.*, 1999), causing high evaporation rate and loss of water. Shoeb and Singh (2004) reported that under favourable conditions, water hyacinth

can achieve a growth rate of 17.5 metric tons per hectare per day.

It is estimated that about 70-75 % of the cost of production is incurred in the procurement of feed and feed ingredients for poultry (Idowu et al., 2005). To minimize this, there is an urgent need for animal scientists and feed nutritionists to address the high cost incurred on feeds and feed ingredients by sourcing for non-conventional feed ingredients that are cheaper and nutritive, and can replace the conventional ones. Fortunately, water hyacinth plant, though a weed, has high nutritive values similar to high quality forage (Dung, 2001). According to Malik et al. (2014), water hyacinth meal (WHM) when compared to wheat offal (WO) has a higher CF content (21 % versus 11 %), lower CP content (14 % versus 16 %), higher ash content (24 % versus 6 %) and comparable ether extract (5 % versus 4 %) and metabolizable energy content (1901 Kcal/kg versus 1845 Kcal/kg). The challenge therefore is converting a weed of nuisance value into a valuable feed resource for poultry, as a non-conventional feed ingredient.

Several exogenous enzymes have been used by different workers in nutrient utilization and retention studies for different categories of farm animals, especially when fed fibrous feedstuffs.

They have been used to increase the availability of nutrients by retarding the adverse effects of anti-nutritional factors present in feed components (Munir and Maqsood, 2013). Enzymes are added to animal ration with the goal of increasing its digestibility, removing anti-nutritional factors and improving nutrient availability; and a large number

Table 1: Proximate composition of water hyacinth meal (WHM)

Parameter	Composition (g/kg)
Dry matter	949.00
Crude protein	144.10
Crude fibre	213.30
Ether extract	30.00
Ash	231.10
Nitrogen free extract	330.50

of carbohydrases, proteases, phytases and lipases are used for this purpose (McCleary, 2001). Therefore, the objective of this research study is to evaluate the growth performance and nutrient digestibility of cockerels fed diets containing varying levels of WHM (as a replacement for wheat offal) supplemented with exogenous enzymes at the chick phase.

Materials and Methods

Preparation of Water Hyacinth Meal

Water hyacinth meal (WHM) was prepared using the procedure of Malik et al. (2014). Whole plants of water hyacinth were collected from the surface of the River Niger by hand and using local canoes at Nkukuso village, near Bida, in Lavun Local Government Area of Niger State. The harvested green plants were sun-dried for a few days at the river bank until they were properly dried. Foreign extraneous materials such as leather wastes, empty tin containers, iron materials and dried weeds were manually removed. They were then packaged in polythene sacks and transported to the Animal Production Laboratory of the Federal University of Technology, Minna, for further processing. In the lab, they were oven dried at a temperature of 80 °C for between 18 to 24 hours to a moisture content of less than 10 %. The dried samples were then milled with an attrition mill and sieved using a 2 mm sieve to obtain the water hyacinth meal (WHM); which was stored in plastic containers with tight-fitting lids until needed for use in formulating the experimental diets. Table 1 gives the proximate composition of the WHM.

Experimental Location

This Research Study was carried out at the Poultry Unit of the Animal Production Teaching and Research Farm, Federal University of Technology, Minna. Minna is the capital city of Niger State of Nigeria, with an estimated population of about 500,000 and a land area of about 6,784 square kilometers. It is located in the Southern Guinea Savanna Vegetation Zone, between Latitude 9°37' North and Longitude 6°33' East. Its mean annual rainfall is 1300 mm, taken from an exceptionally long record of 50 years. Temperature rarely falls below 22 °C; the peaks are 40 °C (February - March) and 35 °C (November - December). The rainy season starts in April and lasts between 190 and 200 days (2009-2013 Postgraduate School Prospectus, Federal University of Technology, Minna).

Experimental Diets

The experimental diets were formulated as follows: Diet 1, 2 and 3 contained 0, 100 and 200 g/kg dietary inclusion levels of WHM (replacing 0, 50 and 100 % wheat offal respectively) with no exogenous enzymes (Maxigrain®) added; while

Diet 4, 5 and 6 contained 0, 100 and 200 g/kg dietary inclusion levels of WHM (replacing 0, 50 and 100% wheat offal respectively) with Maxigrain® added at the rate of 100 g per tonne of the mixed feed. Other feed ingredients used were purchased from Minna Central Market and from other feed ingredients depots within Minna metropolis. Table 2 shows the gross composition of the experimental diets (g/kg diet).

Experimental Animals and their Management

A total of 288 day-old *Black Harco* cockerels were obtained from Minna branch depot of Avian Specialties Limited, Ibadan. Before the arrival of the birds, the deep litter house was thoroughly washed and disinfected and the floor was covered with wood shaving up to 10 cm depth. Drinkers and feeders were set in place in the pens and sources of heat provided using stove and coal pot. The deep litter house was then covered with polythene sheets which provided adequate brooding environment for

the birds (ambient temperature was about 32 °C). On the arrival of the chicks, the birds were unboxed and counted; and antibiotics, multivitamins, vitamin C and glucose were administered in their drinking water as anti-stress and source of energy to stimulate feed consumption. The birds were then fed with a commercial pullet chick mash for two weeks, after which they were weighed, before being introduced to the six different experimental diets. 48 birds were randomly selected and allotted to each treatment; with 16 birds per replicate and three replicates per treatment. They were fed for six weeks. During the course of the experiment, clean drinking water and fresh feed were supplied to the birds *ad libitum*. A standard vaccination programme was administered to the birds from day old till the end of the brooding period to ensure that the birds were immunized against major poultry diseases. Records of feeds intake, body weight gain, and left over feed were taken.

Table 2: Composition of the experimental diets (g/kg)

WO replacement	Without enzymes			With enzymes		
	0%	50%	100%	0%	50%	100%
Ingredients (g/kg)	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	Diet 6
Maize	453.90	453.90	453.90	453.90	453.90	453.90
Groundnut cake	251.10	251.10	251.10	251.10	251.10	251.10
Wheat offal (WO)	200.00	100.00	0.00	200.00	100.00	0.00
Water hyacinth meal	0.00	0.00	200.00	0.00	100.00	200.00
Fish meal	20.00	20.00	20.00	20.00	20.00	20.00
Palm oil	10.00	10.00	10.00	10.00	10.00	10.00
Lysine	5.00	5.00	5.00	5.00	5.00	5.00
Methionine	5.00	5.00	5.00	5.00	5.00	5.00
Bone meal	45.00	45.00	45.00	45.00	45.00	45.00
Limestone	5.00	5.00	5.00	5.00	5.00	5.00
*Premix	2.50	2.50	2.50	2.50	2.50	2.50
Salt	2.50	2.50	2.50	2.50	2.50	2.50
Determined Analysis						
Dry matter	893.00	901.00	902.00	905.00	879.90	899.00
Crude protein	196.00	200.00	206.50	206.50	206.50	203.00
Crude fibre	60.00	86.70	93.30	60.00	80.00	86.70
Ether extract	160.00	155.00	145.00	165.00	165.00	160.00
Ash	105.00	100.00	126.70	100.00	100.00	111.50
Nitrogen free extracts	372.00	359.30	330.50	373.50	328.40	337.80

*Each 2.5 kg of premix is made of the following: Vitamin D₃, 10,000,000.00 IU; Vitamin E, 2,000,000.00 IU; Vitamin K₃, 23,000.00 mg; Vitamin K₁, 2,000.00 mg; Vitamin B₁, 1800.00 mg; Vitamin B₂, 5,500.00 mg; Niacin, 27,500.00 mg; Panthothenic acid, 7,500.00 mg; Vitamin B₆, 3,000.00 mg; Vitamin B₁₂, 15.00 mg; Folic acid, 750.00 mg; Biotin, 60.00 mg; Choline, 300,000 mg; Cobalt, 20.00 mg; Copper, 2,000.00 mg; Iodine, 1,000.00 mg; Iron, 20,000.00 mg; Manganese, 40,000.00 mg; Selenium, 200.00 mg; Zinc, 30,000.00 mg; Antioxidant, 1,250.00 mg.

Parameters Determined

The following parameters were determined using the procedures of Adesida *et al.* (2010). Average feed intake was obtained by subtracting the quantity of the left-over (uneaten) feed from the quantity of feed supplied to the birds per day. Weekly body weight gain was determined by subtracting the body weight of the birds in the preceding week from the body weight of the birds in the following week. Feed conversion ratio (FCR) was obtained by dividing the average feed intake per bird per week by the average body weight gain per bird per week for each treatment while protein efficiency ratio (PER) was determined by dividing the body weight gain by the birds by their protein intake.

Digestibility Studies

At the end of the six weeks' feeding trial, three birds per replicate and a total of nine birds per treatment, were randomly selected, removed from the floor and placed in metabolism cages for three days, for them to adjust to the conditions in the cages. Measured quantities of fresh feed were then given to the birds daily, while the droppings were collected 24 hours later (and for four days) using aluminum foils, based on the total collection method. Collected droppings were then oven-dried at 80 °C for 24 hours to get a constant weight. The oven dried faeces were then packaged in polythene bags and stored in plastic containers in a deep freezer until needed for analysis. Apparent digestibility of nutrients (ADN) was calculated using the formula described by Isikwenu *et al.* (2010) thus:

$$ADN = \frac{\text{Nutrient intake} - (\text{Nutrient in faeces voided})}{\text{Nutrient intake}} \times 100$$

Nutrient intake

Total digestible nutrient (TDN) was calculated using the formula given by Church and Pond (1988) thus:

$$TDN = \text{Digestible crude protein} + \text{Digestible NFE} + 2.25 \times \text{Digestible ether extract}$$

Chemical Analysis

The proximate composition of the water hyacinth meal (WHM), the experimental diets and the collected faecal droppings were determined using the procedures of AOAC (1990).

Statistical Analysis

Data obtained from this research study was subjected to a one-way analysis of variance (ANOVA) based on a 2 x 3 factorial model using the Statistical Analysis Software (SAS, 2000, Version

6). Where means were significant, they were separated using the Duncan Multiple Range Test (Duncan, 1955).

Results and Discussion

There were no significant ($P > 0.05$) differences in initial body weight (IBW), final body weight (FBW), total body weight gain (TBWG), average body weight gain (ABWG), total feed intake (TFI), feed conversion ratio (FCR) and protein efficiency ratio (PER) between enzyme-supplemented and non-supplemented diets; and between 0, 100 and 200 g/kg dietary inclusion levels of WHM (Table 3). This result does not quite agree with the findings of Alam *et al.* (2003) who reported that cockerels fed on enzymatic diets had a tendency to convert feed more efficiently to live weight than those without enzyme supplementation. However, according to an earlier study by Malik *et al.* (2014), the crude fibre component of WHM is made up of 24.60% cellulose, 26.08% hemicellulose and 12.86% lignin; and has NDF and ADF values of 63.54 and 37.46 % respectively. Each gramme of Maxigrain^{*} contains 10,000 IU cellulase, 200 IU β-glucanase, 10,000 IU xylanase and 2,500 FTU phytase. This enzyme composition may not be very effective in degrading the high cellulose (24.60%), high hemicellulose (26.08%) and high lignin (12.86%) fibre component of WHM at high dietary inclusion levels of WHM (100 and 200 g/kg diet) in cockerel diets, hence the non-significant difference in growth performance among the birds fed those diets.

In terms of nutrient digestibility, crude fibre and nitrogen free extracts were better utilized by cockerels fed the enzyme-supplemented diets than those fed diets without enzyme supplementation; while total digestible nutrient (TDN) was significantly ($P < 0.05$) higher for birds fed diets with 0 g/kg WHM than birds fed the 100 and 200 g/kg WHM diets. This may be due to the fact that exogenous enzymes supplement the digestive enzymes of monogastric animals by aiding the breakdown of non-starch polysaccharides, protein and anti-nutritional factors thereby increasing their nutritional value by making available the nutrients to the birds (Chot, 2006; Giraldo *et al.*, 2008). According to Buchanan *et al.* (2007), exogenous enzymes hydrolyze non-starch polysaccharides (NSPs)

Table 3: Main effects of exogenous enzymes supplementation and feeding graded levels of water hyacinth meal on the growth performance of cockerels

Treatment	IBW (g)	FBW (g)	TBWG (g)	ABWG (g)	TFI (g)	AFI (g)	FCR	PER
ENZYME (E)								
0	31.94	491.31	459.36	8.21	1922.00	34.32*	4.29	1.19
1	31.59	479.98	448.39	8.00	2073.00	37.02*	4.67	1.54
SEM	0.55	23.56	23.97	0.43	106.13	1.89	0.42	0.45
LOS (0.05)	NS	NS	NS	NS	*	NS	NS	NS
HYACINTH (H)								
0%	32.81	508.93	476.12	8.51	1969.30	35.17	4.26	1.22
10%	31.25	484.07	452.82	8.08	1968.30	35.15	4.38	1.17
20%	31.55	463.94	432.69	7.75	2054.90	34.66	4.79	1.72
SEM	0.67	28.85	29.35	0.52	129.95	2.52	0.61	0.55
LOS (0.05)	NS	NS	NS	NS	NS	NS	NS	NS
INTERACTION								
EX-H.	NS	NS	NS	NS	NS	NS	NS	NS

*Means in the same column with different superscripts were significantly ($p<0.05$) different

IBW (g/bird) = Initial body weight.

ABWG (g/bird/day) = Average body weight gain.

ICR = Feed conversion ratio.

LOS = level of significance

FBW (g/bird) = Final body weight.

TFI (g/bird) = Total feed intake.

PER = Protein efficiency ratio

NS = Not significantly different

TBWG (g/bird) = Total body weight gain
AFI (g/bird) = Average feed intake
SEM = Standard error of means

Table 4: Main effects of exogenous enzymes supplementation and feeding graded levels of water hyacinth meal on the nutrient digestibility of cockerels (%)

Treatment	Dry matter	Crude Protein	Crude Fibre	Ether Extract	Ash	NFE	TDN
ENZYME (E)	0	88.93	87.69	69.94 ^b	96.01	90.16 ^a	81.42 ^b
SEM	1.27	89.27	86.88	86.67 ^a	96.71	80.70 ^b	86.52 ^a
LOS (α .05)	NS	1.71	4.38	0.45	2.23	0.81	0.73
HYACINTH (H ₂)	0%	*	NS	*	*	*	*
10%	89.67	85.76	75.14	96.09	90.24	90.69	86.42 ^a
20%	88.62	85.81	77.36	96.21	82.15	91.14	83.54 ^b
SEM	89.02	88.40	82.41	96.79	83.01	92.23	82.16 ^b
LOS (α .05)	1.55	2.10	5.37	0.56	2.73	0.99	0.89
INTERACTION E X H	NS	NS	NS	NS	NS	NS	*
EX H	NS	NS	NS	NS	NS	NS	*

^a Means in the same column with different superscripts were significantly ($p<0.05$) different

SEM = Standard error of the means

LOS = Level of significance

TDN = Total digestible nutrient

NS = not significantly different

which might be potentially used by the animal, increasing the usage of feed energy, therefore increasing the digestibility of all nutrients.

Recommendations

It is therefore recommended that based on growth performance parameters, WHM can be included up to 200 g/kg in the diet of cockerels at the starter phase (effectively replacing wheat offal 100 % in the diets of cockerels) without exogenous enzymes addition. However, better nutrient digestibilities are obtained when the WHM-based diets are treated with exogenous enzymes from Maxigrain®.

References

- ADESIDA, S. A., AGUNBIADE, J. A. & ENJIORUNDA, O. O. 2010. Effect of palm kernel cake and deoiled palm kernel cake on the performance of cockerels. *Nigerian Journal of Animal Production*, 37(1): 34-39.
- ALAM, M. J., HOWLIDER, M. A. R., PRAMANIK, M. A. H. & HAQUE, M. A. 2003. Effect of exogenous enzyme in diet on broiler performance. *International Journal of Poultry Science*, 2, 168-173.
- A.O.A.C. 1990. Association of Official Analytical Chemists. *Official Methods of Analysis*, 15th edition, Washington DC.
- BUCHANAN, N. P., KIMBLER, L. B. & PARSONS, A. S. 2007. The effects of non-starch polysaccharide enzyme addition and dietary energy restriction on performance and carcass quality of organic broiler chickens. *Journal of Applied Poultry Research*, 16, 1-12.
- CENTER, T. D., DRAY JNR, G. A., JUBINDSKY, G. P. & DRODOWITZ, M. J. 1999. Biological control of water hyacinth under conditions of maintenance management: Can herbicides and insects be integrated? *Environmental Management*, 23: 241-256.
- CHIOT, M. 2006. Enzymes for the feed industry, past, present and future. *World's Poultry Science Journal*, 62(2): 5-15.
- CHURCH, D. C. & POND, W. G. 1988. *Basic Animal Nutrition and Feeding* (3rd edition). New York, USA: John Wiley & Sons.
- DUNCAN, D. B. 1955. Multiple Ranges and Multiple F-Tests. *Biometrics*, 11: 1-42.
- DUNG, N. N. X. 2000. *Evaluation of green plants and by products from the Mekong delta with emphasis on fibre utilisation by pig*. Dissertation Uppsala: Swedish University of Agricultural Sciences, Agraria 285.
- GIRALDO, I. A., TEJIDO, M. I., RAMILLA, M. J., RAMOS, S. & CARRO, M. D. 2008. Influence of direct-fed fibrolytic enzymes on diet digestibility and ruminal activity in sheep fed grass hay-based diet. *Journal of Animal Science*, 86: 1617-1623.
- IDOWU, O. M. O., BAMGBOSE, A. M., JEGEDE, A. V., IDOWU, O. A., SULE, R. AND GRUVBETIM, D. 2005. Effect of replacement of cassava peels with cassava roots on performance and egg quality characteristics of laying hens. *Proceedings of the Annual Conferences of the Animal Science Association of Nigeria (ASAN)*, Sept. 12th-15th, 2005, pp. 136-139.
- ISIKWENU, J. O., AKPODIETE, O. J., OMEJE, S. I. & OKAGBARE, G. O. 2010. The effects of replacing groundnut cake with urea-treated and fermented brewers' dried grains on nutrient digestibility, retention and carcass characteristics of broiler finishers. *Nigerian Journal of Animal Production*, 37(1): 1-12.
- MALIK, A. A., AREMU, A., AYANWALE, B. A., IJAIYA, A. T., BADMUS, A. H. A. & DIKKO, A. H. 2011. An evaluation of water hyacinth [*Eichhornia crassipes* (Martius) Solms-Laubach] meal as a feedstuff for pullet chicks as determined by carcass and haematological characteristics. *International Journal of Advanced Biological Research*, 4(2): 114-121.
- MCCLEARY, B. V. 2001. *Analysis of feed enzymes*. In: *Enzymes in Farm Animal Nutrition*. Eds. Belford, M. R. & Patridge, G. G. UK: CAB International, p.406.
- MUNIR, K. & MAQSOOD, S. 2013. A review on role of exogenous enzymes supplementation in

- Malik et al. Growth and nutrient digestibility of cockerels fed varying levels of water hyacinth with enzyme
poultry production. *Emirate Journal of Food and Agriculture*, 25(1): 66-80.
- SAS 2000. Statistical Analytical Systems. *User's Guide*, SAS Institute, Cary, North Carolina, USA.
- SHOEB, F. & SINGH, H. J. 2000. Kinetic studies of biogas evolved from water hyacinth. 2nd International Symposium on New Technologies for Environmental Monitoring and Agro-applications, p. 138.
- SIL, G. C. DAS, P. M. ISLAM, M. R. & RAHMAN, M. M. 2002. Management and disease problems of cockerels in some farms of Mymensing, Bangladesh. *International Journal of Poultry Science*, 1: 102-105.