

TITLE PAGE

**APPLICABILITY OF AERIAL PHOTOGRAPHS TO
URBAN CHANGE DETECTION IN NARAGUTA AREA
OF THE JOS PLATEAU**

By

**Shehu Yunusa Adam
M. TECH/SSSE/128/96/97**

**Being Thesis submitted to department of Geography Federal
University of Technology Minna**

**In Partial Fulfillment for the award of
Masters of technology degree in remote sensing
Applications
JUNE, 2004**

CERTIFICATION

This is to certify that this research was carried out by Shehu Yunusa Adam (M. Tech/SSSE/128/96/97) as part of the requirement for Master of Technology (M. Tech) degree in Remote Sensing Applications, Geography Department, School of Science and Science Education, Federal University of Technology, Minna.

Dr. Akinyeye, P. S
Supervisor


.....
Signature and Date

Dr. M.T. Usman
H.O.D (Geography Dept)


.....
Signature and Date

Prof. J. A. Abalaka
Dean, Post Graduate School


.....
Signature and Date

Dr. S. A. Mashi
External Examiner

.....
Signature and Date

APPROVAL

This thesis has been read and approved as meeting the requirements of the school of
nce and science education (Geography Department)

.....
External Examiner

.....
Date

.....
Head Department

.....
Date

DEDICATION

This work is dedicated to my beloved Sayyada Rahmat and to my parents; Alhaji Shehu
Amad Adamu and Hajiya Jamila Shehu Adamu.

ACKNOWLEDGEMENT

A research work of this kind normally owes much to a great many people. I owe my gratitude to Almighty Allah, the most beneficent and the most merciful for giving me the will, the energy and the resources to carry out this great work.

I am greatly indebted to my project supervisor; Dr. Akinyeye, P. S for his guidance, tolerance, detailed and valuable criticisms and suggestions inspite of his busy schedule.

My Sincere gratitude goes to my lecturers, Dr. M. T. Usman (H.O.D), Dr. Okhumamhe A. A, Dr. G. N. Nsorfor, Dr. Umoh T. Umoh and Dr. Abubakar Sadauki. I owe my sincere gratitude to Dr. Halilu S. Ayuba for his inspiration, patience and guidance in this work

I most sincerely thank my elder brothers; Engr. Musa Hassan, Engr. Sabo Shehu and Sheikh Suleiman Adamu for their support and encouragement throughout the programme. I also thank my younger brothers; Sheikh Mohammed Alhaji and Ahmed Sani for all their assistance from the beginning of this programme up to the end.

To my friends Mohammed Alli, Shehu Inuwa, Musa Isah, Mohammed Ndanusa, Audu Bello, Yakubu Inuwa, Sani Aliyu, Miss Aishatu Mohammed, Miss Philomena Oyekezie and Mallam Musa B. Umar (ADP Computer Supervisor). I Say Thanks.

To Captain K. M. Maikundi, Tukur Kele, Idris Mohammed Kontagora, Sani Tanko, Mustafa Gambo, Alhaji Garba O. Ibrahim and Kabiru Ahmadu. I remain grateful to my friends and colleagues Mr. Akin Buari, Mr. Solomon Ndache and Abdulkadir. I

thank you. I also thank Mrs. Huleira Mohammed Ali and my uncles; Shehul Hadi Zariff and Sheikh Ibrahim A. Bako.

And to my friends and relations who in one way or the other contributed not only to this research work but also the programme whose name owing to lack of space did not feature here. I thank you and may Allah bless you all.

Yunusa Shehu Adamu

June, 2004

ABSTRACT.

Urban areas grow upwards and outwards. These growth are accompanied by both positive and negative consequences on the environment as well as the population living within. Problem such as poor road networks, poor health and sanitary conditions, decaying social amenities, traffic congestion, in adequate housing, conflicting land uses etc. These are the results of poor urban planning and uncontrolled development, which characterized many towns and cities in Nigeria. To understand the present urban landscape, we have to discover why towns and cities have grown to their present pattern. For this study aerial photographs of 1975 and 1991 were used to determine the extent and amount of change and the consequences on the study area, Naraguta area of Jos Plateau. These photographs were interpreted manually and a classification scheme was developed and used to classify the different urban categories. A change map produced from the overlay of both aerial photographs and the changes within the period was calculated from the statistics generated from the techniques employed. These changes were graphically represented to show the distribution and change in the categories. These result showed that 8.384 hectares (45.8%) of the study area had changed over the period to 6.592 hectares (21.14%) which was a decrease in the under developed areas. This study has demonstrated that remote sensing can provide the bio-physical information necessary for monitoring urban change detection.

TABLE OF CONTENTS

	<i>page</i>
ification - - -- - - - - -	i
roval - - - - - - - -	ii
ication - - - - - - - -	iii
nowledgement - - - - - - - -	iv
tracts - - - - - - - -	vi
le of Contents - - - - - - - -	viii
of figures - - - - - - - -	ix
of table - - - - - - - -	x
of plates - - - - - - - -	xi
 APTER ONE	
Introduction - - - - - - - -	1
Background to the Study - - - - - - - -	1
Problem Statement - - - - - - - -	2
Aim and Objectives of Study - - - - - - - -	3
4 Justification for the use of Remote Sensing to the Study- -	4
5 Description of the Study Area- - - - - - - -	5
6 Structure of the Project - - - - - - - -	10
 HAPTER TWO:	
0 Literature Review -- - - - - - - -	12
 HAPTER THREE	
0 Research Methodology and Materials- - - - - - - -	19
.1 Research Materials - - - - - - - -	19
.2 Principles of Interpretation - - - - - - - -	20
.3 Methodology - - - - - - - -	21
.4 Data Analysis - - - - - - - -	22
.5 Classification Scheme - - - - - - - -	22

Ground Truth	-	-	-	-	24
APTER FOUR					
Results and Introduction	-	-	-	-	25
Interpretation of 1975 Aerial Photographs	-	-	-	-	25
Interpretation of 1991 Aerial Photographs	-	-	-	-	30
Urban Landuse Changes Between 1975 - 1991	-	-	-	-	33
Discussion of Results	-	-	-	-	38
Consequences of the changes	-	-	-	-	42
CHAPTER FIVE					
0 Summary, Recommendation And Conclusion	-	-	-	-	44
1 Summary	-	-	-	-	44
2 Conclusion	-	-	-	-	45
3 Recommendation	-	-	-	-	45
lates	-	-	-	-	47
ferences	-	-	-	-	52

LIST OF FIGURES

I	Plateau State Map Showing The Study Area -	-	6
4.1	Urban Map of Naraguta, Jos 1975	-	28
4.2	Percentage distribution of Urban Categories -	-	29
4.3	Urban map of Naraguta, Jos 1991.	-	31
4.4	Percentage distribution of Urban Categories 1991	-	32
4.5	Urban Change Map of Naratuta, Jos 1975-1991	-	36
4.6	Relative distribution of Urban Categories (1975-1991)	-	37

LIST OF TABLES

Urban Landuse Classification Scheme	-	-	23
Urban Categories Aerial Coverage and percentage 1975 and 1991	-	-	26
Distribution of Urban Change Categories and Magnitude 1975-1991			34
Percentage Distribution of Urban Categories 1975 and 1991	-	-	35
Proportion of Change of Urban Category Classes to overall Change-			35

LIST OF PLATES

nd c.	Negative impact of Urbanisation on Streams across Bauchi Road	49
	Racecourse of Naraguta Jos after 1975	49
	Bauchi Road Motor Park	49
	Main Entrance of University of Jos	49
	Well -organised Parking Space	49
b, c.	Modern Road Networks	49
nd b	Commercial Areas	50
b and c	New Building Structures	51

CHAPTER ONE

1.0 INTRODUCTION

1.1 BACKGROUND TO THE STUDY

One of the basic issues in the development process is resource allocation and a major resource, which serve as a common base to the entire development efforts is land. Land is a fundamental necessity of life for it is the very framework on which the socio-political and economic activities of a people function. Thus, an essential prerequisite to any development effort therefore, according to Adeniyi, (1981) is the appraisal of the existing land use situation and how it has been changing overtime.

The term Land use relates to the human activities associated with a piece of land; (Lillesand and Kiefer, 1979), while land cover have been described by Lo, (1986) as the vegetational and artificial constructions covering the land surface. Although landuse is usually not directly visible from the imagery, landcover is directly visible from the remotely sensed imagery. Landuse and landcover can be described as urban or built up lands which may be residential, commercial, industrial, transportation or institutional.

Urban areas can be described as those areas that constitute part of a town or a city. Urban areas have higher population density; they are centres of agglomeration of socio-cultural and economic functions. They are devoid of primary economic activities and are the center of political administration. Urbanization and modernization processes with the increase in demand for land, housing and transportation induced radical change in their pattern.

The technological advancement of the world today has reached a point where man can disrupt and change totally the environment in which he lives. The occurrence of these

potentials to disrupt and change the environment has served to stimulate and focus public attention upon environmental issues. One major way in which human activities easily disrupts the working of the environment is through urban development. To understand the extent of such disruptions, there is need for monitoring of changes occurring in area of urban development. A good way to carry out such a monitoring is to employ remote sensing techniques.

The application of remote sensing techniques for urban change detection, monitoring, mapping and analysis requires continuous acquisition of data for the formulation of policy and programmes. There is therefore an urgent need to have timely, accurate and cost effective source of data and this can be obtained through satellite imageries or aerial photographs.

Aerial photography for urban change detection and analysis can be facilitated through the use of zoom transfer scope or a video/graphic system as an aid to comparing photographs of two different dates with a map.

This study intend to take advantage of historical and sequential aerial photographs for Naraguta, Jos area as an analytical tool to carry out change detection in the area.

1.2 PROBLEM STATEMENT

Cities are centres of attraction of large population concentration and tend to increase due to specialization and diversification of industrial and other urban economic activities. The discovery of and mining of Tin in Jos was the initial pull-factor for its population concentration. This, has since been overtaken by events.

The decline in the Tin mining saw to the diversification of the economy of the people but it's physical effect have always been obvious on the land scope. For many years, Jos Plateau has witnessed incessant soil degradation by mining operations. Mine failing,

neglected excavations, unfilled sample pits and hill of mine spoilt have dotted the scenery of a once rich and beautiful region.

Naraguta is part of the metropolitan Jos but located at the periphery. The position of the topography of the area in relation to other parts of the metropolis have contributed to its influx of population and hence, activities. This, in consequence therefore, contributed to the social, economic and physical problems of the Jos North. The problem existing today include over congestion of population and activities, competition for space, unplanned and haphazard urban development, environmental pollution, deterioration of social amenities.

So far research information is lacking on urban change detection using the remote sensing techniques on Naraguta area of Jos. There is therefore the need for such researches on urban change detection to know what changes occurred in the area between 1975 and 1991. Where urban changes has occurred most frequently there is a great need to determine access and detect how much change has occurred from time to time. To achieved this, the use of remote sensing application techniques can provide a synoptic, timely, consistent, regular and reliable source of data for urban planners and managers and can determine how improvement can be executed.

1.3. AIM AND OBJECTIVES OF THE STUDY

The study aims at using manual techniques to analysis urban changes over the period 1975 to 1991 with a view to understand the nature and extent of the changes in urban parameters of Naraguta area of Jos.

- (i) To assess and detect how much change has occurred in Naraguta area of Jos Plateau.
- (ii) To determine the implications of this changes on the urban area.

- (iii) To produce urban change inventory maps of the study area for the years 1975 and 1991.
- (iv) To suggest how improvement can be focused towards more positive direction and to serve as a guide for future policy decisions.
- (v) To make recommendations from the use of remote sensing techniques in modern urban planning growth.

4. JUSTIFICATION FOR THE USE OF REMOTE SENSING TO THE STUDY.

One of the main aims of urban change detection is to determine how much development has occurred in an area after a long period of time, the impact of the development and solution to the existing problems and also the provision of accurate information gathering system.

Inventory of urban changes within specific number of years is very important if any meaningful development process is to occur. Therefore the use of remote sensing technique is very advantageous in the sense that real time spatial analysis of an area can be done using either aerial photographs or satellite imageries or both. There is no doubt that satellite imageries as a remote sensing tool has advantages over aerial photograph.

Aerial photography is however, one of the oldest and most common techniques of remote sensing used even today. It is economical, more readily available and has good spatial resolution with good stereoscopic capability. Vertical aerial photograph tend to have properties that are similar to those of map with an approximately constant scale over the whole photograph. Its result could readily be used for mapping and measurement of urban areas and hence, it is very good for studies in urban change detection.

Due to the above and also the non availability and access to the satellite imageries of the area, this study is relying on the manual technique of aerial photographs of 1975 and 1991

to detect urban changes in Naraguta area of Jos. This study also tend to fill the research gap between 1975 and 1991 for there might not have been a study of change detection in the area over the period. The result of this study will provide a knowledge of the extent of urban changes in the area and a basis upon which further research can be conducted in urban change detection using satellite imageries

1. 5. DESCRIPTION OF THE STUDY AREA.

1. 5. 1. LOCATION AND EXTENTS

Jos plateau lies in the central part of the Nigeria and covers a total surface area of about 7, 762 km² (Awogbade, 1983). The plateau is located at latitudes 10° 11' N and 8° 55'N by longitudes 8°21' E and 9° 30' E (fig 1.). The study area; Naraguta (fig 1) lies to the northern - part of the Jos metropolis and it covers total land area of about 25, 536h.a




Fig. 2. PLATEAU STATE SHOWING NARAGUTA, JOS;
THE STUDY AREA.

Source After Okhimamhe A (2000) with modifications.

5.2. GEOLOGY

Geologically, the Jos plateau is representative of pre-Cambrian basement complex; the ancient group of crystalline rocks which form part of the main African continental mass (Buchanan, 1995). The crystalline shield or rocks are of two main categories: metamorphic and igneous (John, et al 1977). However, after the period of continuous erosion, volcanic activities in the mid-tertiary period resulted in vast deposit of basaltic materials, filling alleys and burying tin-bearing alluvium. Naraguta area of Jos lies to the North – West of the Shere hills.

The study area Naraguta area of Jos is not geological unit for it cannot be demarcated on a distinct geological criteria it is purely a morphological or relief unit with elevation above the ground level as a feature distinguishing it from the rest of the crystalline shields.

5.3. CLIMATE

The Jos plateau is one of the coolest regions in Nigeria. The region is characterized by four to five months of dry season and seven to eight months of wet season. The dry and wet seasons in the area as in other parts of the country are greatly controlled by the annual migration of Inter Tropical Zone of Convergence (I T Z C). In the Jos plateau the influence of the northeast trade wind is strong particularly during the months of November and March when the harmattan; a strong dry and dusty wind prevails.

The dry tropical continental air mass arrives the country from the north till October and by January, its effect is felt in almost every part of the country as the dry and cold harmattan season. It retreats in March and by July it s completely out of the country.

The dry season is characterized by the dust laden hamattan wind coming from across the Sahara desert. This occur between November to March to area where relative humidity is low, night temperature is cool, vegetation growth decreases to minimum, the soil dries out.

The wet season set in by the month of May and lasts till September. It is characterized by frequent storms, which coincides with the planting season. During this period the moist south-westerly wind brings rain.

1.5.3.1 RAINFALL

Relief is a significant factor in the rainfall pattern of Jos it has a total of eight months of rainfall from April to November. The Jos plateau has a higher annual total of rainfall than the average for its latitude. It is also termed as a monsoon area due to its heavy rainfall. Orographic effects are also obvious in the rainfall pattern and type. The study area has a total rainfall of between 1520mm to 2030mm per annum, which is more than other areas on the same latitude. Jos is characteristics of a line squall area. It is noted that in spite of its distance from the coast, the study area has fairly high possibility of more than 24% of rainfall occurring for hours during the day in raining season.

1.5.4. LANDUSE

Urbanisation and modernisation processes with the increased demand they pose for land, housing and transportation can effect significant changes in the settlement structure. Factors affecting this change could be attributed to changes in the land use pattern as well as changes associated with socio - economic development. These changes in socio - economic development brings about high expectations; change in taste and higher standards of living. Improvement in transport and communication facilities plays a crucial role in the organisations and changes of urban areas. Since the urban area is an area of transition, there is therefore a conflict between agricultural, residential, commercial and transportation uses of land.

In the study area, high-rise buildings are found in the midst of bungalow and substandard structures. These residential types lie side by side with other land uses such as commercial, institutional and transportation routes. Other built-up areas, clustered residential lands transitional land, areas of commerce, park or open spaces and refuse dumps. The result is that of a mixture of urban landuses and refuse dumps. Industries and factories in few instances are found intermixed with other landuse types in spite of the fact that specific areas are designated for industries and factories. Agricultural landuse occur at places that are quite distant from the residential areas

From the above characteristics of the landuse pattern of the study area, it is evident that planning problems abound and these include, uneconomic use and waste of land, lack of basic data for planning purposes and lack of planning by the local authorities for urban growth. This in most cases result in productive land lying idle for several years awaiting development or being destroyed as sewage or refuse dumps including industrial waste.

1.5.5 SOIL

The high intensity of microbial activities due to the climate of Jos has led to an abundance of humus and thus a high fertility rate of the soils. The soils of the study area range from dark to light brown in colour. Despite this fact, during the rainy season, the high altitude of the area has led to large-scale erosion, leaching and laterite formation. Plants with shallow roots do well during the dry season and these include Lettuce, cabbages, garden eggs and vegetables.

1.5.6 VEGETATION

The study area can be termed as Savannah region. This is because of the type of climate, rainfall, relative humidity and temperature ranges, which encouraged the growth of grasses. Though some trees grow well in the study area in the past, but the activities of man and animals have destroyed much of the natural vegetation. Presently, the existing trees and grasses seen are only survivals of the original plant cover.

1.7 STRUTURE OF THE PROJECT

This project work is divided into five parts or Chapters. Chapter One titled, "introduction" presents the introduction of the project topic, statement of research problem, aim and Objective, Justification for the use of remote sensing technique to the study, description of the study area and the structure of the thesis itself.

Chapter Two titled "Literature review" deals extensively with the relevant and related literatures consulted in the study.

Chapter three titled "Research materials and Methodology" comprises the materials used, principle of interpretation, types of interpretation; Aerial photograph interpretation and growth.

Chapter four titled "Discussion of result" presents analysis, discussion and results

Chapter five titled "Summary, Conclusion and Recommendations" of the research

CHAPTER TWO

2.0 LITERATURE REVIEW

2.1 A number of studies have shown that landuse mapping for various purpose and at widely varying degrees of detail is possible through remote sensing techniques. Assessing overall landuses on small-scale imagery first because popular with the advent of astronaut photography and more recently, land sat imagery.

The urban land is always in a dynamic state, thus to keep tracks of development requires a real-time system of analysis. In examining the development of urban areas, the conventional methods are not flexible in monitoring the trends in the growth of urban environment, hence this research advocates the uses of remote sensing techniques.

Research into the application of aerial photography and other remotely sensed imageries to urban studies dates perhaps from the late forties (Adeniyi, 1984) in developed countries some development have been made in methods of estimating dwelling units (Eyre,1970); Lindgren, 1971 and Hsu, 1971) estimation of population density (Nkambwe, 1983; Adeniyi, 1984), the estimation of characteristics of residential buildings (Adeniyi, 1984) to the estimation of building construction as indication of spatial urban growth (Ajayi 1981).

Dunfard,(1980) illustrated the effectiveness of remote sensing for rural development planning in Africa. The major aim of their study was to attempt to minimize the time, cost and personnel using a combination of remote sensing techniques for rural development planning. Using aerial photography and existing reports on vegetation and

soil conditions, they were able to provide land resources and landuse information for rural development planning in Arusha region, Tanzania for a period of three months.

Rao, (1994) used remote sensing data (aerial photography and satellite imageries) and information obtained from the Census department in India. He also identified the causal variable and using three techniques: Causal modeling techniques, cross-classification method and rating method, he was able to forecast the future quantity of built up areas.

Sogunle, (1990) studied the landuse/land cover of Ibadan area by SPOT image of 1986 with aerial photographs taken in 1993: A six category landuse scheme was adopted for use in the interpretation. The SPOT image was manually interpreted at a scale of 1:50 000 using prom-2 projector. Identification of landuse/land cover type was based upon characteristics of the signature observed.

The results showed that the city has witnessed a tremendous urban expansion largely concentrated along the central strip of the study area where urban landuse increase by 173% from 8.33km² in 1973 to 22.74km² in 1986. The urban expansion has serious consequences on the soil in terms of exposure to hazardous torrential rainfall, severe erosion and run-off, high soil compaction due to excess human traffic and surface sealing. This study is applicable to this present research work because one of the objectives is to assess and detect how much change has occurred in Naraguta area of Jos.

Adeniyi, (1980) used a combination of computer and sequential aerial photographs of 1962 and 1974 at a nominal scale of 1:40 000 and 1:20 000 respectively to detect changes in landuse of Lagos. The study area encompassed urban built up areas,

urban vacant land and non-urban land. A landuse classification scheme was devised as well as a minimum mapping unit for interpretation and subsequent storage of landuse data by computer. The result shows that the landuse change revealed the rapid increase of residential landuse and strong expansion of the urban area of Lagos. This study is also relevant to this research for one of the objectives is to detect urban change in the study area.

Lo, (1972) also used sequential aerial photographs at a nominal scale of 1:20 000 to study the landuse changes in Clarke county, Georgia between 1970 and 1983. By using a simple technique, he matched two landuse maps compiled for two different dates over a light table to delineate landuse changes the resultant map shows an increase in the urban or built up land particularly in residential, commercial and services as well as a decrease in the agricultural land and forest. This study is also relevant to this present research for one of the objectives is to detect urban changes and also to provide a map showing the changes.

Adeniyi, (1981) used sequential aerial photographs for 1974 and 1976 at a nominal scale of 1:20 000 and 1:25 000 respectively using simple visual method to derive timely and quantitative data on the rate of building construction in a rapidly urbanizing areas of Lagos. Agege, one of the urban fringe areas was used as a case of study. Grids of one hectares in size were carefully drawn on stable transparent film and super imposed on the right hand photo of a pair of stereo-photographs and then reproduced photographically prior to interpretation. Two interpreters using a mirror stereoscopes were used to record data on each of the cells. Such as the total number of completed

buildings under construction, total number of completed buildings and the specific uses of cells without any structure.

The result shows an increase in the rate of building construction of 37.1% between 1974 and 1976. It also shows the speed at which such type of data could be obtained over the conventional methods. This study is also relevant to this research for it shows that there are some changes in the urban landuse in the form of building constructions both completed and uncompleted when compared with the previous aerial photographs of the study area.

Nkambwe, (1984) used multi-date aerial photographs for the study of physical urban growth of Ile-ife. Sequential aerial photographs of 1950, 1960 and 1970 at a scale of 1:3 600, 1:9 600 and 1:10 000 respectively were used to obtain data on the extent of the build up area of the town total floor area above the ground and the total floor area on the ground for the Study periods. The data was obtained by a combination of the three sets of aerial photographs with the aid of a mirror stereoscope. The result shows three forms of physical growth in the study area, these are, the lateral growth the vertical growth and, the intensification or infilling of spaces between buildings with buildings. This study is also relevant to this research for it how changes are detected in the study area in form of urban growth.

Ikhuoria, (1993) applied remote sensing techniques of sequential black and white aerial photographs to study the vegetation and landuse changes in a rainforest ecosystem of northeast Edo State.

The result shows that within a decades (1967 and 1977) the ecosystem experienced a drastic reduction or depletion of forested lands rapid territorial expansion of settlements and tremendous increase in agricultural lands.

Okhimamhe, (1993) also carried out a study using SPOT HRV acquired in 1986. A combination of Satellite imageries with aerial photographs of 1974 was used to detect the changes in the landuse/landcover in the Burum/Tiga area over a period of 1 years of after the construction of Tiga dam. The result shows that 38,897 hectares of changes have taken place where crop/pasture land, wooded shrub land had increased by 104 percent. The study also showed that the sandy area has increased which are indicators of desertification.

Roger, (1985) carried a study to assess the change in the land surface occurring as result of the activities of man or climatic variability. Land sat MSS imageries of 1970 and 1979 covering St. Lawrence Valley in Quebec Canada were used in a time series analysis to monitor seasonal and long term changes in the landuse and land cover. the result showed that deforestation occurred and there was significant decrease in farmlands and a marked increase in urbanisation

Avery, (1980) utilized aerial photographs at 1:20 000 scale in 1954 and 1960 to evaluate landuse changes in Georgia, U.S.A. For each period, a landuse map was determined with the aid of grid. The study discovered a shift of the agricultural pattern of the dark country from heavy emphasis on cotton to poultry production, Livestock farm wood lot management while crop land was reverted back to forest land as result of the influx of manufacturing industry to provide employment.

Lo and Shipman, (1990) carried out a study on Geographical information system (G.I.S) approach to landuse change dynamics detection. This was applied to assess the impact of new town development in Twen Mon, New territories, Hong Kong through integrating of past and current aerial photographs which were taken in 1976 and 1987 at a scale of 1:25 000 and 1:40 000 respectively image overlaying and masking techniques were used. A low-cost microcomputer based GIS IDRISI soft ware was used in landuse change detection. The result shows that the binary masking method reveals the dynamics of landuse change.

Owolabi (1998) used sequential aerial photographs of 1982 and 1988 at a nominal scale of 1:10 000 and 1:4 000 respectively to detect changes in urban landuse of Victoria Inland, Lagos over period of 16years. Using a binary masking techniques and over laying of the two aerial photo mosaic with a manual interpretation technique he was able map a total land area of 268.0ha of the area. The result shows that about 102.64ha (38.3% of total area covered) was the changes discovered over the period 1982 to 1988. He identify the causes and consequence of the urban growth of the area.

Ojigi (1997) used a combination of multi-temporal SPOT satellite imageries (Panchromatic) and aerial photographs (mosaic) of 1980, 1984, 1988,1992, and 1994 to manually produce the sequential urban growth maps of Abuja and identified the growth rate of five districts of the Federal Capital Territory using urbanization curves and histograms. The result based on graphical and multiple regression analysis revealed that Abuja has growth substantially over the years and that urban development and growth is a function of financing

Lo and Dale (2000) used a combination of Landsat TM and sequential aerial photographic imageries for 1973, 1979, 1983, 1987, 1992 and 1997 with computer to detect a dramatic changes in landuse and landcover of Atlanta metropolitan area of Georgia, USA. The

result shows a dramatic change in landuse and landcover with loss forest and cropland to urban use. In particular, low density urban use, which include largely residential use, has increased by over 119% between 1973 and 1997. These landuse and landcover changes have drastically altered the land surface characteristic of the area.

From the foregoing review of literature, it is discovered that some of the techniques applied by various studies are also applicable to this present research. The attempt made in this study is to take advantage of the advance in remote sensing application technology as shown in the case studies described in the review to further enhance our knowledge on urban change detection in Naraguta area of Jos.

The application of remote sensing techniques in monitoring, evaluating and detecting changes in urban landuse in Nigeria is still at its embryonic stage. Problems of remote sensing applications in developing countries including Nigeria are; Manpower shortage, high cost of product acquisition, inadequate finance and lack of institutional support on the part of the government.

CHAPTER THREE

3.0 RESEARCH METHODOLOGY AND MATERIAL

3.1 THE RESEARCH MATERIALS USED

Aerial photographs are one of the best sources of data for landuse change detection in urban areas at different time intervals, This is due to the advantages which they offer over other sources such as the on-ground observation.

Two sets of aerial photographs of the study area acquired from the photogrammetry department of the Ministry of Lands and Survey, Jos was used. Those of 1975 have a nominal scale of 1:10 000 while those of 1991 have a nominal scale of 1:8 000. The 1975 sets were numbered as follows; 76626-109,76626-110 and, 76626-111. The 1991 sets were numbered as follows; 8748,8749 and 8750.

The quality of the 1975 sets of aerial photographs was fair, the problem was that they were exceptionally whitish (bright) and thus, there was some difficulties in delineation and identification of features. The 1991 sets of aerial photographs on the other hand were of good quality and thus, delineation and identification was much easier.

Other materials used include; mirror stereoscope, graph sheets, light table, cartographic (rotrin) pens and lettering stencils. Also complementing these aerial photographs were still pictures of some scenes (urban features) of the study area and fieldwork.

1.2 PRINCIPLES OF INTERPRETATION

The characteristics used in analyzing and interpreting remote sensing data had helped in recognizing, differentiating and qualifying objects on the aerial photographs. Using such elements as size, tone, texture, locational site, shape and shadows were so useful for the interpretation.

The sizes of buildings were used in determining the type of building. For example, in differentiating between schools, and residential houses. The texture; roughness and smoothness of objects were used to differentiate rock out crops from a bare soil and among other features such as racecourse open forest.

The pattern which refers to the arrangement of structure were used in differentiating university lecture halls from student hostels and staff quarters more accurately because of familiarity with the area.

The location of objects or structures in relation to others had also helped to ascertain many relevant features. Association of elements was very important here because the areas with residential quarters were never associated with cultivation.

Shades casted by features such as staff quarters, lecture halls, residential compounds, student hostels, mosques and churches due to their sizes have also assisted in the interpretation. The shades casted by trees had helped in differentiating them from open ground or rock out crops.

3.3 METHODOLOGY

Aerial photography have been employed by planners to detect changes over a period of time in a region. The advantage of aerial photographs is that when it is taken as overlapping pairs it can provide a three-dimensional view (stereoscope view) of the earth's surface.

In this study, each set of photographs were sequentially arranged on table according to their fiducial points and flight line to overlap each other by 60%. This gave a stereoscopic view of the features contained in the pair of photographs. These arrangements made possible the production of mosaic of the study area for the two periods under study. The mosaic for each of the years (1975 and 1991) were traced from maps into an acetate transparent sheet and onto a tracing paper using a light table, pencils and rotting pens.

The scale of the 1975 aerial photograph (i.e. 1:10 000) is smaller than those of the 1991 sets of aerial photographs (i.e. 1:8 000). For this reason, the researcher had to enlarge the scale of the 1975 aerial photographs Mosaic to correspond to that of the 1991 set before interpretation was done. Square grid enlargement method was used to transfer information from the former 1:10 000 scale mosaic to the new 1:8 000 scale outline. Permanent features such as rock outcrops and even roads were first registered before the rest information.

Using the elements of interpretation, features such as the University, Federal School of forestry racecourse, senior staff quarters, streams, rocks outcrops, vegetation, footpath, roads etc were delineated and mapped from the 1975 mosaic. Similar procedure

was applied in the case of the 1991 mosaic. Conventional pictures of scenes in the study area was used to aid the interpretation

3.4 DATA ANALYSIS

The landuse maps produced from the mosaic of the aerial photograph of 1975 and 1991 were overlaid on the light table. From these overlay of 1975 and 1991 maps an urban change map was then produced to detect and signify the changes that occurred in the study area between 1975 and 1991 (sixteen year period) Before the mapping was done, the changes detected were field checked ground truth.

Quantitative aerial data analysing the urban change categories were then compiled for the study area. This was achieved by placing a transparent square grid of 5mm x 5mm cell over the map to measure areas of urban changes.

3.5 CLASSIFICATION SCHEME

Landuse interpretation accuracy depends on the classification scheme, which had to be designed with regard to the cultural character of the study area. However, the USGS scheme designed by Anderson et al (1976) has provided a valuable guide for the classification scheme used in this research. The concept of different levels of details of urban landuse are interpretable according to the different scales of the imagery and hence, the spatial resolution quality is normally adopted by the researcher.

However, each of the urban landuse categories was identified by a two digit symbol indicating the level II category interpreted from the aerial photographs of 1975 and 1991 respectively. The classification scheme used for this visual manual interpretation is given in table below,

Table.3 urban landuse classification scheme

S/N	LEVEL I	LEVEL II
1.	Built up Areas	1.1 University of Jos 1.2 Federal School forestry 1.3 Commercial Areas 1.4 Residential Areas 1.5 Industrial Areas 1.6 Senior Staff Quarters 1.7 Parks 1.8 Race course
2.	Underdeveloped Areas	2.1 Bares course 2.2 Rock outcrops

Source:- Compiled by the author (2003)

3.6 GROUND TRUTH

During the course of the study, field checks were conducted first to familiarize the researcher with the study area. During the course of interpretation of the aerial photographs, field check was also carried out several time to identify and verify the features seen on the aerial photographs so as to minimise interpretation errors. All the areas involved in the study visited were photographed. The areas visited included packs, the Federal school of Forestry, Jos, the University of Jos, Racecourse and other areas of interest to the study area. All these areas were considered as samples and represent the rate and type of change in the recent years.

CHAPTER FOUR

4.0 RESULTS AND INTRODUCTION

In this chapter, the results of the interpretation of the aerial photograph of 1975 and 1991 of Naraguta area of Jos Plateau are shown in tables and figures. Table 4.1 shows the urban categories of the study area and their percentage aerial coverage as interpreted from 1975 and 1991 air photo mosaic. Figure 4.1 shows the urban map of Naraguta as interpreted from 1975 air photo mosaic while figure 4.2 shows graphically its percentage distribution. Figure 4.3 shows urban map of the area as interpreted from the 1991 air photo mosaic while figure 4.4 shows graphically the percentage distribution of each of the urban categories of the area. Table 4.2 shows the distribution of urban categories in hectares (ha) for 1975 and 1991 and the magnitude of change for each of the categories over the period. Table 4.4 shows in proportion the changes for each of the urban category class to the overall change. Figure 4.5 shows the urban change map of the study area as interpreted from the overlays of the 1975 and 1999 air photo mosaic while figure 4.6 graphically compared the relative distribution of each category for the two period of study.

4.1 INTERPRETATION OF 1975 AERIAL PHOTOGRAPH

The interpreted map of the study are which was obtained directly from the air photo mosaic by the author is shown in figure 4.1a. The results of the interpretation showed that the under developed areas made up the largest category in the classification covering a total area of 14000 hectares representing about 60.34% (Table 4.1). The interpreted map showed that there was no Motor Park and so it is represented by 0% (Table 4.1). The only major road (Bauchi road) was quite narrow. It was also noted that the Senior staff Quarters was quite small

occupying an area of about 1500 hectares representing about 6.46% (Table 4.1) of the study area. The racecourse can be clearly identified at the northern tip of the map (figure 4.1) occupying an area of about 9.05% (Table 4.1). This can be associated with the presence of expatriates who used it for recreational purpose.

The University of Jos is found at the North Central part of the map (figure 4.1) to the left hand side of the Bauchi road. It covered an area of about 900 hectares representing about 3.87%. The federal School of Forestry Jos is found directly opposite the University (fig. 4.1) Covered an area of about 700 hectares representing about 3.0% (Table 4.1)

Table 4.1 Urban Categories Aerial Coverage and percentage 1975 and 1991.

S/N	Urban Category	1975		1991	
		Area (Hectares)	Percentage (%)	Area (Hectares)	Percentage (%)
1.	University of Jos	900	3.87	1792	7.0
2.	Federal School of Forestry	700	3.0	1536	6.0
3.	Race Course	2100	9.05	1088	4.26
4.	Senior Staff Quarters	1500	6.46	3264	12.78
5.	Parks	*	*	384	1.5
6.	Built up Areas (Developed)	4000	17.24	10880	42.60
7.	Under Developed Areas	14000	60.34	6592	25.8
Total		23200	100	25536	100


Source:- Author Analysis of Aerial Photographs of 1975 and 1991 of Naraguta area of Jos

In general, the built up areas accounted for about 4000 hectares of land of the study area and representing about 17.24% (Table 4.1). A stream which cut across the Bauchi road flowing from Dogon-Dutse area down to Anguwan-rogo to the North west of the map (fig. 4.1) had lots of vegetation cover along its banks. There were neither farms nor human settlement at the

western side of the university. The houses seen on the aerial photographs especially those found in the Senior Staff Quarters and Dogon-Dutse area were well spaced out and arranged in rows due to low demand for land.

The road network is not well developed and mostly narrow and untarred. The occurrence of footpath in the map (fig. 4.1) as seen on the aerial photograph indicated the stage or level of development in the study area. People mostly moved on foot and on animal backs. Most of the trees shown on the photographs were well arranged and demarcates buildings. This suggested that there is Government Reservation Area (GRA) in the area. The drainage pattern of the area can be clearly identified with its dendrite pattern taken their sources from the hills around the northeastern part of the map (fig 4.1).

The build-up areas have also expanded greatly. The building structures have been modernized; an indication of rises in living standard and taste of the people. The build up areas though scattered about the map (fig 4.1) accounted for 10880 hectares representing about 42.6% (Table 4.1), while the underdeveloped areas had drastically reduced in size due to encroachment by buildings and farmlands. This category (under developed Areas) accounted for about 6592 hectares presenting about 25.8% (Table 4.1) of the land categories in the study area (fig. 4.1a). The expressway to Bauchi State (Bauchi ring road) had also been constructed (plate 3c) Large number of motor vehicles were noted on the aerial photograph plying the ring road. This suggested rapid development in transportation in the area. The drainage network can hardly be noticed due to blockage by building structures in the area.



Jos University (1) Federal School of Forestry (2) Race Course (3) Senior Staff Quarters (4)
 U.D.A) Under Developed Areas: Rock Out Crops/Hills Vegetation
 Streams — Major Roads — Minor Roads - - - Foot Path

4.1: URBAN MAP OF NARAGUTA, JOS (1975),

Source Author's Analysis of Air Photographs (1975)

NARAGUTA JOS
1975




Fig 4.1b Percentage distribution of urban categories 1975

Source:- Author Analysis of Aerial Photography of Naraguta Jos (2003)

The Racecourse had also shrunk in size and lacks maintenance. Building structures and some farming activities had encroached on the racecourse (plate 1b). It now only account for 1088 hectares representing about 4.26% (Table 4.1) of the land categories in the map (fig 4.1) of the study area. The graphical representation is seen in Figure 4.2

2 INTERPRETATION OF 1991 AERIAL PHOTOGRAPH

The 1991 Aerial photo mosaic (fig 4.3) shows a great contrast of what was the study area in 1975. A lot of developments have taken place over the sixteen-year period. Petrol Stations now lined up the boundary of Bauchi road, which also witness expansion with even streetlights. Other roads that led into the major part of the town have also been expanded and tarred. The stream that cut across the Bauchi road had shrunk to less than half its original size and there was an absence of vegetation on either side of its bank. This was because refuse have been continuously dumped in most parts of the river banks while at other areas of the river banks structure had been erected there.

A new motor park have been constructed (Bauchi Park) covering an area of about 384 hectares representing about 1.5% (Table 4.1) of the land categories in the map (fig 4.3). The Jos University had also witnessed expansion and it now accounted for about 1792 hectares representing about 7% (Table 4.1) of the land categories in the map (fig. 4.3). The Senior Staff Quarters of the University located at the southeastern part of the map (fig. 4.3) also expanded and it now covered an area of about 3264 hectares representing about 12.78% (Table 4.1) of the land categories in the study area. The graphical representation is seen in fig. 4.4.




Fig.4.3 URBAN MAP OF NARAGUTA, JOS (1991)

Source:- Author's Analysis of Air Photographs of (1991)

NARAGUTA JOS 1991




Fig 4.4 Percentage distribution of urban categories 1991

Source:- Author Analysis of Aerial Photography of Naraguta, Jos

4.3 URBAN LANDUSE CHANGE BETWEEN 1975 – 1991

In this study, urban change refers to the reduction or increase of the various urban categories used in the classification scheme (table 3).

Table 4.1 shows the aerial coverage and percentages of the categories of the urban landuse of the study area. Table 4.2 shows the distribution of urban changes based on each category and the magnitude of changes between the two period, 1975 and 1991. This makes urban change analysis much easier. Fig 4.6 shows the graphical representation of the relative distribution of urban change categories over the two periods.

Table 4.3 shows the percentage changes in the distribution of the urban categories in the study area over the two period (1975 –1991). As can be seen in Table 4.3, the underdeveloped areas recorded the highest amount of change from 60.34% to 25.8% and a decrease of –34.54%. The Built up areas increased from 17.24% in 1975 to 42.6% in 1991. This therefore shows an indication of an upward rise in terms of development in the study area. More lands have been used up, building expanded and structures had been erected including the expansion in road networks. These changes are shown in the urban change map (fig 4.5) of the study area.

The Jos University also recorded a positive change from 3.87% in 1975 to 7.0% in 1991 and a percentage increase of 3.13% (table 4.3). The racecourse on the other hand recorded a negative change from 9.05% in 1975 to 4.26% in 1991 and decrease of –4.79% (table 4.3). This can be attributed to the increase demand for building and other forms of urban activities. The Senior Staff Quarters have on the Other hand recorded as positive change from 6.46% in 1975 to 12.78% in 1991; an increase of 6.32% (Table 4.3). The rise in the figures is the result of increase in student's population and hence, the population of the Staff of the University due to the expansion witnessed in the institution over the sixteen year period. The park (Motor) on the

ther hand did not appear in the 1975 urban map because it was non-existent then in the study area (fig 4.1) but it appeared in the 1991 urban map (fig. 4.3). This shows the development that was witnessed over the sixteen-year period after 1975. Therefore the percentage rose from nil or % in 1975 to 1.5% in 1991 (Table 4.2). The Federal School of Forestry also recorded an increase changes from 3.0% in 1975 to 6.0% in 1991 and increase of 3.0% (Table 4.2).

Thus, the Positive and Negative changes noted in the study area between 1975 and 1991 might be the result of population increase, increase in economic and other activities and hence, the rise in living standard of the people of the study area and consequently the demand for land.

Table 4.2 Distribution of Urban categories and magnitude of change between 1975 - 1991.

S/N	Urban Category	1975 Area (Hectares) (A)	1991 Area (Hectares) (B)	Magnitude of change in hectares (b - a) = c	Remarks
1.	University of Jos	900	1792	892	Increase
2.	Federal School of Forestry	700	1536	836	Increase
3.	Race Course	2100	1088	- 1012	Reduction
4.	Senior Staff Quarters	1500	3264	1764	Increase
5.	Parks	*	384	384	Increase
6.	Built up Areas (Developed)	4000	10880	6880	Increase
7.	Under Developed Areas	14000	6592	- 7408	Reduction
Total		23200	25536	19.76	Increase

Note * Urban Change Categories did not appear on the 1975 Aerial Photo.

Source:- Author Analysis of Aerial Photographs of 1975 and 1991 of Naraguta area of Jos
(2003)

Table 4.3 Percentage Distribution of the Urban Categories.

S/N	(A) Urban Categories	(B) 1975 Percentage Distribution	(C) 1991 Percentage Distribution (%)	(D) Percentage Change (C-B) %
1.	University of Jos	3.87	7.0	3.13
2.	Federal School of Forestry	3.0	6.0	3
3.	Race Course	9.05	4.26	-4.79
4.	Senior Staff Quarters	6.46	12.78	6.32
5.	Parks	-	1.5	1.5
6.	Built up Areas (Developed)	17.24	42.60	25.36
7.	Under Developed Areas	60.34	25.8	-34.54
Total		100	100	-

Source:- Authors Analysis of Aerial photographs of, Naraguta area of Jos (2003).

Table 4.4 Proportion of change of Each Urban Category Class to the Overall Change

S/N	Urban Category	Magnitude of Change (C) (Ha)	Change % of Each Class (C/Total C x 100)
1.	University of Jos	892	4.65
2.	Federal School of Forestry	836	4.35
3.	Race Course	- 1012	5.27
4.	Senior Staff Quarters	1764	9.19
5.	Parks	384	2.00
6.	Built up Areas (Developed)	6880	35.87
7.	Under Developed Areas	- 7408	38.63
Total		19176	100

Sources:- Authors Analysis of Aerial photographs of Naraguta area of Jos (2003)

NARAGUTA, JOS (1975 - 1991)




Fig 4.6 Relative distribution of urban categories (1975 –1991)

Source:- Author Analysis of Aerial Photography of Naraguta, Jos (2003)

4.4 DISCUSSION OF RESULTS

Naraguta which was once quite scenic, sparsely peopled and more or less occupied by expatriate who came as a result of mining and other related activities in the area has now change significantly and it is witnessing urban growth. This growth is mostly attributed to the result of influx of migrants form within and without the Jos Plateau. This have a far reaching implications on its environment as well as the resources within.

The study have demonstrated the usefulness of remote sensing applications in analyzing the applicability of aerial photographs in detecting urban changes in the area. From the analyses of the air photo mosaics of 1975 and 1991 it was discovered that so many developments have taken place as can be seen below.

The 1975 air photo mosaic (fig 4.1) revealed that there was no Motor park in the area and hence, a nil was recorded as its aerial coverage and its percentage distribution (tables 4.1, 4.2 and 4.3) its present location was occupied by farm lands. The influx of population and hence, the need for accommodation and other activities stimulated the need for additional Motor parks in Jos and hence, its present location along the Bauchi road at the northern tip of the map (fig 4.3) and plate 2a shows some Motor cars parked in the Motor park. The aerial coverage as obtained from the analysis of the 1991 photo mosaic is 384ha. (table 4.2) which is about 1.5% of the total land area (table 4.3) and its percentage change in all the categories is 2.0 (table 4.4). Fig 4.6 graphically compares its relative distribution from the two air photo mosaic (1975 and 1991). Therefore, there is an increase or growth from nil to 2.0% for the Motor park in relation other urban categories.

The racecourse (fig 4.1) that was relatively large in area, well maintain and well serviced as recreation centre for the expatriate mine workers as now become a shadow of its past due to

neglect and consequently the demand for land (fig 4.3 and 4.5). Its aerial coverage in 1975 mosaic is 2100ha (table 4.3) and its percentage distribution relative to other urban categories is 05% (table 4.3) and a percentage change of -4.79%. Plate 1b shows its present state in 2003 while figures 4.3 and 4.5 show its 1991 mosaic. At present, most part of has been encroached by farmlands and residential uses (Plate 1b). Most of recreational facilities have been vandalized and the structure dilapidated.

The underdeveloped area is 14,000 ha in 1975 mosaic and about 60.34% of the total land area (table 4.1) as can be seen figure 4.1 and its graphical representation in figure 4.2. It however, recorded the highest negative changes over the period from 14,000ha (60.34%) in 1975 to 6,592ha (25.8%) in 1991 mosaics (tables 4.1 and 4.3). Its magnitude of change is about -7408ha and a percentage change from overall total of 38.63% (table 4.4). This can be seen in figure 4.3 and 4.5. Figure 4.6 compared its relative distribution for 1975 and 1991. The negative change however have impacts on the reduction in farmlands and hence agricultural productivity. It also reveals the increase in demand for various urban landuse. This however explains the reason why farming activities are carried out on hillslopes and many highlands in the area.

Anguwan Rogo located to the northwestern part of the map (fig. 4.1 and 4.3) which hitherto was well organised and the planned has now turn to shun or squalo area. Its streets were taken over by commercial and industrial user. Forest found along the stream banks (fig. 4.1) have been cleared (fig. 4.3) and was replaced by residential, commercial and industrial landuser (plate 1a and c). The stream have also shrunk considerably due to dumping of garbage and household refuse to the extent that its flow have been reduced to almost a trickle. If this trend is not controlled its however, feared that in a near future environment pollution and its attendant effects and flood might pose a problem to the area. The role of vegetation cover in reducing impacts of

rain drops on soil, encouraging the infiltration of water and preventing soil erosion is not in doubt. Its destruction is bound to encourage surface run off rain water and soil erosion. Since household sewage are continuously dumped into stream channel flood might result which may have far reaching consequences on the people and those categories of landuse located along the stream banks.

The Dogon Duste area (GRA) located to south west (fig.4.1) has become a shadow of its past and is now considered as a twilight zone (fig. 3). Slums or barchas (Zinc houses) have taken over most of the available spaces in the area. These slums are mostly the residences of the retired and serving low cadre mine workers. Also mining and other allied construction firms now compete for space in the area designated for residential uses. This however, is bound to affect the urban plan of the area.

The University of Jos has also recorded a positive change from 900ha or 3.87% in 1975 photo mosaic (fig. 4.1) to 1,792 or 7.0% in the 1991 photo mosaic (fig. 4.3) as is seen in table 4.1 and 4.2. And magnitude of and proportion of change can be seen in table 4.3 and 4.4. The growth rate of 892ha over the period might be attributed to expansion in virtually all the institutions of higher learning in the country to accommodate the increase rate of student enrollment as well as increase in the courses offered and department. This also might influence the increased of about 1764ha (table 4.2) for the senior staff quarters. This is because as the students population and the number of courses and departments increased so also the number of academic and non academic staff to cope with the demand and hence the need for increase in number of their accommodation.

The Federal School of Forestry has also witnessed an increase change from 700ha in 1975 photo mosaic (fig. 4.3) as is seen in table 4.1. The increase change is 836ha (table 4.2).

This can be attributed to the introduction of new courses as well as increase in student enrollment over the period just as is the case with the University of Jos.

The built-up areas witnessed the highest positive change from its 4000ha (fig. 4.1 and table 4.1) in 1975 mosaic to 10880ha (fig. 4.3, 4.6 and table 4.2) an increase of 6880ha (25.36%) as is seen in tables 4.3 and 4.4. The built-up areas can be seen scattered in most part of the 1991 mosaic (fig. 4.3). This growth rate can be attributed to the influx of population due to increasing role of Jos from its hitherto mining centre to its present administrative, commercial, industrial, educational and other functions.

The study also reveals an overall increase in urban land use of the area from a total of 23200ha in 1975 mosaic to 25536ha in 1991 mosaic, thereby indicating a magnitude change of +10,756ha and -8,420ha for both developed areas and underdeveloped areas respectively. This is however, inline with similar studies conducted elsewhere such as those of Owolabi (1998) in Victoria Island; Lagos, Ojigi (1997) in Federal Capital City; Abuja, Adeniyi (1980 and 1981) in Agege; Lagos and Roger, et al (1985) in St. Lawrence valley, Quebec; Canada.

1.5 CONSEQUENCES OF THE CHANGES

We live in world of dynamics of nature, economics, population and urbanization. These has far reaching consequences on man and his environment. Principal threats to man come from nature (his environment) and from man himself. There is a growing awareness and evidences that continuous, unrestrained and unplanned intervention on the natural ecosystem can only lead to disaster. There now a growing fear that rate at which the world's population is growing and the corresponding rate at which the resources are being exploited is becoming a source of concern that man existence, will within the foreseeable future be threatened.

Analysis of the 1991 photo mosaic of Naraguta (fig. 4.3) as compared with that of 1975 photo mosaic indicated that most of changes detected especially in the built-up areas deviated from the initial landuse pattern of the area. As was pointed out in chapter 4.4, there is an encroachment of farmlands and residential uses into the racecourse which hitherto serve as a recreation center. It is also reveal that there is a massive deforestation especially along the stream banks of Anguwan Rogo. There is also a massive expansion of squalor and slums in both Anguwan Rogo and Dogon Duste areas. The study also revealed the encroachment of commercial and industrial uses in the area meant for residential use thereby creating a sort of mixed or haphazard landuse pattern. The ground truth conducted in May, 2003 also reveal the indiscriminate and uncontrolled dumping of household refuse and industrial sewage into the stream channels thereby blocking flow of water and hence leading to environmental pollution and health hazards.

The overall consequences of the above on man and environment in the area can be summed up under the following perspectives;

- i). Traffic congestion: - Due to high population and number of vehicles particularly along the road between the University of Jos, the Federal School of Forest up to the Bauchi road Motor park, traffic congestion is a daily phenomenon. This is mostly severe during the rush-hour periods i.e. in the morning when the student and children are going to the school and workers to their work places and from the early hour of the afternoon up to evening. This situation is most acute at the road junctions where most road users are impatiently eager to reach their destination.
 - ii). Overcrowding:- population pressure create the desire for more lands as well as the development of and expansion of squalors and slums. This is particularly pronounced in areas of lower cadre mine and construction workers. This study revealed the concentration of slums known as "Barchas" in Anguwan Rogo and Dogon Dutse areas. The consequences of this development are the overstressing of the utilities and social amenities in the area. The situation is so acute that electricity and pipe borne water is being rationed in some areas. Government or public health facilities and schools are over populated resulting in massive decline in the services they provide to the citizens.
 - iii) Flood and environmental pollution:- Due to the increasing rate of deforestation, pavement of houses and roads with concrete and tarmac which encourages surface run off and the reduce the rate of percolation of rain water into the soil. The dump of refuse and industrial waste into the stream channels block them and thereby creating an environmental pollution and consequently, flood during the period of heavy rainfall.
- The foregoing situations therefore calls for the urgent attention of the policy makers and in particular, the urban and town planning authority to control the trend.

CHAPTER FIVE

5.0 SUMMARY, CONCLUSION AND RECOMMENDATIONS

5.1 SUMMARY

To understand and develop the present landscape, we need to discover why town and cities in Nigeria have grown to their present pattern. This study demonstrated the applicability of aerial photographs in detecting landuse changes in Naraguta area of Jos Plateau from 1975 to 1991. The research focused on analyzing the nature and extent of changes in the urban parameters of the area. To achieve the objective, manual technique was used to interpret the aerial photograph of the area for the two periods (1975 and 1991). This was achieved by mapping the urban extent of the area for each category of landuse types for the two periods. The landuse maps generated from the air photo mosaic of the two periods were overlaid and then a change map was produced and the changes detected were calculated. The USGS landuse/landcover classification scheme of Anderson, et al (1976) has provided a valuable guide for designing the landuse classification scheme for this study. This scheme provides for urban landuse details (level II) interpretable according to different scales of the photographs and hence, the spatial resolution quality is adopted.

The results obtained have revealed that the University of Jos has increased by 892ha, Federal School of Forestry by 836ha, the senior staffs by 1,764ha, the Motor park by 384ha and built-up (developed) areas by 6,880ha. The underdeveloped areas and racecourse shows a decline in both aerial coverage and is negative magnitude of change of 7408ha and - 1012ha respectively. Table 4.3 and 4.4 shows the percentage distribution of the urban categories and the proportion of change of each urban category class to the overall change over the 1975 and 1991 period (16years).

5.2 CONCLUSION

Landuse maps derived of change detection derived from aerial photography data allows us to compile an objective inventory of an entire area before proceeding to a detailed study phase. By comparing photographs acquired on different dates or recent imageries with older maps, we can monitor growth rate and detect every change that occur. It is our job as environmental scientists to examine when growth should be allowed to take place and the land are required. We also need to know what resources are threatened and the likely consequences of the planned growth.

In line with the stated objectives, this study has shown that there is a significant changes in the urban landuse of Naraguta area of Jos Plateau over a sixteen year period. That the build up areas increase from 7100ha in 1975 to 17856ha in 1991 showing a positive increase or changes while the underdeveloped areas including the race course show a negative change or decrease growth from 16100ha in 1975 to 7680ha in 1991.

From the foregoing therefore, it is concluded that aerial photographs as a tool of remote sensing can help in providing adequate information necessary for detecting urban changes in landuse and can also be used in monitoring and studying the landuse parameters of on area.

5.3 RECOMMENDATIONS

From the research conducted and the findings made so far, the following recommendations are made;

- i. That there is an urgent need for the policy makers, the town planning authority as well as all the parties concerned to try as much as possible to adhere to the master plan of the area for the purpose of development control.

- ii. There is the need for the State Sanitary and environmental board to monitor and ensure that house-hold refuse and industrial waste are disposed of at the right places to avoid pollution and flood disaster.
- iii. There is also the need for the town planning authority to properly implement the zoning system of land use to avoid conflicting uses of the land as the trend shows in some areas.
- iv. There is also the need to separate commercial/Industrial uses of the land from the residential uses.
- v. There is the need for the town planning authority to work hand in hand with the National Remote sensing center, in Bukuru, Jos to obtain up-to-date and real time data on satellite imageries of the Jos Plateau so that different aspects of the environment can be monitored and studied.
- vi. The launching of the Nigeria satellite 1 in September 2003 is a welcome development and its image should be acquired at regular interval by the state planning authority so that it can facilitate its activities.
- vii. Staff of the state town planning authority should be trained on the current remote sensing techniques of interpretation of imageries and digital image processing techniques to facilitate its activities.

To this end, it can thus be concluded that the use of remote sensing to not only urban applications is inevitable in this epoch of technological advancement. Further research is recommended on the application of satellite imageries in urban change detection.

PLATES

ates 1a, 1c

These two plates are evidence of the negative impacts of urbanization in the study area. Plate 1a and 1c show what is left of the large stream that cuts across the Bauchi road on the maps. Refuse and Garbage have blocked its path of flow and buildings have taken over most of banks of the stream. This stream has been reduced and almost blocked. It is now a refuse dump for both industrial and domestic waste. The complete disappearance of the stream is inevitable.

ate 1b

This plate shows the once large and well-maintained racecourse. Farming activities have now taken over it due to lack of available land for farming. At the top right hand corner of the plate, are the recreational structure which have been dilapidated and neglected.

ates 2a, 2b, 2c

Plate 2c shows more organised pattern of packing space. While plate 2b shows the main entrance of the University of Jos which is some of the positive land use system in the study area. Plate 2a shows the Bauchi park in the study area which is an element of high percentage of development.

Plates 3a,3b, 3c.

These are showing wider and better road networks in the study area. This is a positive development in urban development and evidence of the high percentage of development.

Plates 4a, 4b,

These plates show commercial areas. It should be noted that these areas before were not developed and only were occupied by dense vegetation. This is also a positive trend in urbanization, and it is due to the high rate of development.

Plates 5a, 5b, 5c

Showing the erection, construction and presence of new structures emphasizing high rate of development as against the past.



PLATE 1a



PLATE 2a



PLATE 1b




PLATE 2b



PLATE 1c



PLATE 2c



PLATE 3a



PLATE 4a



PLATE 3b



PLATE 4b



PLATE 3c



PLATE 5a




PLATE 5b



PLATE 5c

REFERENCES

- Adeniyi, P.O. (1981). The Role of Remote Sensing in Land use planning. In Nkambwe,(eds),Application of Remote Sensing Techniques in Nigeria, NISOR pub 1.No. 1 and 2.
- Adeniyi, P.O. (1984). "Application of Aerial photography to the Estimation of the Characteristics of Residential Buildings". Nigeria Geographical Journal, Vol. 27 No. 1 and 2.
- Adeniyi, P.O. (1980). Land use change Analysis using Sequential Aerial photography and Computer Techniques' Photo Eng. and Remote Sensing. Vol. 46, No. 11.
- Ajayi, J.A. (1981). "Estimation of Building Construction as indication of Spatial Urban Growth using sequential Aerial photography." In Nkambwe (eds), application of Remote Sensing Techniques in Nigeria, NISOR publ. No. 2
- Anderson, J. R. Hardy, E. E, Roach, J. T. and Witmer, R. E. (1976). A landuse/landcover classification scheme for use with Remote Sensing Data. Professional paper 1964, United States Geological Survey, Washington D. C

6. Avery, G. (1965). Measuring Land use change in USDA photographs. photo.
Eng. and Remote Sensing. No 45, PP. 661-8.
7. Awogbade, M.O (1983). Fulani Pastoralism; Jos Case Study. Ahmadu Bello
University Press Ltd, Zaria.
8. Buchanan, K and Pugh, J. (1955). Land and people in Nigeria. London
University Press, London.
9. Dunford, C E. (1983). Remote Sensing for Rural development planning in
Africa. ITC journal. No. 4.
10. Eyre, E. E. (1970). Census Analysis and population Studies. Photo. Eng. and
Remote Sensing, vol. 36 No. 5.
11. Hsu, S. (1971). Population Estimation. Photo. Eng. and Remote Sensing.
Vol. 37. No 5.
12. Ikhouria, I.A (1993). Vegetation and Land use changes in a Rainforest
Ecosystem. Remote Sensing Journal (Maiden Edition),
April 1993.
13. John, T. E. (1977). A Land use /Land cover classification System for use with
Remote Sensing data. Geological Survey professional paper 964.
U.S. Government Printing Office, Washington D.C pp. 107-159.
14. Lillesand, T M and Krefer, R. W(1979). Remote Sensing and Image
Interpretation. 3rd edition, John Wiley and Sons Inc. U.S.A.

15. Lindgren, D.T. (1971). Dwelling Unit Estimation from Colour infrared photos.
Photo. Eng. Vol. 37.
16. Lo, C.P (1972). Land use change in Clark County, Atlanta, in Lo, C.P.
Applied Remote Sensing. Longmans, New York.
17. Lo, C.P. (1986). Applied Remote Sensing, Longmans, New York.
18. Lo, C.P and Shipman (1990). Geographical Information System (GIS) Approach
To Land Use Change Dynamic Detection. Photo. Eng. and
Sensing. Vol. 39. No. 5.
19. Lo, C. P and Dale, A. Q (1999) A landuse and landcover change:- A Remote
Sensing Approach. In Journal of American Society for
Photogrammetry and Remote Sensing No 1054. September,
2003.
20. Nkambwe, M (1983). Remote Sensing in Urban Population Estimation in
Nigeria: The Example Of Ile-Ife, Application of Remote
Sensing Techniques in Nigeria. NISOR Publ. No.2.
22. Nkambwe, M. (1984). Application of Multi- date Aerial Photographs for the Study
of Physical Urban Growth of Ile-Ife. Nig. Geographical Journal
Vol. 27. No. 1 and 2.
23. Ojigi, M. L (1997). Assessment and Mapping of Sequential Urban Development
of the Federal Capital City, Abuja with Remotely Sensed
Data (1980-1994), A case study of phase I. An M. Tech
Thesis in Department of Geography, F. U. T, Minna.

24. Okhimamhe, A.A (1993). Assessment of Environmental Impact Of Dam Construction in Nigeria: "A case study Of Tiga Dam in Kano State". An M. Tech Thesis in Department of Geography, Federal University of Technology, Minna.
25. Owolabi, T. (1998). Application of Sequential Aerial Photographs for the Detection of Urban Landuse Changes:- A Case Study of Victoria Island, Lagos State. An M. Tech Thesis in Department of Geography, Federal University of Technology, Minna.
26. Rao, K.M (1995). Remote Sensing for Land use planning. International Journal of Remote Sensing. Vol. 16 No. 1 PP. 53-60
27. Roger, B. C. (1985). Preliminary Result of Anthropogenic Albedo Changes Over The Past 15 years In Eastern Canada. Proceedings Of ISLSCP Conference, Rome . PP 497-499
28. Sogunle, W. K. (1990). A land.use/ Land cover Classification System For Use With Remote Sensing Data. U.S Geological Survey. Professional Papers; 904.