Numerical Solution of First and Higher Order IVPs Via a Single Continuous Block Method

Jamiu Garba^{1*}, Khadeejah James Audu ¹, Umaru Mohammed¹, Abd'gafar Tunde Tiamiyu¹

¹Department of Mathematics, Faculty of Physical Sciences, Federal University of Technology, Minna, Nigeria

ABSTRACT

This article focuses on the development and implementation of a single continuous collocation numerical scheme for solving first and higherorder ordinary differential equations (ODEs). By employing the interpolation and collocation technique on power series as basis function, we were able to come up with a continuous scheme from which block methods for effectively solving first and higher order ODEs were derived. This is better and faster than the traditional way of developing a continuous scheme for a specific order of ODE. The method's accuracy is determined to be of order seven, establishing its consistency. Results from the implementation of our method show its applicability on nonlinear equations and application problems from first, second, third, and fourth order ODEs that are of significant implications on various fields in physics, engineering, biology and mathematics. Furthermore, the numerical results generated by the method reveal its effectiveness and accuracy, and also its superiority over some methods that exist in literature.

Keywords: Accuracy, Collocation, Continuous scheme, First order ODEs, Higher order ODEs, Interpolation

1. INTRODUCTION

Mathematical modeling entails the ability to transform issues from a particular field of application into workable mathematical expressions, whose theoretical and numerical evaluations provide understanding, solutions, and recommendations that are advantageous for the initial applications [35]. Numerous fundamental laws of nature in Physics, Chemistry, Biology, and astronomy are most effectively articulated through differential equations [34]. Their applications are widespread in mathematics, particularly in geometry, as well as in engineering, behavioural sciences, industrial mathematics, artificial intelligence. This kind of problem can be represented using either first-order or higher-order ODEs. First-order ODEs are commonly used in studying problems for example, determining the movement of an object that is ascending or descending while experiencing air resistance, and calculating the current in an electrical circuit; population expansion; radioactive decay; mixture problems; and so on. Second-order ODEs are also commonly used while analyzing vibrating systems, electromagnetism, and electrical circuits with capacitors, resistors, and inductors. Third order differential equations can also be used to solve physical problems like thin film flow, electromagnetic waves and gravity-driven flows. In general, solutions to differential equations are used to forecast the behavior of a system at a later time or in an unknown location. But there aren't many analytical ways to solve ODEs in a continuous or closed form, and nonlinear ODEs can be hard to solve or may not have a closed form of solution at all.

Given the challenges associated with solving most of these problems, numerous researchers have focused extensively on applying numerical methods to provide approximate solutions for differential equations. Numerical methods are particularly effective for addressing mathematical problems, leveraging the speed and efficiency of modern digital computers in performing arithmetic operations [42]. The process of solving these problems with high-precision digital computers typically begins with initial data, followed by the execution of suitable algorithms to produce the desired outcomes [43]. Common approaches for solving IVPs are classified into single-step methods, such as the Runge-Kutta methods [21, 43], and multistep methods [1 - 10].

Over the years, various techniques for deriving continuous linear multistep methods (LMMs) aimed at directly solving IVPs have been extensively discussed in the literature. Key techniques include collocation, interpolation, integration,

*Corresponding Author Email: jamiu.garba@st.futminna.edu.ng Submitted: 03.01.2025 Revision Requested: 06.03.2025 Last Revision Received: 10.03.2025 Accepted: 13.03.2025 Published Online: 06.05.2025 Burdur Mehmet Akif Ersoy University Press

https://dergipark.org.tr/sjmakeu



Cite this article as: Garba, J., Audu, K. J., Mohammed, U., Taimiyu, A. T. (2025). Numerical Solution of First and Higher Order IVPs Via a Single Continuous Block Method. Scientific Journal of Mehmet Akif Ersoy University, 8(1): 16-34. **DOI:** <u>https://doi.org/10.70030/sjmakeu.1611387</u>

Content of this journal is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.



and the use of interpolating polynomials. Basis functions like power series, Chebyshev polynomials, Legendre polynomials, and trigonometric functions have been used for this purpose. (see [7, 10,17, 18, 22, 44, 45, 46]). Researchers have found linear multistep method very useful in solving ODEs. Some authors [7, 10, 11, 12] have develop effective methods for first order ODEs while [6, 13, 27, 35, 42, 44] have implemented linear multistep methods on second order ODEs with comparative accuracy. Other higher ODEs have been solved numerically by [2, 14, 15 31, 37, 39, 43, 47, 48, 49]. All of the linear multistep methods developed by these researchers were for a specific order of ODEs. Recently Kuboye and Adeyefa [22] developed a linear multistep method for first, second and third order ODEs.

In this research, we derive a single continuous technique from a hybrid block approach that effectively handles first, second, third, and fourth-order initial value problems. The subsequent portion of this article describes the method's derivation, while the third sets out the analysis that ensures the method's validity. In part four, a variety of numerical problems are addressed, results are displayed graphically, and the method's efficacy is demonstrated by comparing absolute errors to those of recent existing methods in literature. In section five, we offer broad conclusions on our findings.

The general form of the ODE is expressed as follows:

$$z^{(m)} = \left(z(t), z'(t), ..., z^{(m-1)}(t)\right)$$

$$z(t_0) = z_0, z'(t_0) = z'_0, ..., z^{(m-1)}(t_0) = z_0^{(m-1)}$$
(1)

2. DERIVATION OF THE METHOD

As stated in the introduction, the continuous representation of our method will be generated from one of the orthogonal basis functions, preferably the power series, due to its ease of usage. Therefore, the approximate solution to (1) is as follows:

$$Z(t) = \sum_{j=0}^{m+n-1} c_j t^j$$
(2)

where m and n are respectively the number of interpolation and collocation points. We derive a two-step linear multistep method using (2) with degree m + n - 1 = 10. (2) is interpolated at $t = t_n$ and the first, second and third derivatives are collocated at $t = t_{n+2}$. Furthermore, we collocate the fourth derivative at distinct points $t = t_{n+j}; \left(j = 0, \frac{1}{4}, \frac{3}{4}, 1, \frac{5}{4}, \frac{7}{4}, 2\right)$. All these processes lead to a system of nonlinear equations in t_{n+j} with c_j 's as the

unknowns.

$$Z(t_{n}) = z_{n}$$

$$Z'(t_{n+2}) = f_{n+2}$$

$$Z''(t_{n+2}) = g_{n+2}$$

$$Z'''(t_{n+2}) = p_{n+2}$$

$$Z^{iv}(t_{n+j}) = g_{n+j}$$
(3)

We solved for the unknowns using the matrix inversion method via a mathematical software – Maple 2015; and then substitute c_i 's values back into (2) to obtain the continuous scheme in the form:



$$Z(t) = z_{n} + h\alpha'(t) f_{n+2} + h^{2}\alpha''(t) g_{n+2} + h^{3}\alpha'''(t) p_{n+2} + h^{4} \left(\beta_{0}(t)q_{n} + \beta_{\frac{1}{4}}(t)q_{n+\frac{1}{4}} + \beta_{\frac{3}{4}}(t)q_{n+\frac{3}{4}} + \beta_{1}(t)q_{n+1} + \beta_{\frac{5}{4}}(t)q_{n+\frac{5}{4}} + \beta_{\frac{7}{4}}(t)q_{n+\frac{7}{4}} + \beta_{2}(t)q_{n+2}\right)$$

$$(4)$$

Evaluating (4) at the non-interpolating points gives 6 discrete schemes that form the block method as follows:

$$z_{n+\frac{1}{4}} = z_n + \frac{1}{4} h f_{n+2} - \frac{15}{32} h^2 g_{n+2} + \frac{169}{384} h^3 p_{n+2} + \frac{1}{13005619200} h^4 \begin{pmatrix} 1837375q_n - 21942710q_{n+\frac{1}{4}} - 353393782q_{n+\frac{3}{4}} + 131048330q_{n+1} \\ -1219981490q_{n+\frac{5}{4}} - 1715628930q_{n+\frac{7}{4}} - 409914793q_{n+2} \end{pmatrix}$$
(5)

$$z_{n+\frac{3}{4}} = z_n + \frac{3}{4} h f_{n+2} - \frac{39}{32} h^2 g_{n+2} + \frac{129}{128} h^3 p_{n+2} + \frac{1}{4335206400} h^4 \begin{pmatrix} 1070517q_n - 8616430q_{n+\frac{1}{4}} - 163366014q_{n+\frac{3}{4}} + 67852190q_{n+1} \\ -776333082q_{n+\frac{5}{4}} - 1257048730q_{n+\frac{7}{4}} - 312696051q_{n+2} \end{pmatrix}$$
(6)

$$z_{n+1} = z_n + hf_{n+2} - \frac{3}{2}h^2g_{n+2} + \frac{7}{6}h^3p_{n+2} + \frac{1}{793800}h^4 \begin{pmatrix} 283q_n - 1880q_{n+\frac{1}{4}} - 28840q_{n+\frac{3}{4}} + 11585q_{n+1} \\ -150248q_{n+\frac{5}{4}} - 260760q_{n+\frac{7}{4}} - 66265q_{n+2} \end{pmatrix}$$
(7)

$$z_{n+\frac{5}{4}} = z_n + \frac{5}{4}hf_{n+2} - \frac{55}{32}h^2g_{n+2} + \frac{485}{384}h^3p_{n+2} + \frac{1}{2601123840}h^4 \begin{pmatrix} 1221691q_n - 7249910q_{n+\frac{1}{4}} - 88889206q_{n+\frac{3}{4}} + 28468370q_{n+1} \\ -491995826q_{n+\frac{5}{4}} - 906420930q_{n+\frac{7}{4}} - 234924589q_{n+2} \end{pmatrix}$$
(8)

$$z_{n+\frac{7}{4}} = z_n + \frac{7}{4} h f_{n+2} - \frac{63}{32} h^2 g_{n+2} + \frac{511}{384} h^3 p_{n+2} + \frac{1}{1857945600} h^4 \begin{pmatrix} 1082161q_n - 5977070q_{n+\frac{1}{4}} - 58808638q_{n+\frac{3}{4}} + 10672550q_{n+1} \\ -34408610q_{n+\frac{5}{4}} - 664954650q_{n+\frac{7}{4}} - 176256247q_{n+2} \end{pmatrix}$$
(9)

$$z_{n+2} = z_n + 2hf_{n+2} - 2h^2g_{n+2} + \frac{4}{3}h^3p_{n+2} + \frac{1}{99225}h^4 \begin{pmatrix} 29q_n - 160q_{n+\frac{1}{4}} - 1568q_{n+\frac{3}{4}} + 280q_{n+1} \\ -1568q_{n+\frac{5}{4}} - 17760q_{n+\frac{7}{4}} - 4712q_{n+2} \end{pmatrix}$$
(10)

.

3. ANALYSIS OF THE BLOCK METHOD

At this point, following the approaches adopted in [18, 27, 28], the order of accuracy of our method is:

 $(7,7,7,7,7,7)^{T}$ and error constants given as

18

(221815417	66674131		593	184617553	
	3515887072051200	[•] 390654119116	800	2682408960	703177414410240	,
	150066817	251				
$\left(\cdot \right)$	502269581721600	838252800)



We show the working of equation (10) as follows:

$$\begin{split} \varepsilon_{0} &= 1 - 1 = 0 \\ \varepsilon_{1} &= 2(1) - 2 = 0 \\ \varepsilon_{2} &= \frac{1}{2!} (2^{2}) - (2 \times 2) - (-2) = 0 \\ \varepsilon_{3} &= \frac{1}{3!} (2^{3}) - \frac{1}{2!} (2^{2} \times 2) - (-2 \times 2) - (\frac{4}{3}) = 0 \\ \varepsilon_{4} &= \frac{1}{4!} (2^{4}) - \frac{1}{3!} (2^{3} \times 2) - \frac{1}{2!} (-2 \times 2^{2}) - (2 \times \frac{4}{3}) = 0 \\ \varepsilon_{5} &= \frac{1}{5!} (2^{5}) - \frac{1}{4!} (2^{4} \times 2) - \frac{1}{3!} (-2 \times 2^{3}) - \frac{1}{2!} (2^{2} \times \frac{4}{3}) - \\ &\left(\frac{58}{99225} + \frac{64}{19845} + \frac{64}{2025} - \frac{16}{2835} + \frac{2624}{14175} + \frac{2368}{6615} + \frac{9424}{99225}\right) = 0 \\ \vdots \\ \varepsilon_{11} &= \frac{1}{11!} (2^{11}) - \frac{1}{10!} (2^{10} \times 2) - \frac{1}{9!} (-2 \times 2^{9}) - \frac{1}{8!} (2^{8} \times \frac{4}{3}) - \\ &\frac{1}{7!} \left(\left(\frac{1}{4}\right)^{7} \frac{64}{19845} + \left(\frac{3}{4}\right)^{7} \frac{64}{2025} - (1)^{7} \frac{16}{2835} + \left(\frac{5}{4}\right)^{7} \frac{2624}{14175} + \left(\frac{7}{4}\right)^{7} \frac{2368}{6615} + (2)^{7} \frac{9424}{99225} \right) \\ &= \frac{251}{838252800} \end{split}$$

The techniques described in sections (5) through (10) can typically be represented using a matrix-based difference equation, which is detailed in the following steps.:

$$A^{(1)}Z_{w} = A^{(0)}Z_{w-1} + hBF_{w} + h^{2}CG_{w} + h^{3}DP_{w} + h^{4}\left[E^{(0)}Q_{w-1} + E^{(1)}Q_{w}\right]$$
(11)

where

$$\begin{split} & Z_{w} = \left(z_{n+\frac{1}{4}}, z_{n+\frac{3}{4}}, z_{n+1}, z_{n+\frac{5}{4}}, z_{n+\frac{7}{4}}, z_{n+2}\right)^{T}, \\ & Z_{w-1} = \left(z_{n-\frac{1}{4}}, z_{n-\frac{3}{4}}, z_{n-1}, z_{n-\frac{5}{4}}, z_{n-\frac{7}{4}}, z_{n-2}\right)^{T}, \\ & F_{w} = \left(f_{n+2}\right)^{T}, G_{w} = \left(g_{n+2}\right)^{T}, P_{w} = \left(p_{n+2}\right)^{T}, \\ & Q_{w} = \left(q_{n+\frac{1}{4}}, q_{n+\frac{3}{4}}, q_{n+1}, q_{n+\frac{5}{4}}, q_{n+\frac{7}{4}}, q_{n+2}\right)^{T}, \\ & Q_{w-1} = \left(q_{n-\frac{1}{4}}, q_{n-\frac{3}{4}}, q_{n-1}, q_{n-\frac{5}{4}}, q_{n-\frac{7}{4}}, q_{n-2}\right)^{T} \end{split}$$

and the matrices $A^{(1)}, A^{(0)}, B, C, D, E^{(1)}$ and $E^{(0)}$ are matrices whose elements are given by the coefficients of the block method.

Definition 3.1. The newly developed two-step hybrid block method (5 – 10) is considered zero-stable if and only if the first characteristic polynomial $\xi(t)$ has roots that satisfy $|t| \le 1$, and for the roots $|t| \le 1$, their multiplicity does not exceed one. The characteristic function of this newly derived method is presented below:





 $\xi(\lambda) = \lambda A^{(1)} - A^{(0)}$

$$\xi(\lambda) = \lambda \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} - \begin{pmatrix} 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 \\ \end{pmatrix} = (\lambda + 1)\lambda^{5}$$
(13)

the solution of which is $\lambda = (0, 0, 0, 0, 0, 1)$. Hence, our method is zero-stable.

Definition 3.2. If the order of the hybrid block approach is greater than or equal to one, it is consistent. Our method is consistent as a result of order *p*=7.

Theorem 3.1. Zero stability and consistency are adequate requirements for the convergence of a linear multistep method [26, 17, 27]. The new hybrid block technique is convergent due to its zero-stability and consistency.

To analyze the stability characteristics of the developed scheme, it is applied to a standard test problem.

$$z' = \lambda z, \qquad z'' = \lambda^2 z, \qquad z''' = \lambda^3 z, \qquad z^{i\nu} = \lambda^4 z, \qquad \operatorname{Re}(\lambda) < 0$$
 (14)

to yield

$$Z_{w} = \xi(w) Z_{w-1}, \quad w = z\lambda$$
⁽¹⁵⁾

where the matrix $\xi(w)$ is given as:

$$\zeta(w) = \left(Z^{(1)} - wF - w^2G - w^3P - w^4Q^{(1)}\right)^{-1} \left(Z^{(0)} + w^4Q^{(0)}\right)$$
(16)

The Matrix $\zeta(w)$ has eigenvalues $(0, 0, 0, 0, 0, \lambda_6)$, and the dominant eigenvalue $\lambda_6: \mathbb{C} \to \mathbb{C}$ is a rational function with real coefficients given by

$$R(w) = \frac{Y(w)}{X(w)}$$

The stability region is illustrated in Figure 1, indicating that the method is A-stable.



Fig 1. Region of absolute stability of the method



4. NUMERICAL EXPERIMENT

The recently developed hybrid block method is applied to both first and higher-order ODEs. This continuous approach results in a primary discrete two-step multi-derivative block method (as outlined in equations 5 - 10), along with supplementary methods that are integrated and utilized as a block method to generate approximations concurrently. $\{y_{n+1}, y_{n+2}\}$ at a block points $\{x_{n+1}, x_{n+2}\}$, $h = x_{n+1} - x_n$, n = 0, ..., N-2, on a partition [a, b], where $a, b \in \Box$ is the interval of integration, h is the constant step-size, n is a grid index and N > 0 is the number of steps. We obtain initial conditions at $x_{n+2}, n = 0, 1, ..., N-2$, using the computed values y_{n+2} over smaller intervals $[x_0, x_2], ..., [x_{N-2}, x_N]$. For example, when $n = 0, [y_1, y_2]$ are acquired at the same time across the smaller interval $[x_0, x_2]$, as y_0 is established from the initial value problem, for $n = 2, [y_3, y_4]$ are also acquired simultaneously over the smaller interval $[x_3, x_4]$, as is now understood from the earlier section, and so forth. Consequently, the smaller interval $[x_n, x_{n+2}]$ does not overlap, and the solutions derived in this manner are more precise than those obtained through traditional predictor-corrector methods. Similar approach is applied to higher order ODEs considered – the first, second and third derivatives of the continuous scheme are evaluated at all points to cater for the higher derivative terms in the general higher order ODEs considered. We present the graphical solution of the problems and compare absolute errors with some existing methods in the literature.

4.1 First-Order Problems

The block method (5) to (10) is used directly on some first order problems.

Problem 1: The SIR model is an epidemiological framework that tracks the hypothetical number of individuals within a closed population who are affected by an infectious disease over time. This type of model derives its name from the interconnected equations that describe the populations of susceptible individuals S(t), infected individuals I(t), and recovered individuals R(t). It serves as an effective and simple model for various infectious diseases, such as measles, mumps, and rubella. The model is represented by the three interconnected equations provided below.:

$$\frac{dS}{dt} = \mu (1 - S) - \beta IS$$

$$\frac{dI}{dt} = \mu I - \gamma I + \beta IS$$

$$\frac{dR}{dt} = \mu R + \gamma I$$

where μ , β and γ are positive parameters. Define *z* to be:

$$z = S + I + R$$

and adding (13), (14) and (15), the following evolution equation for z is obtained.

$$z' = \mu (1 - z)$$

Kuboye and Adeyefa [22] solved this problem with the following parameters:

$$\mu = \frac{1}{2}, \ z(0) = \frac{1}{2}, \ h = 0.1$$

Exact solution: $z(t) = 1 - \mu e^{-\mu t}$





Problem 2: We consider the Riccati differential equation solved in Khalsaraei et. al. [20]

$$z' = 1 + 2z - z^2; \quad z(0) = 0, \quad 0 \le t \le 10$$

Exact solution: $z(t) = 1 + \sqrt{2} \tanh\left(\sqrt{2}t + \frac{1}{2}\log\left(\frac{\sqrt{2}-1}{\sqrt{2}+1}\right)\right)$

Problem 3: We consider the nonlinear system of stiff chemical problem solved in Akinfenwa et. al. [11]

 $z_{1}' = \lambda z_{1} + z_{2}^{2}, \quad z_{1}(0) = -\frac{1}{\lambda + 2} \\ z_{2}' = -z_{2}, \quad z_{2}(0) = 1$

The exact solution is given as $z_1(t) = -\frac{\exp(-2t)}{\lambda+2}$, $z_2(t) = \exp(-t)$ where $\lambda = 10000$.



Fig 2. The profile solution for problem 1





Fig 4. The profile solution for problem 3



	Table 1. Error comparison for Problem 1						
t	Error in [22]	Error in [19]	Error in [4]	Error in New			
	(h=0.1)	(h = 0.1)	(h=0.1)	$methodig(h\!=\!0.1ig)$			
0.1	3.846×10 ⁻¹³	1.998×10 ⁻¹⁵	9.104×10 ⁻¹⁵	4.736×10 ⁻²²			
0.2	7.319×10 ⁻¹³	3.886×10 ⁻¹⁵	7.105 × 10 ⁻¹⁵	6.225×10 ⁻²²			
0.3	1.044 × 10 ⁻¹²	5.440×10 ⁻¹⁵	8.882×10 ⁻¹⁵	1.021×10 ⁻²¹			
0.4	1.324×10 ⁻¹²	6.994×10 ⁻¹⁵	2.121×10 ⁻¹⁴	1.127×10 ⁻²¹			
0.5	1.575×10 ⁻¹²	8.216×10 ⁻¹⁵	1.368×10 ⁻¹³	1.459×10 ⁻²¹			
0.6	1.797×10 ⁻¹²	9.548×10 ⁻¹⁵	7.983×10 ⁻¹³	1.529×10 ⁻²¹			
0.7	$1.995 imes 10^{-12}$	$1.055 imes 10^{-14}$	3.699×10 ⁻¹²	1.805×10 ⁻²¹			
0.8	2.168×10 ⁻¹²	1.132×10^{-14}	-	1.845×10 ⁻²¹			
0.9	2.320×10 ⁻¹²	1.221×10 ⁻¹⁴	-	1.845×10 ⁻²¹			
1.0	2.452×10 ⁻¹²	$1.288 imes 10^{-14}$	-	2.086×10 ⁻²¹			

Table 2. Error comparison for Problem 2

t	Error in [19]	Error in [4]	Error in New method
	(h = 0.05)	(h=0.1)	(h=0.1)
1	1.418×10 ⁻¹¹	9.104×10 ⁻¹⁵	2.168×10 ⁻¹¹
2	7.234×10 ⁻¹³	7.105×10 ⁻¹⁵	1.052×10 ⁻¹²
3	1.163×10 ⁻¹³	8.882×10 ⁻¹⁵	5.216×10 ⁻¹⁴
4	2.132×10 ⁻¹⁴	2.121×10 ⁻¹⁴	3.144×10 ⁻¹⁵
5	2.664×10 ⁻¹⁵	1.368×10 ⁻¹³	1.970×10 ⁻¹⁶
6	4.441×10 ⁻¹⁶	7.983×10 ⁻¹³	1.233×10 ⁻¹⁷
7	4.441×10 ⁻¹⁶	3.699×10 ⁻¹²	7.213×10 ⁻¹⁹
8	4.441×10 ⁻¹⁶	-	4.702×10 ⁻²⁰
9	4.441×10 ⁻¹⁶	-	2.977×10 ⁻²¹
10	4.441×10^{-16}	-	1.850×10 ⁻²²

Table 3. Error comparison for Problem 3

t	Z _i	Error in [20] $(h = 0.0001)$	Error in [11] $(h = 0.01)$	Error in [26] $(h=0.1)$	Error in New method $ig(h\!=\!0.1ig)$
3	(z_1) (z_2)	1.779×10 ⁻²⁰ 2.079×10 ⁻²⁰	2.030×10 ⁻¹⁹ 1.440×10 ⁻¹⁴	3.790×10 ⁻²² 2.998×10 ⁻¹⁸	1.972×10^{-23} 1.981×10^{-18}
5	$egin{pmatrix} (z_1) \ (z_2) \end{pmatrix}$	2.493×10 ⁻¹⁹ 4.664×10 ⁻¹³	1.200×10^{-20} 3.210×10^{-15}	$\frac{1.60 \times 10^{-21}}{6.740 \times 10^{-19}}$	6.019×10 ⁻²⁵ 4.468×10 ⁻¹⁹
10	$egin{array}{c} (z_1) \ (z_2) \end{array}$	5.743×10 ⁻²⁰ 6.346×10 ⁻¹²	1.110×10 ⁻²⁰ 4.380×10 ⁻¹⁷	7.120×10^{-20} 9.080×10^{-21}	5.466×10^{-29} 6.021×10^{-21}

4.2 Second-Order Problems

In order to implement our method on second order IVPs, we take the first derivative of the continuous scheme (4) and evaluated at points t_{n+j} , $\left(j=0,\frac{1}{4},\frac{3}{4},1,\frac{5}{4},\frac{7}{4},2\right)$ with the block schemes (5 – 10).

Problem 4: Consider the nonlinear second order IVP from Abdelrahim and Omar [1].

23



$$z'' = t(z')^2, \ z(0) = 1, \ z'(0) = 0.5$$

The exact solution is: $z(t) = \arctan h\left(\frac{1}{2}t\right) + 1$

Problem 5: Cooling of a body. Source: Kwanamu et. al. [24]

The temperature z degree of a body t minutes after being placed in a certain room, satisfies the differential equation

$$3\frac{d^{2}z}{dt} + \frac{dz}{dt} = 0, \quad z(0) = 60, \quad \frac{dz}{dt}\Big|_{t=0} = -\frac{80}{9}$$



Fig 7. The profile solution for problem 6

Problem 6: Consider the nonlinear problem:

 $z'' - 2z^3 = 0$ (Source: Ogunlaran & Kehinde [30])

with the following initial conditions



$$z(1) = 1, z'(1) = -1 \text{ and } h = \frac{0.1}{40}$$

Exact solution:
$$Z(t) = \frac{1}{t}$$

Table 4. Error comparison for Problem 4					
t	Error in [1]	Error in [7]	Error in new	Error in New	
	$\left(h=\frac{1}{2}\right)$	(h = 0.1)	method	method $\left(h=\frac{1}{2}\right)$	
	(30)		(n = 0.1)	(30)	
0.1	1.310×10 ⁻¹⁶	1.957×10^{-13}	1.539×10 ⁻¹⁵	6.552×10 ⁻²⁰	
0.2	3.975×10 ⁻¹⁴	6.040×10 ⁻¹³	6.459×10 ⁻¹⁵	2.934×10 ⁻¹⁹	
0.3	1.021×10 ⁻¹⁴	1.262×10 ⁻¹²	1.791×10 ⁻¹⁴	7.771×10 ⁻¹⁹	
0.4	3.304×10 ⁻¹³	3.715×10 ⁻¹²	3.972×10 ⁻¹⁴	1.726×10 ⁻¹⁸	
0.5	-	7.919×10 ⁻¹²	8.698×10 ⁻¹⁴	3.588×10 ⁻¹⁸	
0.6	-	1.416×10^{-11}	1.781×10 ⁻¹³	7.353×10 ⁻¹⁸	
0.7	-	3.616×10 ⁻¹¹	3.987×10 ⁻¹³	1.531×10 ⁻¹⁷	
0.8	-	7.473×10 ⁻¹¹	8.522×10 ⁻¹³	3.306×10 ⁻¹⁷	
0.9	-	1.335×10^{-10}	2.182×10 ⁻¹²	7.538×10 ⁻¹⁷	
1.0	1.293×10 ⁻¹²	4.317×10 ⁻¹⁰	5.160×10 ⁻¹²	1.844×10 ⁻¹⁶	

Table 5. Error comparison for Problem 5, h = 0.1

t	Error in [38]	Error in [41]	Error in [33]	Error in [24]	Error in New
					method
0.1	3.550×10 ⁻¹¹	2.300×10 ⁻¹⁷	7.476×10 ⁻⁰⁶	-	9.000×10 ⁻²³
0.2	4.580×10 ⁻¹¹	1.710×10 ⁻¹⁶	2.939×10 ⁻⁰⁵	4.000×10 ⁻¹⁸	3.600×10 ⁻²²
0.3	7.000×10 ⁻¹¹	4.370×10 ⁻¹⁶	6.480×10 ⁻⁰⁵	9.000×10 ⁻¹⁸	8.300×10 ⁻²²
0.4	6.500×10 ⁻¹¹	8.130×10 ⁻¹⁶	1.128×10 ⁻⁰⁵	1.700×10 ⁻¹⁷	1.450×10 ⁻²¹
0.5	3.330×10 ⁻¹¹	1.290×10 ⁻¹⁵	1.725×10^{-04}	2.600×10 ⁻¹⁷	2.230×10 ⁻²¹
0.6	4.200×10 ⁻¹¹	1.864×10 ⁻¹⁵	2.431×10 ⁻⁰⁴	3.800×10 ⁻¹⁷	3.170×10 ⁻²¹
0.7	4.380×10 ⁻¹¹	2.525×10^{-15}	3.238×10 ⁻⁰⁴	5.100×10 ⁻¹⁷	4.240×10 ⁻²¹
0.8	1.070×10^{-10}	3.269×10 ⁻¹⁵	4.139×10 ⁻⁰⁴	6.500×10 ⁻¹⁷	5.410×10 ⁻²¹
0.9	6.580×10 ⁻¹¹	4.089×10 ⁻¹⁵	5.127×10^{-04}	8.100×10 ⁻¹⁷	6.720×10 ⁻²¹
1.0	1.6900×10 ⁻¹⁰	4.980×10 ⁻¹⁵	$6.195 imes 10^{-04}$	9.700×10 ⁻¹⁷	8.140×10 ⁻²¹

Table 6. Error comparison for Problem 6

t	Exact Solution	Error in [30]	Error in New method
1.0025	0.997506234413965	3.160×10 ⁻¹⁷	5.400×10 ⁻²⁹
1.0050	0.995024875621891	7.550×10 ⁻¹⁷	2.480×10 ⁻²⁸
1.0075	0.992555831265509	1.190×10 ⁻¹⁶	5.650×10 ⁻²⁸
1.0100	0.990099009900990	1.510×10 ⁻¹⁶	1.005×10 ⁻²⁷
1.0125	0.987654320987654	1.810×10 ⁻¹⁶	1.566×10 ⁻²⁷
1.0150	0.985221674876847	2.220×10 ⁻¹⁶	2.241×10 ⁻²⁷
1.0175	0.982800982800983	2.630×10 ⁻¹⁶	3.031×10 ⁻²⁷
1.0200	0.980392156862745	2.930×10 ⁻¹⁶	3.929×10 ⁻²⁷
1.0225	0.977995110024450	3.210×10 ⁻¹⁶	4.936×10 ⁻²⁷
1.0250	0.975609756097561	3.590×10 ⁻¹⁶	6.044×10 ⁻²⁷



4.3 Third-Order Problems

In order to implement our method on third order IVPs, we take the first and second derivatives of the continuous scheme (4) and evaluated at points t_{n+j} , $\left(j=0,\frac{1}{4},\frac{3}{4},1,\frac{5}{4},\frac{7}{4},2\right)$ with the block schemes (5 – 10).

Problem 7: Consider the nonlinear IVP:

$$z''' = (2tz''+1)z'; z(0) = 1, z'(0) = 0.5, z''(0) = 0$$

The exact solution is: $z(t) = \tan^{-1}\left(\frac{1}{2}t\right) + 1$

Source: Yakusak &. Owolanke [47]

Problem 8: Consider the nonlinear ODE

$$z''' = \frac{3}{8z^5}; z(0) = 1, z'(0) = \frac{1}{2}, z''(0) = -\frac{1}{4}$$

Exact solution: $z(t) = \frac{1}{\sqrt{1+t}}$

Source: Adeyeye & Omar [8]

Problem 9:

$$z''' = tz'' - (tz)^{2} + t\sin(t) - \cos(t) + t^{2}\sin^{2}(t); \quad z(0) = 0, \quad z'(0) = 1, \quad z''(0) = 0$$

Exact solution: $z(t) = \sin(t)$

Source: Source: Adeyeye & Omar [8]

1.0

	Table 7. Error comparison for Problem 7					
t	Error in [47]	Error in [9]	Error in [31]	Error in New		
	(h = 0.01)	(h = 0.01)	(h = 0.01)	method		
				(h = 0.01)		
0.1	9.609×10 ⁻¹²	1.931×10 ⁻⁰⁸	2.043×10 ⁻¹⁴	4.401×10 ⁻²⁴		
0.2	7.072×10 ⁻¹⁰	5.608×10 ⁻⁰⁷	8.371×10 ⁻¹⁴	3.661×10 ⁻²³		
0.3	6.693×10 ⁻⁰⁹	3.755×10 ⁻⁰⁶	2.813×10 ⁻¹³	1.353×10 ⁻²²		
0.4	3.142×10 ⁻⁰⁸	1.340×10 ⁻⁰⁵	7.667×10 ⁻¹³	3.665×10 ⁻²²		
0.5	1.051×10 ⁻⁰⁷	3.259×10 ⁻⁰⁵	1.853×10 ⁻¹²	8.554×10 ⁻²²		
0.6	2.852×10 ⁻⁰⁷	5.816×10 ⁻⁰⁵	4.163×10 ⁻¹²	1.855×10 ⁻²¹		
0.7	6.777×10 ⁻⁰⁷	7.152×10 ⁻⁰⁵	8.965×10 ⁻¹²	3.905×10 ⁻²¹		
0.8	1.467×10 ⁻⁰⁶	2.564×10 ⁻⁰⁵	-	8.224×10 ⁻²¹		
0.9	2.983×10 ⁻⁰⁶	1.709×10 ⁻⁰⁴	-	1.774×10 ⁻²⁰		

4.003×10⁻²⁰

 6.706×10^{-04}

 $6.189 imes 10^{-06}$





Fig 8. The profile solution for problem 7





Fig 10. The profile solution for problem 9

Table 8. Comparison of errors for Problem 8				
t	Exact Solution	Error in [8] $\left(h = 0.1 ight)$	Error in New method $ig(h\!=\!0.1ig)$	Error in New method $(h = 0.01)$
0.2	1.095445115010332226913940	2.181×10 ⁻¹¹	3.890×10 ⁻¹³	3.870×10 ⁻²¹
0.4	1.183215956619923208513466	$7.070 imes 10^{-11}$	1.812×10^{-12}	2.242×10^{-20}
0.6	1.264911064067351732799557	1.348×10^{-10}	4.673×10 ⁻¹²	5.891×10 ⁻²⁰
0.8	1.341640786499873817845504	2.106×10^{-10}	9.038×10 ⁻¹²	1.138×10^{-19}
1.0	1.414213562373095048801689	2.964×10^{-10}	$1.490 imes 10^{-11}$	1.871×10 ⁻¹⁹
1.2	1.483239697419132589742279	3.914×10 ⁻¹⁰	2.225×10^{-11}	2.785×10^{-19}
1.4	1.549193338482966754071706	4.947×10 ⁻¹⁰	3.106×10 ⁻¹¹	3.877×10 ⁻¹⁹
1.6	1.612451549659709930473323	$6.059 imes 10^{-10}$	4.128×10 ⁻¹¹	5.142×10^{-19}
1.8	1.673320053068151095956344	7.242×10^{-10}	$5.290 imes 10^{-11}$	6.576×10 ⁻¹⁹
20	1 732050807568877293527446	8.492 × 10 ⁻¹⁰	6.586 × 10 ⁻¹¹	8.174×10 ⁻¹⁹

Table 8. Comparison of errors for Problem 8



Table 9. Error comparison for Problem 9					
t	Exact Solution	Error in [8]	Error in New	Error in New	
		(h = 0.1)	method	method	
		, , , , , , , , , , , , , , , , , , ,	(h = 0.1)	(h = 0.01)	
0.1	0.099833416646828152307	6.661×10 ⁻¹⁶	7.851×10 ⁻¹⁹	3.435×10 ⁻²⁷	
0.2	0.19866933079506121546	3.914×10 ⁻¹⁵	3.028×10 ⁻¹⁸	1.919×10 ⁻²⁶	
0.3	0.29552020666133957511	1.243×10 ⁻¹⁴	8.180×10 ⁻¹⁸	6.472×10 ⁻²⁶	
0.4	0.38941834230865049167	2.887×10^{-14}	1.779×10 ⁻¹⁷	1.538×10 ⁻²⁵	
0.5	0.47942553860420300027	5.601×10 ⁻¹⁴	3.339×10 ⁻¹⁷	3.017×10 ⁻²⁵	
0.6	0.56464247339503535720	9.692×10 ⁻¹⁴	5.667×10 ⁻¹⁷	5.243×10 ⁻²⁵	
0.7	0.64421768723769105367	1.547×10 ⁻¹³	8.933×10 ⁻¹⁷	8.383×10 ⁻²⁵	
0.8	0.71735609089952276163	2.326×10 ⁻¹³	1.333×10 ⁻¹⁶	1.261×10 ⁻²⁴	
0.9	0.78332690962748338846	3.346×10 ⁻¹³	1.904×10 ⁻¹⁶	1.812×10 ⁻²⁴	
1.0	0.84147098480789650665	4.644 × 10 ⁻¹³	2.627×10 ⁻¹⁶	2.509×10 ⁻²⁴	

4.4 Fourth-Order Problems

Problem 10: Consider the oscillatory problem arising from ship dynamics:

$$z^{i\nu} + 3z'' + z(2 + \varepsilon \cos \lambda t) = 0; \quad z(0) = 1, \quad z'(0) = 0, \quad z''(0) = 0, \quad z'''(0) = 0$$

Where $\lambda = 0$ for existence of the exact solution: $z(t) = 2\cos t - \cos(\sqrt{2}t)$

Source: Familua & Omole [16]

Problem 11: Consider the nonlinear problem:

$$z^{iv} = (z')^{2} - zz''' - 4t^{2} + e^{t} (1 - 4t + t^{2}); \quad z(0) = 1, \quad z'(0) = 1, \quad z''(0) = 3, \quad z'''(0) = 1$$

Exact solution: $z(t) = t^2 + e^t$. (Source: Familua & Omole [16])

Problem 12: consider the nonlinear sinusoidal problem:

$$z^{iv} = z^2 + \sin^2(t) - \cos(t) - 1; \quad z(0) = -1, \quad z'(0) = 0, \quad z''(0) = 1, \quad z'''(0) = 0$$

Exact solution: $z(t) = -\cos(t)$. Source: Tiamiyu *et al.* [43]





Fig 12. The profile solution for problem 11





Fig 13. The profile solution for problem 12

Table 10. Little comparison for Fromeni 10 ($n = 0.005125$	Table 10	. Error com	parison for	Problem 10	(h = 0.003125))
--	----------	-------------	-------------	------------	----------------	---

		-	(/
t	Error in [16]	Error in [16]	Error in [45]	Error in New Method
	(Block mode)	(P-C mode)		
0.003125	6.686×10 ⁻¹³	5.686×10 ⁻¹⁰	1.900×10 ⁻¹⁹	3.657×10 ⁻³⁶
0.006250	1.458×10 ⁻¹¹	1.768×10^{-10}	2.300×10 ⁻¹⁹	1.262×10 ⁻³⁵
0.009375	1.083×10^{-10}	$5.910 imes 10^{-09}$	8.600×10 ⁻¹⁹	4.233×10 ⁻³⁵
0.001250	3.918×10 ⁻¹⁰	5.768×10 ⁻⁰⁹	$1.380 imes 10^{-18}$	1.042×10 ⁻³⁴
0.015625	$1.025 imes 10^{-09}$	1.100×10^{-08}	3.530×10 ⁻¹⁸	2.271×10 ⁻³⁴
0.018750	2.217×10 ⁻⁰⁹	6.899×10 ⁻⁰⁸	5.310×10 ⁻¹⁸	4.359×10 ⁻³⁴
0.021875	4.226×10 ⁻⁰⁹	4.636×10 ⁻⁰⁸	$8.880 imes 10^{-18}$	7.728×10 ⁻³⁴
0.025000	7.358×10 ⁻⁰⁹	5.788×10 ⁻⁰⁷	3.922×10 ⁻¹⁷	1.276×10 ⁻³³
0.028125	$1.197 imes 10^{-08}$	2.246×10 ⁻⁰⁷	5.846×10 ⁻¹⁷	2.001×10 ⁻³³
0.031250	$1.846 imes 10^{-08}$	2.846×10^{-07}	8.477×10 ⁻¹⁷	3.001×10 ⁻³³

Table 11. Error comparison for Problem 10 using different values of h			
t	Exact solution	Error in New method	Error in New method
		(h = 0.01)	(h=0.1)
0.1	0.99999167499652860438	3.477×10 ⁻²⁷	4.196×10 ⁻¹⁸
0.2	0.99986719911195714198	5.225×10 ⁻²⁶	1.437×10 ⁻¹⁷
0.3	0.99933105226749824584	2.522×10 ⁻²⁵	4.756×10 ⁻¹⁷
0.4	0.99790057330915505191	8.075×10 ⁻²⁵	1.157×10^{-16}
0.5	0.99492052670511528095	1.938×10 ⁻²⁴	2.4783×10 ⁻¹⁶
0.6	0.98958301770794685383	3.941×10 ⁻²⁴	4.670×10 ⁻¹⁶
0.7	0.98095229007588226219	7.137×10 ⁻²⁴	8.088×10 ⁻¹⁶
0.8	0.96799382462962246366	1.186×10 ⁻²³	1.302×10^{-15}
0.9	0.94960705358355858551	1.844×10 ⁻²³	1.981×10 ⁻¹⁵
1.0	0.92466091697090496135	2.718 [×] 10 ⁻²³	2.874×10 ⁻¹⁵



Table 12. Error comparison for Problem 11			
t	Error in [16]	Error in [23]	Error in New
			method
0.031250	1.149×10 ⁻¹²	1.788×10^{-10}	2.000×10 ⁻²⁴
0.062500	1.885×10^{-11}	1.134×10^{-08}	9.000×10 ⁻²⁴
0.093750	9.780×10 ⁻¹¹	1.196×10^{-07}	2.300×10 ⁻²³
0.125000	3.166×10 ⁻¹⁰	6.401×10 ⁻⁰⁷	4.800×10 ⁻²³
0.156250	7.909×10 ⁻¹⁰	2.349×10^{-06}	9.000×10 ⁻²³
0.187500	1.676×10^{-09}	6.573×10^{-06}	1.500×10 ⁻²²
0.218750	3.169×10 ⁻⁰⁹	1.610×10^{-05}	2.3400×10 ⁻²²
0.250000	5.512×10^{-09}	3.501×10^{-05}	3.450×10 ⁻²²
0.281250	8.995×10 ⁻⁰⁹	$6.985 imes 10^{-05}$	4.870×10 ⁻²²
0.312500	1.396×10^{-08}	1.245×10^{-04}	6.620×10 ⁻²²

 Table 13. Error comparison for Problem 11 using different values of h

t	Exact solution	Error in New method	Error in New method
		(h = 0.1)	(h = 0.01)
0.1	1.11517091807564762481	8.200×10 ⁻¹⁹	2.430×10 ⁻²⁷
0.2	1.26140275816016983392	2.960×10 ⁻¹⁸	1.891×10^{-26}
0.3	1.43985880757600310398	8.000×10 ⁻¹⁸	6.317×10 ⁻²⁶
0.4	1.65182469764127031782	1.706×10^{-17}	1.486×10 ⁻²⁵
0.5	1.89872127070012814685	3.173×10 ⁻¹⁷	2.881×10 ⁻²⁵
0.6	2.18211880039050897488	5.298×10 ⁻¹⁷	4.947×10 ⁻²⁵
0.7	2.50375270747047652162	8.246×10 ⁻¹⁷	7.811×10 ⁻²⁵
0.8	2.86554092849246760458	1.211×10 ⁻¹⁶	1.160×10 ⁻²⁴
0.9	2.86554092849246760458	1.707×10^{-16}	1.645×10 ⁻²⁴
1.0	3.71828182845904523536	2.232×10^{-16}	2.251×10 ⁻²⁴

Table 14. Error comparison for Problem 12				
t	Exact solution	Error in [43] $(h=0.01)$	Error in New method $ig(h\!=\!0.01ig)$	
0.1	- 0.995004165278025766095561987804	1.400×10 ⁻²⁹	5.580×10 ⁻²⁹	
0.2	-0.980066577841241631124196516748	8.240×10 ⁻²⁸	8.465 × 10 ⁻²⁸	
0.3	-0.955336489125606019642310227568	7.825×10 ⁻²⁷	4.236×10 ⁻²⁷	
0.4	-0.921060994002885082798526732052	3.697×10 ⁻²⁶	1.331×10 ⁻²⁶	
0.5	-0.877582561890372716116281582604	1.209×10 ⁻²⁵	3.234×10 ⁻²⁶	
0.6	-0.825335614909678297240952498955	3.148×10 ⁻²⁵	6.675×10 ⁻²⁶	
0.7	-0.764842187284488426255859990192	7.021×10 ⁻²⁵	1.230×10 ⁻²⁵	
0.8	-0.696706709347165420920749981642	1.400×10 ⁻²⁴	2.088×10 ⁻²⁵	
0.9	-0.621609968270664456484716151407	2.564×10 ⁻²⁴	3.323×10 ⁻²⁵	
1.0	-0.540302305868139717400936607443	4.393×10 ⁻²⁴	5.031×10 ⁻²⁵	

Problem 13. And lastly, we also consider the following nearly sinusoidal problem in first-order system of equations. Source: (Akinfenwa *et. al.,* [10])

$z_1' = -21z_1 + z_2 + 2\sin t,$	$z_1(0)=2$
$z_2' = 998z_1 - 999z_2 + 999\cos t - 999\sin t,$	$z_2(0) = 3$



Exact solution: $z_1(t) = 2e^{-t} + \sin t$, $z_2(t) = 2e^{-t} + \cos t$

	HBSDBDF [10]		New Method		
h	Maximum	Relative error	Maximum	Relative error	ROC
	error		error		
0.4	8.9924×10 ⁻⁷	3.6279×10 ⁻⁷	1.2119×10^{-11}	1.8909×10^{-10}	-
0.2	5.9042×10 ⁻⁹	2.6294×10 ⁻⁹	1.6035×10^{-14}	2.2570×10^{-13}	9.56
0.1	4.5695×10 ⁻¹¹	1.8848×10^{-11}	1.7159×10^{-17}	3.0416×10 ⁻¹⁶	9.87
0.05	2.9376×10 ⁻¹³	1.2826×10^{-13}	1.7479×10^{-20}	3.1990×10 ⁻¹⁹	9.94

5. DISCUSSION OF RESULTS

Figures 1–3 respectively display the graphical solution of the SIR model in problem 1, the nonlinear Riccati equation in problem 2 and the nonlinear system of stiff chemical equations in problem 3. In the graphs, we plotted the results generated from our method (red boxes) and the exact solution (blue line). It is easily seen that the new method agrees with the analytical solutions to the problems. Going further, Table 1 shows the comparison of absolute errors for Problem 1. It is shown that the newly derived method has a computational advantage over the methods in ([22], [19] and [4]). Also in Table 2, we show the absolute errors of some methods and ours, for different step sizes in solving problem 2. A comparison of our method with [4] using the same step size (h = 0.1) indicates that our method outperforms the method in [4], and further comparison with the method in [19] (whose step size is smaller) also indicates the superiority of our method over the one in [19]. Table 3 presents a comparative analysis of absolute errors for problem 3 using different step sizes. We solved the nonlinear chemical stiff problem using the step size (h = 0.1)and analysis shows that our method performs better than the methods in [26] (with the same step size), [11] (whose step size is h = 0.01 and [20] (with step size h = 0.0001).

Problems 4–6 are second-order problems considered using the same block method as the first-order problems. Problem 4 is a nonlinear IVP, while Problem 5 is an application problem in the cooling of a body and Problem 6 is also a nonlinear problem. Their behavioural solutions are displayed in figures 4–6 demonstrating agreement in the numerical method that we derived and the exact solutions. To further validate the effectiveness of our method, we compare the absolute errors against those produced by existing techniques. This comparative analysis provides additional evidence supporting the superior performance of our approach. Table 4 shows the comparison of errors for problem 4. It is shown that the new method with h = 0.1 outperforms the method in [7] with the same step size and

that in [1] using the same step size of $\left(h = \frac{1}{30}\right)$. Also, Table 5 shows the comparison of errors for Problem 5, which

indicates the superiority of our method over those in [24, 33, 38, 41] using the same step size. And in Table 6, we show the numerical solution made by the new method, the absolute error and that in [30], it is shown that the new method is better in terms of accuracy.

The nonlinear third-order initial value problems were the next class of ODEs we looked at in this paper. The graphical solutions of problems 7–9 are displayed in Figures 7–9. Furthermore, comparative analysis for Problem 7 in Table 7 indicates that our method is found to give better accuracy than the methods in [31], [47], and [9]. Similarly, analyses in Tables 8 and 9 also show that the newly derived method in this paper gives better accuracy than the method derived in [8] for problems 8 and 9, respectively.

Finally, a class of fourth-order ODEs is also solved by our method. Figures 10–12 display the graphical solutions for problems 10–12, respectively. Problem 10 is solved by [16] using block mode predictor–corrector (P–C) of the linear



multistep method. Also, [45] solved this particular problem in the interval [0.003125, 0.03125] with h = 0.003125. Table 10 shows the comparison of errors between our method, [16] and [33] as mentioned above. It has been demonstrated that the new block method provides a more accurate approximation for the application problem related to ship dynamics. We further compare our method for variable step sizes over [0, 1] using h = 0.1 and h = 0.01 and Table 11 shows that the method gives better approximation as h becomes smaller. Similarly, the nonlinear equation in problem 11 is also solved by [16] and [23] and the results in Table 12 assert that the newly derived method is superior in terms of accuracy to the methods in [16, 23]. Table 13 gives the comparison of errors in our method based on different step sizes. And in Table 14, the comparison of errors for problem 12 is made between our method and that in [43]. It is shown that as the numerical iteration progresses, our method gets closer to the exact solution than the method in [43].

Lastly, we incorporated the sinusoidal problem into a system of first-order equations, as addressed in [10]. We showcased the method's accuracy, rate of convergence (ROC), and strong stability characteristics. We calculated the

maximum error, $(\max |z(t_i) - z_i|)$, relative error $\max_i \frac{|z(t_i) - z_i|}{|1 + z(t_i)|}$ and compared with the method in [10], which

indicates that the new method has higher accuracy than the method in [10]. Also, the $ROC = \log_2\left(\frac{e^{2h}}{e^h}\right)$, for different

step sizes *h*, where e^h is the maximum absolute error for each *h* is calculated as shown in Table 15.

6. CONCLUSION

In this paper, a novel numerical method in the class of linear multistep method is developed for first order and higher order ODEs. The method is derived through the usual interpolation and collocation techniques with power series used as the basis function. Basic numerical properties as established in section 3 show that the method converges and A-stable. The method conveniently solves first, second, third and fourth order IVPs as shown in Tables 1–15. Thirteen numerical problems were considered in all, with nonlinear equation as majority and some application problems. The numerical results generated by our method show its superiority over some existing methods as compared therein. The innovative aspect of this method lies in its capability to effectively solve first, second, third, and fourth order ordinary differential equations (ODEs) without requiring distinct numerical schemes for each type. Therefore, this new approach is proposed as a practical numerical algorithm for tackling first and higher order ODEs. In our upcoming research, we plan to expand this method to address partial differential equations.

ORCID

Jamiu Garba Description https://orcid.org/0000-0002-2850-7730 Khadeejah Audu James Description https://orcid.org/0000-0002-6986-3491 Umaru Mohammed Description https://orcid.org/0000-0002-0777-6536 Tiamiyu Abd'gafar Tunde Description https://orcid.org/0000-0003-1641-7196

REFERENCES

- [1] R. Abdelrahim, and Z. Omar, Direct solution of second-order ordinary differential equations using a single-step hybrid block method of order five, *Math. Comput. App.*, (2016); doi:10.3390/mca21020012.
- [2] Y. A. Abdullahi, Z. Omar, and J. O. Kuboye, Derivation of block predictor–block corrector method for direct solution of third order ordinary differential equations, *Global J. Pure App. Math.*, 11(1)(2016), 343-350.
- [3] C. E. Abhulimen, and L. A. Ukpebor, A new class of second derivative methods for numerical integration of stiff initial value problems, *J. Nig. Math. Soc.*, 38(2)(2019), 259-270.
- [4] O. E. Abolarin, G. B. Ogunware, and L. S. Akinola, An efficient seven-step block method for numerical solution of SIR and growth model, *FUOYE J. Engine. Tech.*, 5(1)(2020), 31–35.
- [5] K. M. Abualnaja, A block procedure with linear multistep methods using Legendre polynomial for solving ODEs, *App. Math. Sci. Res.*, 6(2015), 717-723.



- [6] E. O. Adeyefa, Orthogonal-based hybrid block method for solving general second order initial value problems, *Italian journal of pure and applied mathematics.* 37(2017), 659-672.
- [7] O. Adeyeye and Z. Omar, Maximal order block method for the solution of second order ordinary differential equations, *IAENG Int. J. App. Math.*, 46(4)(2016), 46-4-03.
- [8] O. Adeyeye and Z. Omar, Solving third order ordinary differential equations using one-step block method with four equidistant generalized hybrid points, *IAENG Int. J. App. Math.*, 49(2)(2019), 42 56.
- [9] L. O. Adoghe, B. G. Ogunware and E. O. Omole, A family of symmetric implicit higher order methods for the solution of third order initial value problems in ordinary differential equations, *Theoretical Math. App.*, 6(3)(2016), 67-84.
- [10] O. A. Akinfenwa, R. I. Abdulganiy, B. I. Akinnukawe and S. A. Okunuga, Seventh order hybrid block method for solution of first order stiff systems of initial value problems, *J. Egypt. Math. Soc.*, 28(34)(2020), 28-34. <u>https://doi.org/10.1186/s42787-020-00095-3</u>.
- [11] O. A. Akinfenwa, R. I. Abdulganiy and S. A. Okunuga, A Simpsons 3/8 type block method for stiff systems of ordinary differential equation, Journal of the Nigerian Mathematical Society. 36(3)(2017), 503-514.
- [12] M. O. Alabi, M. T. Raji and M. S. Olaleye, Off-grid collocation four step initial value solver for second order ordinary differential equations, *International Journal of Mathematics and Statistics Studies*, 12 (4)(2024), 1-15
- [13] M. Al-Kandari, Enhanced criteria for detecting oscillations in neutral delay Emden Fowler differential equations, *Kuwait Journal of Science*, 50(4)(2023), 443 – 447.
- [14] A. M. Badmus, Y. A. Yahaya, and Y. C. Pam, Adams type hybrid block methods associated with Chebyshev polynomial for the solution ordinary differential equations, *British J. Math. Comp. Sci.*, 6(1)(2015), 464-474.
- [15] H. G. Debela and M. J. Kabeto, Numerical solution of fourth-order ordinary differential equations using fifth-order Runge–Kutta method, *Asian J. Sci. Tech.*, 8(2)(2017), 4332-4339.
- [16] A. B. Familua and E. O. Omole, Five points mono hybrid linear multistep method for solving nth order ordinary differential equations using power series function, *Asian Res. J. Math*, 3(1)(2017), 1-17.
- [17] J. Garba J. and U. Mohammed, Derivation of a new one-step numerical integrator with three intra-step points for solving first order ordinary differential equations. *Nig. J. Math. App.*, 30(2020), 155 172.
- [18] M. A. Islam, Accurate solutions of initial value problems for ordinary differential equations with the fourth order Runge-Kutta method, *J. Math. Res.*, 7(3)(2015), 41 53.
- [19] S. H. B. Kashkaria and M. I. Syamb, Optimization of one step block method with three hybrid points for solving first-order ordinary differential equations, *Results in Physics*, 12(1)(2019), 592–596.
- [20] M. M. Khalsaraei, A. Shokri, and M. Molayi, The new class of multistep multi-derivative hybrid methods for the numerical solution of chemical stiff systems of first order IVPs, *Journal of Mathematical Chemistry*, (2020). <u>https://doi.org/10.1007/s10910-020-01160-z</u>
- [21] K. A. Koroche, Numerical solution of first order ordinary differential equation by using Runge-Kutta method, *Int. J. sys. Sci. App. Math.*, 6(1)(2021), 1-8.
- [22] J. O. Kuboye and E. O. Adeyefa, New developed numerical formula for solution of first and higher order ordinary differential equations, *J. Interdisciplinary Math.* (2021). <u>https://doi.org/10.1080/09720502.2021.1925453</u>
- [23] J. O. Kuboye, O. R. Elusakin and O. F. Quadri, Numerical algorithm for direct solution of fourth-order ordinary differential equations, *J. Nig. Soc. Phys. Sci.*, 2(2020), 218-227.
- [24] J. A. Kwanamu, Y. Skwame, and J. Sabo, Block hybrid method for solving higher order ordinary differential equation using power series on implicit one-step second derivative, *FUW Trends in Sci. Tech. J.*, 6(2)(2021), 576 582.
- [25] A. I. Ma'ali, U. Mohammed, K. J. Audu, A. Yusuf and A. D. Abubakar, Extended block hybrid backward differentiation formula for second order fuzzy differential equations using Legendre polynomial as basis function, J. Sci., Tech., Math. Edu, 16(1)(2020), 100-111.
- [26] U. Mohammed, J. Garba and M. E. Semenov, One-step second derivative block intra-step method for stiff system of ordinary differential equations *J. Nig. Math. Soc.*, 40(1)(2021), 47-57.
- [27] U. Mohammed, O. Oyelami and M. E. Semenov, An orthogonal-based self-starting numerical integrator for second order initial and boundary value problem of ODEs, J. Phy. Conf. Ser. 1145(2019), 012040. Doi:10.1088/1742-6596/1145/1/012040.
- [28] A. Ndanusa and F. U. Tafida, Predictor-corrector methods of high order for numerical integration of initial value problems, *Int. J. Sci. and Inno. Math. Res*, 4(2)(2016), 47-55.



- [29] G. C. Nwachukwu and T. Okor Second derivative generalized backward differentiation formulae for solving stiff problems, *IAENG Int. J. App. Math.*, 48(1)(2018).
- [30] O. M. Ogunlaran and M. A. Kehinde, A new block integrator for second order initial value problems, *Neuroquantology*, 20(17)(2022), 1976 – 1980.
- [31] B. G. Ogunware, E. O. Omole and O. O. Olanegan, Hybrid and non-hybrid implicit schemes for solving third order odes using block method as predictors, *J. Math. Theory and Modelling*, 5(3)(2015), 10-25.
- [32] B. T. Olabode and A. L. Momoh, Continuous hybrid multistep methods with Legendre basis function for direct treatment of second order stiff ODEs, *Amer. J. Compt. App. Math.*, 6(2)(2016), 38-49.
- [33] O. O. Olanegan, B. G. Ogunware and C. O. Alakofa, Implicit hybrid points approach for solving general second order ordinary differential equations with initial values, *J. Advan. in Math. Comp. Sci.*, 27(3)(2018), 1-14.
- [34] O. O. Olanegan, B. G. Ogunware, E. O. Omole, T. S. Oyinloye and B. T. Enoch, Some variable hybrids linear multistep methods for solving first order ordinary differential equations using Taylor's series, *IOSR J. Math.* 11(2015), 08-13.
- [35] Z. Omar and R. Abdelrahim, New uniform order single step hybrid block method for solving second order ordinary differential equations, *Int. J. App. Engineering Res.*, 11(4)(2016), 2402-2406.
- [36] Z. Omar and O. Adeyeye, Numerical solution of first order initial value problems using a self-starting implicit twostep obrenchkoff-type block method, *J. Math. Stat.* 212(2)(2016), 127-134.
- [37] Z. Omar and J. O. Kuboye, Computation of an accurate implicit block method for solving third order ordinary differential equations directly, *Global J. Pure App. Math.*, 11(1)(2015), 177-186.
- [38] E. O. Omole and B. G. Ogunware, 3-point single hybrid block method (3PSHBM) for direct solution of general second order initial value problem of ordinary differential equations, *J. Sci. Res. Reports*, 20(3)(2018), 1-11.
- [39] E. O. Omole and L. A. Ukpebor, "A step by step guide on derivation and analysis of a new numerical method for solving fourth-order ordinary differential equations, *J. Math. Letter*, 6(2)(2020), 13-31.
- [40] G. F. Simmons, *Differential equations with applications and historical notes*, CRC Press, Taylor & Francis Group Boca Raton, London New York, 2017.
- [41] Y. Skwame, J. Z. Donald, J. Sabo, T. Y. Kyagya and A. A. Bambur, The numerical applications of implicit second derivative on second order initial value problems of ordinary differential equations, *Dutse J. Pure and Appl. Sci.*, 6(4)(2020), 1-14.
- [42] J. Sunday, A. A. James, M. R. Odekunle and A. O. Adesanya, Solution to free undamped and free damped motion problems in mass-spring systems, *Amer. J. Comput. App. Math.,* 6(2)(2016), 89-91.
- [43] A. T. Tiamiyu, A. T. Cole and K. J. Audu, A backward differentiation formula for fourth–order initial or boundary values problems using collocation method, *Iranian J. Opt*, 13(2)(2021).
- [44] P. Tumba, J. Sabo and M. Hamadina, Uniformly order eight implicit second derivative method for solving secondorder stiff ordinary differential equations *Acad. Res. Pub. Group*, 4(1)(2016), 43-48.
- [45] L. A. Ukpebor, E. O. Omole and L. O. Adoghe, An order four numerical scheme for fourth-order initial value problems using Lucas polynomial with application in ship dynamics, *International Journal of Mathematical Research*. 9(1)(2020), 28-41.
- [46] L. A. Ukpebor, Implicit four-step approach with application to non-linear third order ordinary differential equations, *FUDMA J. Sci. (FJS)*, 5(4)(2021), 406 412.
- [47] N. S. Yakusak and A. O. Owolanke, A class of linear multi-step method for direct solution of second order initial value problems in ordinary differential equations by collocation method, *J. Advances in Math. Comp. Sci.*, 26(1)(2018), 1-11.
- [48] A. T. Tiamiyu, K. I. Falade, Q. O. Rauf and S. A. Akande, A numerical technique for direct solution of special fourthorder ordinary differential equation via hybrid linear multistep method, *Cankaya University of Journal of Science and Engineering*, 18(1), (2021).
- [49] K. I. Falade, Numerical solution of higher-order singular initial value problems (SIVP) by exponentially fitted collocation approximate method, *American International Journal of Research in Science, Technology, Engineering & Mathematics*, 21(1)(2018).

Techno-Science Paper ID:1611387