



Original article

## ENHANCED DEGRADATION OF CRUDE OIL WITH *Alcaligenes faecalis* ADY25 AND IRON OXIDE NANOPARTICLE

\*Oyewole, O.A., Raji, R.O., Musa, I.O., Enemanna, C.E., Abdulsalam, O. N., Yakubu, J. G.

Department of Microbiology Federal University of Technology Minna Nigeria

Submitted: October, 2019; Accepted: November, 2019; Published: December, 2019

### ABSTRACT

Crude oil is composed of hydrocarbons, which forms a large group of chemicals that have caused a major concern to various forms of life. The advent of nanotechnology in conjunction with microbial cells produced nano-scale products with more efficient reactivity and larger surface area than its bulk phase facilitating the degradation of petroleum products. This study was conducted to determine the synergistic effect of *Alcaligenes faecalis* ADY25 and iron oxide nanoparticle on biodegradation of crude oil. The iron oxide nanoparticle was synthesized using corn silk extract. Pure isolate of *A. faecalis* ADY25 was collected from microbiology laboratory, Federal University of Technology Minna, Nigeria. The culture was grown on mineral salt medium containing crude oil as carbon source. This was supplemented with different amounts (0, 50, 100, 150, 200 mg) of iron oxide nanoparticles. The rate of biodegradation was determined by spectrophotometry and bacterial count was determined during biodegradation. The highest absorbance (1.418) and bacterial count ( $9.4 \times 10^6$  cfu/g) were obtained on day 9 and 12 by *A. faecalis* ADY25 supplemented with 200 mg of iron oxide nanoparticle. The results of this study showed that different amounts of iron oxide nanoparticle influenced the biodegradation of crude oil by *A. faecalis* ADY25.

**Key words:** *Alcaligenes faecalis* ADY25, crude oil, iron oxide, nanoparticles, biodegradation

\*Corresponding author: oa.oyewole@futminna.edu.ng, +234-8140031875

## INTRODUCTION

Crude oil also known as petroleum is composed of hydrocarbons, which forms a large group of chemicals that have caused a major concern to humans due to their widespread distribution into the environment causing harmful effects to various forms of life [1]. The presence of petroleum and its derivatives in the environment impair the ecosystem and human as well through the transfer of toxic organic materials including polycyclic aromatic hydrocarbons (PAHs) into the food chain [2]. Presence of polycyclic aromatic hydrocarbons (PAHs) in the environment has raised great concern to the health of humans since most of them are recalcitrant in nature. Petroleum and its derivatives being an important energy resources used by industries and in our daily life, at the same time, become a major pollutant of the environment [3]. Because of its complicated nature, petroleum can produce multiple types of toxic effects; it can cause acute lethal toxicity, sub-lethal chronic toxicity, or both depending on the dosage, and metabolism of the organism. Many petroleum components have the ability to bioaccumulate within susceptible aquatic organisms and can be transferred to other food chain levels by trophic transfer [4, 5].

As such, the importance of cleaning a contaminated petroleum environment has led to the development of techniques involving physical and chemical methods such as volatilization, photooxidation, chemical oxidation, and bioaccumulation that are rarely successful in rapid removal and clean-up of PAHs. However, physical and chemical methods to remove and clean PAHs are expensive, rarely successful, and produce toxic by-products [6, 7].

Nanotechnology is an emerging field to produce nano-scale products with more efficient reactivity and larger surface area than its bulk phase. These unique attributes of nanoparticles offer immense potential for their application to clean up hydrocarbons, pesticides and metals contaminated sites. The use of these nanoparticles in the physical and chemical methods of synthesizing iron oxide nanoparticles such as sol-gel, anodization, hydrothermal and microwave assisted process, have shortcomings such as long synthesis procedure, high energy requirement and production of toxic unreactive materials. Also, most of the available commercial reducing agents are expensive and toxic [8, 9, 10].

Microbial bioremediation simply involves the use of microorganisms such as bacteria, fungi and algae in the remediation of pollutants from the environment. Among the aforementioned microorganisms, bacteria have long been considered as one of the predominant hydrocarbons degrading agents found in the environment, which are free living and ubiquitous [6, 11]. Bioremediation has been drawing an increasing attention due to its economical, eco-friendly and self-propelling attributes as compared to the conventional physico-chemical methods of remediation of contaminated sites [6].

The success of bioremediation technologies applied to hydrocarbon-polluted environments highly depends on the biodegrading capabilities of native microbial populations or exogenous microorganisms used as inoculants [3]. The presence of microorganisms with the appropriate metabolic capabilities is the most important requirement for oil spill bioremediation. Unlike higher organisms,

Oyewole et al.

some microorganisms possess the catabolic capacity to use hydrocarbons as carbon and energy source. Bioremediation has been identified as a potential emerging technology for removing different spills, where physical washing and collection could not help. Highly hazardous oily materials can be mineralized to harmless products using suitable microorganisms [12]. The communities that were exposed to hydrocarbons become adapted, exhibiting selective enrichment and genetic changes. The adapted microbial communities can respond to the presence of hydrocarbon pollutants within hours and exhibit higher biodegradation rates than communities with no history of hydrocarbon contamination [13]. Crude oil biodegradation involves a mixture of various bacterial groups or consortia to degrade a wider range of hydrocarbons. The ability to isolate high numbers of certain oil-degrading microorganisms from an area is generally taken as evidence that these microorganisms are the most active oil degraders in such environment and can be used in the bioremediation of contaminated sites [14].

Nanoparticles (NPs), which is gradually gaining wide prominence in bioremediation can be applied directly for removal of organic contaminants through adsorption or chemical modification [9]. It can also serve as a facilitator in microbial remediation of contaminants either by enhancing the microbial growth or by immobilizing the remediating agents or through induced production of remediating microbial enzymes. Besides, nanoparticles induced enhanced production of biosurfactants in microorganisms, also contribute to improved solubility of hydrophobic hydrocarbons and thereby, create a conducive environment for microbial

International Journal of Applied Microbiology  
degradation of these contaminants in the environment [9].

Bacteria used in bioremediation are ubiquitous and readily available. Some bacteria used to commercial reducing agents can degrade crude oil compare to physical and chemical methods of degradation, and the method of synthesizing iron oxide nanoparticles is simple, cheap and eco-friendly and requires short reaction time compare to physical and chemical methods. In addition, Plant materials used for reduction, capping and stabilization of nanoparticles are in abundance. This technique provides a great advantage over chemical and physical methods since it is cost-effective, relatively reproducible and often result in more stable materials [15]. The aim of this study was to determine the synergistic effect of *Alcaligenes faecalis* ADY25 and iron oxide nanoparticle on biodegradation of crude oil.

## MATERIALS AND METHODS

### Sample Collection and Pre-treatment

Corn silk was randomly collected from different locations within Bosso, Minna, Nigeria. This was followed by removal of sands from the corn silk, and other debris after which the corn silks were air dried for three days at  $28^{\circ}\text{C} \pm 2$ .

### Collection of Bacterial Isolate

Pure isolate of *A. faecalis* ADY25 was collected from Microbiology laboratory, Federal University of Technology Minna, Nigeria. The bacterium was isolated from soil and was reported to have a huge potential for biosurfactant production [15]. The purity of the isolate was confirmed by streaking on nutrient agar plate and incubation at  $37^{\circ}\text{C}$  for 24 h.

Oyewole *et al.***Aqueous Extraction of Corn Silk**

An aqueous extract of corn silk was prepared by weighing 30 g of the corn silk into a 1000 ml beaker followed by addition of 500 ml of distilled water. The mixture was heated at 80°C for 30 min and a dark brown coloration of the aqueous solution was obtained. The extract was then allowed to cool to room temperature (28°C±2) and filtered using Whatman No1 filter paper and the quantity produced was measured using a measuring cylinder [16].

**Synthesis of Iron Oxide Nanoparticle**

The aqueous extract from a corn silk was added to 0.1 M of iron sulphate solution at the volume ratio of 2:1 in a sterile conical flask with constant stirring at room temperature (28°C±2) [17]. Then 1.0 M of NaOH solution was added to the above mixture to raise the pH (HANNA instrument) to 9 and the solution was heated at 60-65°C for 4 h and cooled. The resulting solution was centrifuged at 4000 rpm for 5 min and the dark greenish sediment was dried at 50°C using a hot air oven for 24 h and a brownish black residue of iron oxide was collected and kept in a universal bottle. The UV spectrophotometry of the synthesized iron oxide nanoparticles was carried out using UV-1800 series spectrophotometer at 200 to 800 nm

**Preparation of Mineral Salt Medium**

Mineral salt medium (MSM) was prepared using the following composition: K<sub>2</sub>HPO<sub>4</sub> - 1.8g, KH<sub>2</sub>PO<sub>4</sub> - 1.2g, NH<sub>4</sub>CL - 4.0g, NaCl<sub>2</sub> - 0.1g, FeSO<sub>4</sub>.7H<sub>2</sub>O - 0.01g. This was dissolved in 1000 mL distilled water in a conical flask [6].

**Biodegradation of Crude Oil using *A. faecalis* ADY25 and Iron Oxide Nanoparticles**

Nutrient broth was prepared and autoclaved at 121°C for 15 minutes, the broth was allowed to cool and 1 ml culture of *A. faecalis* ADY25 was inoculated inside 25 ml of the medium and incubated at 37°C for 24 hours in the incubator. Different amounts (50, 100, 150, 200 mg) of iron oxide nanoparticles were dissolved in four different conical flasks containing 50 ml of mineral salt medium. One milliliter (1 ml) of crude oil was introduced using a syringe into test tubes containing 9 mL of minimal salt medium with the different amounts of iron oxide nanoparticles after which it was autoclaved at 121°C for 15 minutes. The composition was allowed to cool and 0.1 mL of 24 h old broth culture of the organism was introduced in the test tubes, a control was set up without the organism and the absorbance was taken using a spectrophotometer (Model 752, China), every 3 days for 21 days.

**The Count of *A. faecalis* ADY25 during Biodegradation**

One milliliter of the 24 h old MSM broth culture of *A. faecalis* ADY25 was serially diluted tenfold, using a sterile distilled water. The 0.1 mL of 10<sup>-4</sup> dilution factor was placed on a sterile Petri dish and 15 ml of sterile Nutrient agar medium was poured on it. It was gently swirled and allowed to gel after which it was incubated at 37°C for 24 h and the count was taken and expressed as cfu/mL of the sample. This procedure continues every 3 days for 21 days.

**RESULTS**

The wet iron oxide nanoparticle and oven dried iron oxide nanoparticles is shown in Plates 1 and 2 while the UV spectrum of

Oyewole *et al.*

International Journal of Applied Biological Research, 2017

synthesized iron oxide nanoparticle is shown in Figure 1:




Plate 1. Wet iron oxide nanoparticle, Plate 2: Oven dried iron oxide nanoparticles




Figure 3: The effects of varying amount of iron-oxide nanoparticle on the count of *A. faecalis* ADY25 during crude oil degradation

**DISCUSSION**

*Alcaligenes faecalis* ADY25 used in this study is a Gram-negative rod previously identified as crude oil degrading bacterium with potential to degrade petrochemical products [18]. The isolate was able to degrade crude oil utilizing it as carbon source. The reason for this may be that the organism has enzyme system to degrade and utilize hydrocarbons and it is also capable of attaching to the substrate and produce biosurfactants, which help to break the oil into tiny micelle [15]. When the medium was supplemented with different amounts of iron oxide nanoparticles, an increase in the rate of degradation was observed. This indicates

the ability of the iron oxide nanoparticles to enhance the biodegradable ability of the isolate. Report from [19] indicates that nanoparticles are a supportive nutritional component, which influences lag phase and stationary phase and acts by reducing the duration of lag phase and increasing the duration of exponential and stationary phase. From the result, it was observed that the highest absorbance observed when 200 mg of iron oxide nanoparticles was used. This suggests that increase in amounts of iron oxide nanoparticles may increase the rate of biodegradation. This work is similar to that of [20] who obtained the best degradation using a microcosm containing 0.2 g of iron oxide nanoparticles and biosurfactant to degrade paraffins.

The organism was able to adapt and grow in the presence of crude oil multiplying in cell densities, however bacterial growth was best supported by 100 mg and 200 mg of iron oxide nanoparticle. The highest count ( $9.4 \times 10^6$  cfu/mL) was observed on the 18<sup>th</sup> day by cultures supplemented with 200 mg of iron oxide nanoparticle, this was followed by 100 mg ( $9.0 \times 10^6$  cfu/mL) after 12 days' incubation period. The test organism utilized crude oil as carbon and energy source and as the quantity of the crude oil decreases, microbial cell generation increases. This suggest that the test organisms utilize the crude oil for their growth and metabolism thereby removing the crude oil from the environment [21]. The observed growth and degradation potential of the isolate on the concentration of crude oil may be as a result of its previous exposure to pollutant. Furthermore, the growth of *A. faecalis* ADY25 in crude oil induced the secretion of extracellular protein and carbohydrate [22]. Previous studies have also shown hydrocarbon degrading bacteria to produce extracellular emulsifying agents generally consisting polysaccharide associated with protein. Some examples include alasan and emulsan, derived from *Acinetobacter*.

### CONCLUSION

This study showed the potential of *A. faecalis* ADY25 and iron oxide nanoparticles to degrade crude oil. Different amount of iron oxide nanoparticles (50, 100, 150 and 200 mg) enhance the process of degradation by the isolate. The use of the isolate and 200 mg of iron oxide nanoparticle to clean up crude oil contaminated environments is encouraged.

### REFERENCES

1. Al-Hawash, A. B., Li, S., Zhang, X., Zhang, X., & Ma, P. (2018). Productivity of  $\gamma$ -Linoleic acid by oleaginous fungus *Cunninghamella echinulata* using a pulsed high magnetic field. *Food Bioscience*, 21(5): 1-7.
2. Sel, A., & Fathepure, B. Z. (2009). Biodegradation of BTEX at high salinity by an enrichment culture from hypersaline sediments of Rozel Point at Great Salt Lake, *Journal of Applied Microbiology*, 107(6): 2001-2008.
3. Al-Wasify, R. S. & Hamed, S. R. (2016). Bacterial Biodegradation of Crude Oil Using Local Isolates. *International Journal of Bacteriology*, 2016, Article ID 6013871. Doi:<http://dx.doi.org/10.1155/2016/6013871>.
4. Aboelwafa, A.M. & Alwasify, R.S. (2009). Biodegradation of crude oil using local isolates. *Australian Journal of Basic and Applied Sciences*, 3(4): 4742-4751.
5. Almeda, R., Wambaugh, Z., Chai, C., Wang, Z., Liu, Z., & Buskey, E. J. (2013). Effects of crude oil exposure on bioaccumulation of polycyclic aromatic hydrocarbons and survival of adult and larval stages of gelatinous zooplankton. *PloS one*, 8(10): e74476. doi:10.1371/journal.pone.0074476.
6. Abioye, O. P., Agamuthu, P. & Abdulaziz, A. R. (2012). Biodegradation of Used Motor Oil in Soil Using Organic Waste Amendments. *Biotechnology Research International*. Doi:10.1155/2012/587041.

7. Ghosal, D., Ghosh, S., Dutta, T. K. & Ahn, Y. (2016). Current State of Knowledge in Microbial Degradation of Polycyclic Aromatic Hydrocarbons (PAHs): A Review. *Frontier in Microbiology*, 7, 1369. Doi:10.3389/fmicb.2016.01369.
8. Machado, S., Pacheco, J. G., Nouws, H. P. A., Albergaria, J. T., & Delerue-Matos, C. (2015). Characterization of green zero-valent iron nanoparticles produced with tree leaf extracts. *Science of the Total Environment*, 533, 76-81.
9. Kumari, K., & Singh, A. (2016). Genetically modified Cotton species detection by LISS-III satellite data. *Nature Precedings*, 34(4): 13-15.
10. Cecchin, I., Reddy, K. R., Thome, A., Tessaro, E. F., & Schnaid, F. (2017). Nanobioremediation: integration of nanoparticles, and bioremediation for the sustainable remediation of chlorinated organic contaminants in soils. *International Biodeterioration & Biodegradation*, 119, 419-428.
11. Dasgupta, D., Ghosh, R., & Sengupta, T. K. (2014). Biofilm-Mediated Enhanced Crude Oil Degradation by Newly Isolated Pseudomonas Species. *ISRN Biotechnology*, 6(5): 1-13.
12. Ismail, S., & Dadrasnia, A. (2015). Biotechnological Potential of Bacillus salmalaya 139SI: A Novel Strain for Remediating Water Polluted with Crude Oil Waste. *PLoS ONE*, 10(4): 34-35.
13. Shah, M., Fawcett, D., Sharma, S., Tripathy, K. S. & Poinern, G. E. J. (2015). Green Synthesis of Metallic Nanoparticles via Biological Entities. *Materials (Basel)*, 8(11): 7278-7308.
14. Al-Saleh, M. A., Ndip, J., & Akinjogunla, A. E. (2015). Impact of crude oil production on the petrochemical industry in Saudi Arabia. *Biotechnology and Biotechnological Equipment*, 25(5): 50-55.
15. Raji, R.O. & Oyewole, O.A. (2019). Isolation of Bacteria from Soil to Produce Biosurfactant. Book of Abstract of 40th Annual National Conference of the 42nd Annual Nigerian Society for Microbiology (NSM) Conference, 3rd - 6th September, 2019, Crawford University, Igbesa, Ogun State.
16. Hajinasiri, R., Norozi, B., & Ebrahimzadeh, H. (2016). Biosynthesis of ZnO Nanoparticles using corn silk of Zea mays L. Extract. *Chemistry Letters*, 45(11): 1238-1240.
17. Ali, A., Zafar, H., Zia, M., Haq, I., Phull, A. R., Ali, J. S. & Hussain, A. H. (2016). Synthesis, characterization, applications, and challenges of iron oxide nanoparticles. *Nanotechnology Science and Application*, 9, 49-67.
18. Asadpour, R., Sapari, N. B., Isa, M. H., Kakooei, S. (2019). Further Study of Adsorption of Crude Oils onto Acetylated Corn Silk and its Kinetics and Equilibrium Isotherm. *International Journal of Engineering* 32(2): 229-235.
19. Bhatia, M., Girdhar, A., Chandrakar, B., & Tiwari, A. (2013). Implicating nanoparticles as potential biodegradation enhancers: a review.

Oyewole et al.

*Journal of Nanomedicine and Nanotechnology*, 4(175): 2.

International Journal of Applied Biological Research 2019  
3, 159-169. DOI:  
10.3923/rjet.2009.159.169.

20. El-Sheshtawy, H. S., and Ahmed, W. (2017). Bioremediation of crude oil by *Bacillus licheniformis* in the presence of different concentration of nanoparticles and produced biosurfactant. *International Journal of Environmental Science and Technology*, 14(8): 1603-1614.

21. Igwo-Ezikpe, M.N., Gbenle, O.G., Ilori, M.O., Okpuzor J. & Osuntoki, A.A. (2009). Evaluation of *Alcaligenes faecalis* Degradation of Chrysene and Diesel Oil with Concomitant Production of Biosurfactant. *Research Journal of Environmental Toxicology*,

22. Alizadeh-Sani M., Hamishehkar H., Khezerlou A., Azizi-Lalabadi M., Azadi Y., Nattagh-Eshtivani E., Fashi M., Ghavami A., Aynehchi A. & Ehsani A. (2018). Bioemulsifiers derived from microorganisms: Applications in the drug and food industry. *Adv Pharm Bull.* 8(2): 191-199. DOI: [10.15171/apb.2018.023](https://doi.org/10.15171/apb.2018.023)

23. Ron E., Rosenberg E. (2010) *Acinetobacter and Alkanindiges*. In: Timmis K.N. (eds) *Handbook of Hydrocarbon and Lipid Microbiology*. Springer, Berlin, Heidelberg