



# ANIFS 2019

## THEME CAPTURE FISHERIES AND AQUACULTURE RESEARCH IN THE 21ST CENTURY NIGERIA'S ECONOMY

Proceedings of the 2nd Conference of  
Association of Nigerian Fisheries Scientists

July 9-11, 2019

at the Main Auditorium,

Abubakar Tafawa Balewa University, Sokoto

### EDITORS

Agbani, O.A., Ipinjoku, J.K., Ajan, P.K.,  
Magawata, I., Shinkafi, B.A., Ayoola, S.O.,  
Abubakar, M.Y. and Oribasoro, O.

Scanned with CamScanner

Dr. S. U. Ibrahim

442/570



# ANIFS 2019

Proceedings of the 2nd Conference of  
Association of Nigerian Fisheries Scientists  
is published annually

All correspondence to:

Prof. Fagbenro, O.A. (Editor)  
Department of Fisheries and Aquaculture Technology,  
Federal University of Technology, Akure

Prof. J.K. Ipinjolu (Editor)  
Department of Aquaculture and Fisheries Management,  
Usmanu Danfodiyo University, Sokoto

Prof. Ajani, E.K. (Editor)  
Department of Aquaculture and Fish Management,  
University of Ibadan, Ibadan

Prof. Magawata, I. (Editor)  
Department of Fisheries and Aquaculture,  
Usmanu Danfodiyo University, Sokoto

Prof. (Mrs) Shinkafi, B.A. (Editor)  
Department of Biological Sciences,  
Federal University, Gusau

Dr. Ayoola, S.O. (Editor)  
(Associate Professor)  
Department of Marine Sciences,  
University of Lagos, Lagos

Dr. Abubakar, M.Y. (Editor)  
Department of Aquaculture and Fisheries Management,  
Usmanu Danfodiyo University, Sokoto

Dr. Orisasona, O. (Editor)  
Department of Animal Science,  
Obafemi Awolowo, University, Ile-Ife

For further enquiries online,  
Please visit:  
[www.anifs.org.ng](http://www.anifs.org.ng)

ISBN: 978-978-58234-9-3 (Prints)  
978-978-58234-8-6 (Online)

All rights reserved (2019)

### Editorial Disclaimer

The description employed does not imply the expression of any opinion on the part of the Association of Nigerian Fisheries Scientist (ANIFS); authors are solely responsible for their articles in this publication. Information in this publication may be freely quoted or reprinted, but must be properly acknowledged together with a copy of the publication containing the quotation or reprint to be sent to the Secretariat of the Association.

This publication has been edited externally



USMANU DANFODIYO  
UNIVERSITY,  
SOKOTO



UMARU ALI SHINKAFI  
POLYTECHNIC,  
SOKOTO



Designed & Printed by: Artsmostfare Prints - Tel: 08028618360

Dr. S. U. Ibrahim

443/570





ANIFS2018



**PROCEEDINGS**  
OF 1ST CONFERENCE OF  
ASSOCIATION OF NIGERIAN  
FISHERIES SCIENTISTS



SCIENCE, INNOVATION AND AQUABUSINESS: A TRIPPOD FOR  
SUSTAINABLE FISHERIES AND AQUACULTURE DEVELOPMENT  
IN NIGERIA



## Publications of ANIFS

Access online at:  
[www.anifs.org.ng](http://www.anifs.org.ng)

Dr. S. U. Ibrahim

444/570

**Welcome Address by the Chief Host and Vice-Chancellor,  
Professor A. A. Zuru, at the Opening Ceremony of the Second  
Annual Conference of the Association of Nigerian Fisheries  
Scientists (ANIFS) held at the Main Auditorium,  
Usmanu Danfodiyo University, Sokoto, on 9 July, 2019.**

**Protocol**

I am very delighted to be here today for the opening ceremony of this epoch-making Conference of the Association of Nigerian Scientists (ANIFS). The association intends to create the avenue for scientists of like-minds to interact with one another. It is my pleasure to seize this opportunity to welcome you to the Seat of the Caliphate and to Usmanu Danfodiyo University, Sokoto. We take pride in the institution's uniqueness as the most peaceful University in Nigeria.

This Conference is a very distinctive one in the sense that it has gathered erudite scholars from different parts of Nigeria and the world over to interact and share ideas. The goal of this conference is simple goal towards growth and improvement of fisheries resources. And that is why it is themed 'Capture Fisheries and Aquaculture Research in the 21st Century Nigeria'.

Your choice of this University as the host of this Conference could not have been better. The contribution of this University to the development of fisheries and aquaculture as the pioneer university to kick off a programme in the field in the whole of Northern Nigeria. Since then, the institution has produced professionals and erudite scholars who are greatly contributing to national development. Against this background, the hosting right is well deserved.

We give you the assurance of continuous commitment to excellence, quality programmes and researches that are intended to greatly impact the potential of our younger ones through training and research for growth and development of Nigeria.

I understand that this association is a very young one. Notwithstanding, I want you to hold on to the tenets of what it stands for and never derail from the excellent standards you have set and always work towards. We are ready as a University to support the growth of any association with progressive goals like yours.

The report I received from the Maiden Edition proved beyond doubt that the contributions of our scholars towards the growth and expansion of fisheries and aquaculture in Nigeria cannot be over-emphasized. Nevertheless, do not rest on your efforts.

Permit me to use this opportunity to share an observation I noticed here in the North-West zone of Northern Nigeria generally. The activities of cattle rustlers and bandits in Zamfara and Katsina axis have significantly caused huge depletion in the supply of terrestrial animal protein. A development that has made it increasingly less affordable to the populace. This scenario, coupled with the dwindling promise of the capture fisheries in the available freshwater environments, constitutes a serious danger to our food security efforts. Personally, I think this situation has not only made speedy popularization and simplification of the aquaculture techniques important but the obvious solution. ANIFS may wish to take this challenge seriously and head-on. Once again, on behalf of the University and the entire State, I deeply appreciate the hosting right privilege and wish the participants a successful, memorable and fruitful deliberations. I also wish you safe return to your various destinations after the Conference.

Professor A. A. Zuru  
Vice-Chancellor  
Usmanu Danfodiyo University, Sokoto

Dr. S. U. Ibrahim

J. MAMMAL  
15/07/2019  
445/570

**Address by the President of the Association of Nigerian Fisheries Scientists (ANIFS), Professor J. K. Ipinjolu, at the Opening Ceremony of the Second Annual Conference held July 9, 2019 at the Main Auditorium, Usmanu Danfodiyo University, Sokoto**

**Protocol**

On behalf of the National Executive Council (NEC) of the Association of Nigerian Fisheries Scientists (ANIFS), I humbly welcome you all to the opening ceremony of the Second Annual Conference of the Association.

The Association is a body of scientists in Capture Fisheries and Aquaculture discipline who are committed to the advancement of sustainable fish production through research, innovation and community service. Kindly visit our website ([www.anifs.org.ng](http://www.anifs.org.ng)) for further information on the goal, mission, objectives and activities of the Association.

The first conference of the Association was held precisely one year ago (10-12 July, 2018) at the University of Ibadan, Ibadan. The focus of the maiden Conference was on **Science, Innovation and Aquabusiness: A Tripod for Sustainable Fisheries and Aquaculture Development in Nigeria**. The contributions of the conference are already in circulation in form of Proceedings and a Journal.

In line with the mission of the Association, this 2<sup>nd</sup> conference has been designed to focus on **Capture Fisheries and Aquaculture Research in the 21st Century Nigerian Economy**. Available statistics (FDF, 2015) shows that the country's fisheries sub-sector contributes about 0.88% to the Agricultural GDP, employs about 8.63 million and 19.55million Nigerians in its primary and secondary sectors, respectively. However, a huge annual import bill of about \$1.13 billion is incurred on importation of about 806,000 metric tons of assorted fish products. The country is able to exploit only 32.6% of the estimated potential fish yield from all sectors, and only 12.7% in the aquaculture sector. Consequently, with total domestic production of only 31.6% of the demand, the country has to battle with a fish production deficit of about 2.22 million metric tonnes. Therefore, there is an urgent need to step-up current research efforts and innovation in all areas of the fisheries sector in order to enhance sustainable fish production and improve human nutrition, livelihoods and food security.

The keynote speaker for the topical issue of this conference is a renowned scholar and fisheries scientist, Professor S. J. Oniye of the Department of Zoology, Ahmadu Bello University, Zaria. The technical sessions will look at scientific findings and technologies in fifty nine research papers distributed as follows:

Sub-Theme	Percentage
Fish Nutrition and Feed Technology	24
Fish Genetics, Breeding and Biotechnology	17
Ecology and Management of Fisheries in Wetlands	15
Fish Health-Management	15
Socio-economic, Policy and Gender Issue	12
Fish Processing and Product Development	9
Other Sub-themes	9

The Association is always ready to partner with agencies of government and the private sector on their fisheries activities. In this Conference, our team of scientists and professionals will meet with representatives of such agencies, fish farms, fish processors and other stakeholders immediately after the opening ceremony.

Dr. S. U. Ibrahim  
vi

446/570

**Address by the President of the Association of Nigerian Fisheries Scientists (ANIFS), Professor J. K. Ipinjolu, at the Opening Ceremony of the Second Annual Conference held July 9, 2019 at the Main Auditorium, Usmanu Danfodiyo University, Sokoto**

**Protocol**

On behalf of the National Executive Council (NEC) of the Association of Nigerian Fisheries Scientists (ANIFS), I humbly welcome you all to the opening ceremony of the Second Annual Conference of the Association.

The Association is a body of scientists in Capture Fisheries and Aquaculture discipline who are committed to the advancement of sustainable fish production through research, innovation and community service. Kindly visit our website ([www.anifs.org.ng](http://www.anifs.org.ng)) for further information on the goal, mission, objectives and activities of the Association.

The first conference of the Association was held precisely one year ago (10-12 July, 2018) at the University of Ibadan, Ibadan. The focus of the maiden Conference was on **Science, Innovation and Aquabusiness: A Tripod for Sustainable Fisheries and Aquaculture Development in Nigeria**. The contributions of the conference are already in circulation in form of Proceedings and a Journal.

In line with the mission of the Association, this 2<sup>nd</sup> conference has been designed to focus on **Capture Fisheries and Aquaculture Research in the 21st Century Nigerian Economy**. Available statistics (FDF, 2015) shows that the country's fisheries sub-sector contributes about 0.88% to the Agricultural GDP, employs about 8.63 million and 19.55 million Nigerians in its primary and secondary sectors, respectively. However, a huge annual import bill of about \$1.13 billion is incurred on importation of about 806,000 metric tons of assorted fish products. The country is able to exploit only 32.6% of the estimated potential fish yield from all sectors, and only 12.7% in the aquaculture sector. Consequently, with total domestic production of only 31.6% of the demand, the country has to battle with a fish production deficit of about 2.22 million metric tonnes. Therefore, there is an urgent need to step-up current research efforts and innovation in all areas of the fisheries sector in order to enhance sustainable fish production and improve human nutrition, livelihoods and food security.

The keynote speaker for the topical issue of this conference is a renowned scholar and fisheries scientist, Professor S. J. Oniye of the Department of Zoology, Ahmadu Bello University, Zaria. The technical sessions will look at scientific findings and technologies in fifty nine research papers distributed as follows:

Sub-Theme	Percentage
Fish Nutrition and Feed Technology	24
Fish Genetics, Breeding and Biotechnology	17
Ecology and Management of Fisheries in Wetlands	15
Fish Health-Management	15
Socio-economic, Policy and Gender Issue	12
Fish Processing and Product Development	9
Other Sub-themes	9

The Association is always ready to partner with agencies of government and the private sector on their fisheries activities. In this Conference, our team of scientists and professionals will meet with representatives of such agencies, fish farms, fish processors and other stakeholders immediately after the opening ceremony.

Dr S. U Ibrahim

447/570

### Appreciation

I seize this opportunity, on behalf of the NEC and the entire members of the Association to express our appreciation to organisations and individuals who have contributed to the hosting of this Conference. Our special thanks to the Chief Host and Vice-Chancellor, Usmanu Danfodiyo University, Sokoto, Professor A. A. Zuru, for accepting to host this Conference and for providing support in cash and kind. We are also grateful for the cash donations from the Dean, Prof. B. Z. Abubakar, and all the Departments in the Faculty of Agriculture, UDU, Sokoto; Shehu Shagari College of Education, Sokoto; Sokoto State Polytechnic; and Professor (Mrs) B. A. Shinkafi, Deputy-Vice-Chancellor, Federal University Gusau. We equally acknowledge the commitment and sacrifices of the Head and members of staff of the Department of Fisheries and Aquaculture as well members of the Local Organising Committee drawn across the Faculty of Agriculture, UDU, Sokoto.

We thank our invited Guests of Honour and the Royal Father of the day, HRH Alhaji Samaila Mohammad Mera (CON), who is also a Patron of the Association for finding time to honour our invitation despite their tight schedules. We also appreciate the presence of the Deans and Directors and members of the University community.

Thank you.

**Prof. J. K. Ipinjolu**  
President, ANIFS

Dr S. u Ibrahim

# Contents

Welcome Address by the Chief Host and Vice-Chancellor, Professor A. A. Zuru, at the Opening Ceremony of the Second Annual Conference of the Association of Nigerian Fisheries Scientists (ANIFS) held at the Main Auditorium, Usmanu Danfodiyo University, Sokoto, on 9 July, 2019.....	v
Address by the President of the Association of Nigerian Fisheries Scientists (ANIFS), Professor J. K. Ipinjolu, at the Opening Ceremony of the Second Annual Conference held July 9, 2019 at the Main Auditorium, Usmanu Danfodiyo University, Sokoto.....	vi
Logos of Four Institutions that supported the Conference.....	viii

## Ecology and Management of Fisheries in Wetlands

Diversity and Morphometric Variables of Fish Caught from River Rima in Sokoto State, Nigeria Magami, I. M., Usman, R., and Haidara, A. M.....	3
Some Biological Parameters of <i>Hepsetus odoe</i> from Aiba Reservoir, South-Western Nigeria Ipinmoroti, M.O. and Oyewusi, M.A.....	11
Influence of Weight and Sex on Intestinal Bacteria of Wild <i>Clarias gariepinus</i> (Burchell, 1822) from River Rima in Sokoto State, Nigeria Yusuf, M. A., Obaroh, I.O., Magawata, I., and Sadauki, M.A.....	19
Levels of some Chemical Parameters of Three Water Bodies in Bakolori Irrigation Project, Zamfara State, North Western Nigeria Aminu, M. U., Ipinjolu, J. K. and Shinkafi, B. A.....	27
Length-Weight Relationship and Condition Factor of Commercially Important Fish Species of Usuma River in Gwagwalada, FCT, Nigeria Dan-kishiya, A. S., Omeiza, E. O. and Idowu, R. T.....	37
Assessment of some Physico-Chemical Parameters of Tagwai Reservoir in Minna, Niger State, Nigeria Ibrahim, S.U., Umar, F., Yakubu, U. P., Adama, B. S. and Olubobokun, A. E.....	45
Seasonal Variation in Abundance and Condition Factor of Fish Species in Osun River, Nigeria Iyiola, A. O. and Jenyo-Oni, A.....	51

## Advances in Fish Genetics, Breeding and Biotech

Genetic Variability in Wild Populations of <i>Clarias Gariepinus</i> using Random Amplified Polymorphic DNA (RAPD) Fingerprinting Technique Umar, B. D., Suléiman, B. and Balogun, J. K.....	61
Influence of Age of Donor Fish on the Success of Hypophysation in <i>Clarias gariepinus</i> Egwenomhe, M., Igbo, N. U. and Osaigbovo, P. O.....	69
Effect of Ovulin Hormone Suspended in Saline on Induced Spawning Performance of African Catfishes ( <i>Clarias anguillaris</i> and <i>Clarias gariepinus</i> ) Maradun, H. F., Argungu, L. A., Abubakar, M. Y., Umar, F., Ahmad, M., and Sahabi, A. M.....	75
Reproductive Indices of <i>Clarias gariepinus</i> Broodstock Fed Different Inclusion Levels of Composite Medicinal Plant Powder Onyia, L. U., Ochokwu, I. J., and Jampeo, J. J.....	85

Dr S. U. Ibrahim

ix

449/570

## Fish Nutrition and Field Technology

Growth Response and Feed Utilization of *Heterobranchius longifilis* Fingerlings Fed Diets Fortified with

*Lactobacillus paracasei*

Nghede, O. B., Ajani, E. K. and Soetan, K. O.....

Effect of Black Cumin (*Nigella sativa*) Seed Extracts on Growth Performance of *Coptodon zillii*

Sani, K.A., Obaroh, I.O., Adamu, I., Joseph, E.K., Arilewo, F.V. and Hafsai, A.I.....

Evaluation of Fermented *Mucuna pruriens* Leaf Meal as an Alternative Soybean Meal Diet *Clarias gariepinus*

(Burchell 1822) Fingerlings

Ibrahim, R., Gana, A.B., Iriobe, T., Jega, I.S., Ahmad, S.M. and Inuwa, M.B.....

Growth Performance of Sex Reversed Tilapia *Oreochromis niloticus* Fed Probiotics Based Diets

Kareem, O. K., Ajao, F. S., and Oyenekan, S. I.....

Effects of Honey on Survival and Growth Performance of *Clarias gariepinus* (Burchell, 1822) Fry

Ramayi, C., Balogun, J. K. and Auta, J.....

Haematological Parameters of Juvenile Catfish (*Clarias gariepinus*) Fed Processed Catfish Offal Diet

Iriobe, T., Ajani, E. K., Ibrahim, R., Gana, A. B., and Jega, I. S.....

Growth Response and Carcass Composition of Catfish (*Clarias gariepinus*) Fingerlings Fed Varying

Inclusion Levels of Toasted Wild Groundnut (*Calapogonium muccunoides*) Seed Meal

Bake, G. G., Oghonnaya, C. C., Yusuf, J., Adam, A., Gana, A. B., Nwangwu, D. C. and Sadiku, S. O. E.....

Growth Response of *Clarias gariepinus* Juveniles Fed Diets Containing Varying Inclusion Levels of Fresh

and Dried Maggot (*Musca domestica*) Meals

Akanmu, O. A., Ipinmoroti, M. O., Iyiola, A., Fayemi, O., J. A., Ayanboye, A. O. and Kolawole, A. S.....

Growth Performance of *Clarias gariepinus* Juveniles Fed Diets Incorporated with Graded Levels of Roasted

Cashew (*Anacardium occidentale*) Nut Meal

Adesina, S.A. and Amoo, B.O.....

Evaluation of Grasshopper Meal as Alternative Protein Source in the Diet of Nile Tilapia

(*Oreochromis niloticus*)

Mamman, T., Ipinjolu, J. K. and Hassan, W.A.....

## Fish Processing and Product Development

Effect of Silica Gel and Storage Media on the Sensory and Bacteria Attributes of Smoked Catfish

(*Heterobranchius bidorsalis* Geoffroy Saint Hilaire 1809)

Sahabi, A. M., Magawata, I., Argungu, L.A. and Rabah, A.B.....

Effect of Silica Gel and Storage Media on Proximate Composition of Smoked Catfish (*Heterobranchius*

*bidorsalis* Geoffroy Saint Hilaire 1809)

Sahabi, A. M., Magawata, I. and Abubakar, M. Y.....

Effects of Processing Methods on Nutritional Profile of African Locust Bean (*Parkia biglobosa*) Seeds

Abdullahi, U. U., Garba, B., Mairo, A., Ndako, A. M. and Abraham, O. V.....

Sensory and Chemical Characteristics of *chinchin* made using Wheat Flour and Fish (*Protopterus*

*annectens*, Owen 1839) Powder

Magawata, I., Umar, F. and Makusidi, F.I.....

Dr S.U. Ibrahim

450/570

x

## Socio-Economic, Policy and Gender Issues in Cultures and Capture Fisheries

Constraints to Fish Marketing in Gombe State, Nigeria Tasir, R. W., Ezekail, B. and Akinbote, R.....	205
Utilization of ICTs by Fish Farmers in Bayelsa State, Nigeria Ogunu, Eboye, U. G. ....	211
Fishing Gear and Status of Fisheries of Forcados in Burutu Local Government Area, Delta State, Nigeria Abolagba, O. J., Oghoru, R.O. and Idele, E.O.....	215
Demographic Characteristics and Potentials of Fish Farming in Ibi Local Government Area, Taraba State, Nigeria Ogunremi, J. B., Igbani, F., Uruku, N. M. and Shetur, C. Y.....	223
Profitability and Efficiency of Dried Fish Marketing in Sokoto North Local Government Area, Nigeria Obalola, T.O., Abubakar, B.B., Agboola, B.O., Odum, E.B.E., Sadauki, M.A and Yusuf, M.A.....	229

## Fish Health Management and Environmental Toxicology

Isolation of <i>Citrobacter freundii</i> from Farmed Catfish in Lagos State, Nigeria Akinlayo, I. A., Ezeri, G. N. O., Bankole, M. O., Akinyemi, A. A. and Adeosun, F.I.....	237
Probit Modelling for LC <sub>50</sub> and LC <sub>90</sub> Determination Using Neem Seed Extract and Moringa Root Bark on <i>Clarias gariepinus</i> (Burchell 1822) Juveniles Shamaki, S.B. and Oladimeji, A.O.....	243
Toxicity Effect of Urea Fertilizer on the Juveniles of Wild <i>Clarias gariepinus</i> (Burchell 1822) in Sokoto, Nigeria Argungu, L. A., Magami, I. M., Hudu, A. and Yusuf, M. A.....	249

Dr S. U Ibrahim

451/570


# Ecology and Management of Fisheries in Wetlands



**ANIFS2019**  
ASSOCIATION OF NIGERIAN FISHERIES SCIENTISTS  
[www.anifs.org.ng](http://www.anifs.org.ng)

452/570

S. Y. Ibrahim



## Assessment of some Physico-Chemical Parameters of Tagwai Reservoir in Minna, Niger State, Nigeria

Ibrahim, S.U., Umar, F\*, Yakubu, U. P., Adama, B. S. and Olubobokun, A. E.  
Department of Water Resources, Aquaculture and Fisheries Technology, Federal University of Technology, Minna, Niger State.  
Department of Fisheries and Aquaculture, Usmanu Danfodiyo University, Sokoto.

\*Corresponding author E-mail:  
ulruk915@gmail.com

### Abstract

This study on the assessment of physico-chemical parameters of Tagwai Dam in Minna, Niger State, Nigeria, was carried out from February 2016 to July 2016. Five sampling stations were randomly selected, namely; Tasabo (Station 1), Maigan-ga (Station 2), Lokoto (Station 3), Dutsen-kura (Station 4), and the dam site named Gmukpa (Station 5). The physico-chemical parameters were determined using standard methods, procedures, and instruments. The result revealed that; Water temperature ( $26.36 \pm 0.70 - 32.40 \pm 4.17^\circ\text{C}$ ), pH ( $6.53 \pm 0.20 - 7.91 \pm 0.11$ ), Transparency ( $99.60 \pm 2.70 - 135.40 \pm 25.25$ ), Conductivity ( $74.60 \pm 1.14 - 91.60 \pm 1.67 \mu\text{S/cm}$ ), Total Dissolved Solids ( $0.09 \pm 0.03 - 0.26 \pm 0.38 \text{mg/L}$ ), Nitrate-nitrogen ( $0.16 \pm 0.03 - 0.43 \pm 0.17 \text{mg/L}$ ), Total hardness ( $31.46 \pm 1.64 - 43.38 \pm 0.85 \text{mg/L CaCO}_3$ ), Dissolved Oxygen ( $5.76 \pm 2.10 - 13.60 \pm 3.85 \text{mg/L}$ ), Phosphate-phosphorus ( $6.53 \pm 0.20 - 0.30 \pm 0.15 \text{mg/L}$ ), and water depth ( $6.38 \pm 2.75 - 11.18 \pm 4.80 \text{m}$ ) varied with months and seasons. Analysis of variance showed significant difference between seasons ( $P < 0.05$ ). Water quality of the dam is influenced by anthropogenic activities such as runoffs of inorganic fertilizers and pesticides. Dam water is suitable for irrigational and domestic purposes as indicated by most of the physico-chemical parameters analysed in this research. Hence, there is the need for effective anthropogenic inputs control programme in the dam.

Keywords:  
Physico-chemical parameters,  
Tagwai Reservoir,  
Anthropogenic activities.

### Introduction

Nigeria is blessed with about 853,600 hectares of freshwater capable of producing over 1.5 million metric tonnes of fish annually (FAO, 2009). There is the need to exploit means of using these precious resources even though there are some hindrances, which include effects of domestic and agricultural wastes on the water quality and aquatic life, physical and chemical factors like temperature, turbidity, pH, dissolved gases and carbon dioxide, salts and nutrients. It is no doubt that dams have contributed to the economic growth of many nations with Nigeria inclusive.

Dams built in several parts of the world have played important role in helping communities to harness water resources for several uses. An estimated 30-40% of irrigated land worldwide now relies on reservoir water (Mustapha, 2011). In Nigeria, many researchers have conducted works on different water bodies, some of them include, Balogun *et al.* (2005) on some aspects of the limnology of Makwaye Lake in Ahmadu Bello University Farm, Samaru, Zaria; Balarabe (2001), on effect of limnological characteristic on zooplankton composition and distribution in Dumbi and Kwangila ponds, Zaria; Ibrahim *et al.* (2009) on an assessment of the physico-chemical parameters of Kontagora reservoir, Niger State; Hassan *et al.* (2010) on the algal diversity in relation to physico-chemical parameters of three ponds in Kano metropolis and Abubakar (2009) on the limnological studies for the assessment of Sabke Lake, Katsina State. This study however established physical, chemical, and biological parameters of Tagwai dam, and provided better understanding of the dam's ecosystem.

Dr S. U Ibrahim

453/570

### Materials and Methods

**Study Area**  
Tagwai dam is located in Chanchaga Local Government Area in southwest zone of Minna. The dam is at the east of Tungan Goro about 10km southeast of mobile market and northeast of Paiko. It is an earthen dam constructed in 1978 by the Kano State Water Resources Engineering Construction Agency (NSWB, 1991). The dam lies on latitude  $9^{\circ} 34' N$  and longitude  $6^{\circ} 39' E$ . Fishing activities are carried out on the water body and also serves as a primary reservoir for the city of Minna under the supervision of Niger State Water Board (NSWB).




Figure 1: The geographical location of the study area showing Tagwai Reservoir




Figure 2: The google image of the study area showing the sampling stations

Dr S.U. Ibrahim

454/570

**Water Sampling and Analysis**

Water samples were collected from each station once every month for six months. 100ml dissolved oxygen bottle was used for dissolved oxygen and one litre plastic bottle was also used for the analysis of other water quality parameters which include; conductivity, alkalinity, pH, water temperature. The parameters were analysed at the Water Resources, Aquaculture and Fisheries Technology Laboratory of the Federal University of Technology, Minna. Dissolved oxygen was determined using the Winklers method. The pH of the water samples was determined using a B. Brn Scientific pH- meter (pHS-25). Conductivity was determined using the Jenway 4010 Conductivity Meter. Other methods used in water sampling, preservation and analysis were those prescribed by APHA (2005).

Water temperature (°C) was measured by dipping the echo sounder (Garmin Fish Finder 400C) into the water for about 1-2minutes so as to know the temperature alongside the depth at each station. The turbidity of water was measured with turbidity tube. The tube was calibrated at the bottom with "X" mark in black colour. The water sample was measured in 200ml beaker and poured gradually into the turbidity tube while at the same time observing the calibration mark at the bottom of the tube from the upper side of the tube until the calibrated line disappeared. The depth at which it disappeared was recorded in Nephelometric Turbidity Unit (NTU) from the graduated readings of the turbidity tube (Nathanson, 2003).

**Statistical Analysis**

The statistical analysis of the samples was carried out using the one way analysis of variance (ANOVA) to determine the variation of parameters within the months and stations. The Duncan multiple range test for the separation of mean was used to compare parameters between the months and stations.

**Results**

The physico-chemical parameters of the dam water (Table 1) revealed stations mean variation of Tagwai Dam, with respect to water temperature, dissolved oxygen, pH, alkalinity and conductivity. Table 2 shows the mean monthly variation of the parameters. The water temperature monthly variation ranged from 26.36±0.70° to 30.32±0.81° °C; the pH values ranged between 6.53±0.20° and 7.91±0.11°; transparency of the reservoir fluctuated with mean value of 99.60 ± 2.70° to 135.40 ± 25.25°. The dissolved oxygen values in the reservoir ranged from 5.76 ± 2.10° to 13.60 ± 3.85°. The electrical conductivity ranged from 74.60±1.14° to 91.60±1.67°. The hardness of the dam water ranged from 36.92±1.21° to 43.38±0.85°; Nitrate-nitrogen indicated the mean value ranged from 0.16±0.03° - 0.43±0.17° during the period of study. Total dissolved solids in the dam had the peak value of 0.26±0.38° which was recorded in the month of July while the least value of 0.09±0.03° was also recorded in the month of June; and the mean value of Phosphate-phosphorus ranged from 0.10±0.01° - 0.13±0.05°. The mean value of depth ranged from 6.38±2.75° - 11.18±4.80°.

Table 1: Physico-chemical parameters of Tagwai Reservoir

Parameters	ST1	ST2	ST3	ST4	ST5
DO (mg/l)	9.00±3.03 <sup>bc</sup>	10.40±4.57 <sup>c</sup>	5.33±1.54 <sup>a</sup>	9.33±3.72 <sup>bc</sup>	7.83±2.56 <sup>b</sup>
Conductivity (µS/cm)	86.67±6.25 <sup>a</sup>	87.17±7.08 <sup>a</sup>	88.67±6.95 <sup>b</sup>	87.67±6.38 <sup>ab</sup>	86.67±5.35 <sup>a</sup>
TDS (mg/l)	0.14±0.03 <sup>a</sup>	0.23±0.35 <sup>a</sup>	0.12±0.02 <sup>a</sup>	0.08±0.03 <sup>a</sup>	0.13±0.02 <sup>a</sup>
Transparency (cm)	117.33±28.75 <sup>a</sup>	110.50±12.19 <sup>a</sup>	109.00±13.18 <sup>a</sup>	106.00±2.00 <sup>a</sup>	109.83±9.17 <sup>a</sup>
pH	7.19±0.61 <sup>a</sup>	7.02±0.65 <sup>a</sup>	7.06±0.60 <sup>a</sup>	7.18±0.66 <sup>a</sup>	7.20±0.64 <sup>a</sup>
Nitrate	0.29±0.12 <sup>ab</sup>	0.21±0.05 <sup>a</sup>	0.30±0.09 <sup>b</sup>	0.34±0.16 <sup>b</sup>	0.32±0.13 <sup>b</sup>
Phosphate	0.19±0.16 <sup>a</sup>	0.15±0.07 <sup>a</sup>	0.18±0.11 <sup>a</sup>	0.13±0.03 <sup>a</sup>	0.17±0.06 <sup>a</sup>
Total Alkalinity (mg/l)	23.35±9.20 <sup>a</sup>	22.63±6.62 <sup>a</sup>	23.00±7.46 <sup>a</sup>	23.92±9.90 <sup>a</sup>	23.00±5.45 <sup>a</sup>
Total hardness (mg/l)	38.63±3.30 <sup>a</sup>	38.63±4.70 <sup>a</sup>	37.63±4.90 <sup>a</sup>	36.88±4.47 <sup>a</sup>	38.03±4.29 <sup>a</sup>
Temperature (°C)	29.90±3.52 <sup>a</sup>	28.90±5.54 <sup>a</sup>	29.07±2.18 <sup>a</sup>	27.05±2.07 <sup>a</sup>	28.02±2.05 <sup>a</sup>
Depth (m)	5.33±1.20 <sup>a</sup>	4.35±0.91 <sup>a</sup>	13.11±3.39 <sup>d</sup>	7.52±2.55 <sup>b</sup>	10.33±2.98 <sup>c</sup>

Dr. Sul Ibrahim

455/570

Table 2: Monthly variation of water physico-chemical parameters of Tagwai Reservoir

Parameters	February	March	April	May	June	July
DO(mg/l)	8.60±1.95 <sup>b</sup>	7.60±1.82 <sup>ab</sup>	5.76±2.10 <sup>a</sup>	8.80±1.03 <sup>b</sup>	13.60±3.85 <sup>c</sup>	5.92±1.34 <sup>a</sup>
Conductivity(µS/cm)	91.60±1.67 <sup>c</sup>	90.80±1.48 <sup>bc</sup>	89.20±0.84 <sup>bc</sup>	89.80±1.30 <sup>bc</sup>	88.20±0.84 <sup>b</sup>	74.60±1.14 <sup>a</sup>
TDS (mg/l)	0.15±0.04 <sup>a</sup>	0.13±0.03 <sup>a</sup>	0.11±0.03 <sup>a</sup>	0.10±0.03 <sup>a</sup>	0.09±0.03 <sup>a</sup>	0.26±0.38 <sup>a</sup>
Transparency(cm)	135.40±25.25 <sup>ab</sup>	108.40±2.30 <sup>a</sup>	108.00±1.41 <sup>a</sup>	107.40±0.55 <sup>a</sup>	104.40±1.82 <sup>a</sup>	99.60±2.70 <sup>a</sup>
pH	7.04±0.30 <sup>b</sup>	7.76±0.47 <sup>c</sup>	7.91±0.11 <sup>c</sup>	6.53±0.20 <sup>a</sup>	6.99±0.47 <sup>b</sup>	6.55±0.06 <sup>a</sup>
Nitrate	0.21±0.02 <sup>ab</sup>	0.43±0.17 <sup>d</sup>	0.35±0.09 <sup>cd</sup>	0.32±0.03 <sup>c</sup>	0.29±0.03 <sup>bc</sup>	0.16±0.03 <sup>a</sup>
Phosphate	0.10±0.03 <sup>a</sup>	0.10±0.01 <sup>a</sup>	0.19±0.05 <sup>a</sup>	0.14±0.03 <sup>a</sup>	0.15±0.03 <sup>a</sup>	0.30±0.15 <sup>b</sup>
T. Alkalinity(mg/l)	21.62±0.42 <sup>b</sup>	18.86±0.94 <sup>a</sup>	19.26±0.48 <sup>ab</sup>	20.64±0.21 <sup>ab</sup>	19.90±0.60 <sup>ab</sup>	38.80±4.15 <sup>c</sup>
T. Hardness(mg/l)	41.00±1.95 <sup>c</sup>	31.46±1.64 <sup>a</sup>	43.38±0.85 <sup>d</sup>	36.92±1.21 <sup>b</sup>	37.82±1.21 <sup>b</sup>	37.20±2.70 <sup>b</sup>
Temperature(°C)	27.40±1.53 <sup>ab</sup>	26.36±0.70 <sup>a</sup>	28.60±4.55 <sup>ab</sup>	30.32±0.81 <sup>bc</sup>	32.40±4.17 <sup>c</sup>	26.44±0.55 <sup>a</sup>
Depth(m)	6.38±2.75 <sup>a</sup>	6.46±2.82 <sup>a</sup>	6.38±2.75 <sup>a</sup>	7.92±3.52 <sup>a</sup>	10.44±5.16 <sup>b</sup>	11.18±4.80 <sup>b</sup>

### Discussion

The water temperature of the reservoir fluctuated within months, which was between 26.36°C and 32.40°C. The low water temperature recorded in the reservoir was in the dry season, which could be as a result of seasonal changes in air temperatures associated with the cool dry North-East winds. This is in line with the findings of Indabawa (2009) which reported variations in water temperature in the dry season and can be attributed to intensified heat radiation and effect of harmattan. In the water temperature, no significant difference ( $p > 0.05$ ) was observed between the months, which was similar with the observation of Tisser *et al.* (2008), who reported the lack of significant difference ( $p > 0.05$ ) in monthly variations of water temperature as characteristic of the tropical climate. Temperature influences the oxygen content of water, quantity and quality of autotrophs, while affecting the rate of photosynthesis as well as the quality and quantity of heterotrophs. The water pH in the dam was within 6.53 to 7.76, which makes the water of the dam to be circum-neutral during the study. This was similar to the results of Ibrahim *et al.* (2009) that hydrogen ion concentration (pH) was nearly neutral throughout both seasons, and it was within the range for inland water (pH 6.5 - 8.5) in Kontagora Reservoir, Niger State, Nigeria, which makes it suitable for optimal biological activity. The little increase in pH during the dry season was due to decaying and decomposition of living organisms in the water coupled with the reduction in the water level during the dry season.

The little decrease in pH during the rainy season might be due to the effect of incoming rainwater. According to Janjua *et al.*, (2009) the drop in pH can also be due to the stirring effect of the incoming flood from the rivers and streams that converged towards the lake. It results in the mixing of the poorly alkaline or acidic bottom water with alkaline surface water to reduce pH as evidenced in Shahpur Dam, Pakistan. The transparency of the reservoir was high during the dry season; the higher values of transparency in the dry season may be due to the settling effect of surface run-offs and suspended materials that followed the cessation of rainfall. Ayoade *et al.* (2006) observed the onset of rain decreased the Secchi-disc visibility in two mine lakes around Jos. This supports the observation of Mustapha (2008) that the transparency of water is affected by the amount of the suspended solids in it, and it reduces the light penetrating depth, and hence, reduces the growth of the plants. Dissolved oxygen in the dam indicates two peaks, high in the wet season and low in the dry season. Dissolved oxygen supply in water mainly comes from atmospheric diffusion and photosynthetic activity of plants (Akomeah, *et al.* 2010). The drop of oxygen values from February to April may be due to low temperature in the reservoir. Oxygen plays the most important role in determining the potential biological quality of water.

The highest value was recorded in the dry season while the lowest was recorded in the wet season. The value may be due to chemical fertilizers from irrigated farmlands around the dam coupled with higher rate of evaporation that reduced the level of the water during the dry season. Thus conductivity of water depends upon the concentration of ions and its nutrients status.

Dr S. U. Ibrahim

Table 2: Monthly variation of water physico-chemical parameters of Tsoawa Reservoir

Parameters	February	March	April	May	June	July
D.O (mg/l)	8.60±1.95 <sup>b</sup>	7.60±1.82 <sup>ab</sup>	5.76±2.10 <sup>a</sup>	8.80±3.03 <sup>b</sup>	13.60±3.85 <sup>c</sup>	5.92±1.34 <sup>a</sup>
Conductivity (µS/cm)	91.60±1.67 <sup>c</sup>	90.80±1.48 <sup>bc</sup>	89.20±0.84 <sup>b</sup>	89.80±1.10 <sup>cd</sup>	88.20±0.84 <sup>b</sup>	74.60±1.14 <sup>a</sup>
TDS (mg/l)	0.15±0.04 <sup>a</sup>	0.13±0.03 <sup>a</sup>	0.11±0.03 <sup>a</sup>	0.10±0.01 <sup>a</sup>	0.09±0.03 <sup>a</sup>	0.26±0.38 <sup>a</sup>
Transparency (cm)	135.40±25.25 <sup>ab</sup>	108.40±2.30 <sup>ab</sup>	108.00±1.41 <sup>a</sup>	107.40±0.55 <sup>a</sup>	104.40±1.82 <sup>a</sup>	99.60±2.70 <sup>a</sup>
pH	7.04±0.30 <sup>b</sup>	7.76±0.47 <sup>c</sup>	7.91±0.11 <sup>c</sup>	6.53±0.20 <sup>a</sup>	6.99±0.47 <sup>b</sup>	6.55±0.06 <sup>a</sup>
Nitrate	0.21±0.02 <sup>ab</sup>	0.43±0.17 <sup>d</sup>	0.35±0.09 <sup>cd</sup>	0.32±0.03 <sup>c</sup>	0.29±0.03 <sup>bc</sup>	0.16±0.03 <sup>a</sup>
Phosphate	0.10±0.03 <sup>a</sup>	0.10±0.01 <sup>a</sup>	0.19±0.05 <sup>a</sup>	0.14±0.03 <sup>a</sup>	0.15±0.03 <sup>a</sup>	0.30±0.15 <sup>b</sup>
T. Alkalinity (mg/l)	21.62±0.42 <sup>b</sup>	18.86±0.94 <sup>a</sup>	19.26±0.48 <sup>ab</sup>	20.64±0.21 <sup>ab</sup>	19.90±0.60 <sup>ab</sup>	38.80±4.15 <sup>c</sup>
T. Hardness (mg/l)	41.00±1.95 <sup>c</sup>	31.46±1.64 <sup>a</sup>	43.38±0.85 <sup>d</sup>	36.92±1.21 <sup>b</sup>	37.82±1.21 <sup>b</sup>	37.20±2.70 <sup>b</sup>
Temperature (°C)	27.40±1.53 <sup>ab</sup>	26.36±0.70 <sup>a</sup>	28.60±4.55 <sup>ab</sup>	30.32±0.81 <sup>bc</sup>	32.40±4.17 <sup>c</sup>	26.44±0.55 <sup>a</sup>
Depth (m)	6.38±2.75 <sup>a</sup>	6.46±2.82 <sup>a</sup>	6.38±2.75 <sup>a</sup>	7.92±3.52 <sup>a</sup>	10.44±5.16 <sup>b</sup>	11.18±4.80 <sup>b</sup>

Discussion

The water temperature of the reservoir fluctuated within months, which was between 26.36°C and 32.40°C. The low water temperature recorded in the reservoir was in the dry season, which could be as a result of seasonal changes in air temperatures associated with the cool dry North-East winds. This is in line with the findings of Indabawa (2009) which reported variations in water temperature in the dry season and can be attributed to intensified heat radiation and effect of harmattan. In the water temperature, no significant difference (p>0.05) was observed between the months, which was similar with the observation of Tisser *et al.* (2008), who reported the lack of significant difference (p>0.05) in monthly variations of water temperature as characteristic of the tropical climate. Temperature influences the oxygen content of water, quantity and quality of autotrophs, while affecting the rate of photosynthesis as well as the quality and quantity of heterotrophs. The water pH in the dam was within 6.53 to 7.76, which makes the water of the dam to be ceccum-neutral during the study. This was similar to the results of Ibrahim *et al.* (2009) that hydrogen ion concentration (pH) was nearly neutral throughout both seasons, and it was within the range for inland water (pH 6.5 - 8.5) in Kontagora Reservoir, Niger State, Nigeria, which makes it suitable for optimal biological activity. The little increase in pH during the dry season was due to decaying and decomposition of living organisms in the water coupled with the reduction in the water level during the dry season.

The little decrease in pH during the rainy season might be due to the effect of incoming rainwater. According to Janjua *et al.*, (2009) the drop in pH can also be due to the stirring effect of the incoming flood from the rivers and streams that converged towards the lake. It results in the mixing of the poorly alkaline or acidic bottom water with alkaline surface water to reduce pH as evidenced in Shahpur Dam, Pakistan. The transparency of the reservoir was high during the dry season; the higher values of transparency in the dry season may be due to the settling effect of surface run-offs and suspended materials that followed the cessation of rainfall. Ayoade *et al.* (2006) observed the onset of rain decreased the Secchi-disc visibility in two mine lakes around Jos. This supports the observation of Mustapha (2008) that the transparency of water is affected by the amount of the suspended solids in it, and it reduces the light penetrating depth, and hence, reduces the growth of the plants. Dissolved oxygen in the dam indicates two peaks, high in the wet season and low in the dry season. Dissolved oxygen supply in water mainly comes from atmospheric diffusion and photosynthetic activity of plants (Akomeah, *et al.* 2010). The drop of oxygen values from February to April may be due to low temperature in the reservoir. Oxygen plays the most important role in determining the potential biological quality of water.

The highest value was recorded in the dry season while the lowest was recorded in the wet season. The values may be due to chemical fertilizers from irrigated farmlands around the dam coupled with higher rate of evaporation that reduced the level of the water during the dry season. Thus conductivity of water depends upon the concentration of ions and its nutrients status.

Dr. S. U. Ibrahim

4157/570

Water hardness was higher in the month of April and lower in the month of March and could be as a result of water levels and the concentration of nutrients. The lack of significant difference ( $p > 0.05$ ) between stations and seasons could be because of water low levels and concentration of carbonates. This result is in contrast with the findings of Balogun *et al.* (2005) who observed that water hardness was highly significant ( $p < 0.05$ ) between stations and within the months in Makwaye Lake, Zaria. Nitrate-nitrogen was found to exhibit monthly variation range of 0.43 mgL<sup>-1</sup> to 0.16 mgL<sup>-1</sup>. The mean value recorded was higher in May than in June. The reason for this high concentration in rainy season may be due to excessive influx of nutrients from farmlands where fertilizer is used to boost crop production particularly around the reservoir, as well as input through runoff into the reservoir.

The findings of this study tally with that of Balogun *et al.* (2005) who observed mean monthly variation significant difference ( $p < 0.05$ ) between seasons in Makwaye. The reservoir has higher value of TDS during the dry season; this could be due to decaying of vegetation, higher rate of evaporation caused by increase in air temperature and wind during the dry season. Similar observation was made by Atobatele and Ugwumba (2008) who reported increase in the values of total dissolved solids during the dry season which may be due to most of the vegetation decaying, so giving rise to amount of dissolved solids. However, during the rainy season, the amount of total solids was low, and this may be due to the dilution of water. The values of phosphate-phosphorus in the dam during the dry season may be due to reduced water volume, intensive agricultural activities around the Reservoir as well as the use of fertilizers and pesticides to produce dry season crops like vegetables and maize. Farmers were also using the water from the dam for domestic activities including washing of clothes with detergents, which increased the phosphate-phosphorus level of the water. The result of phosphate-phosphorus variation within seasons also conform with the result of Balogun *et al.* (2005), who observed high significant phosphate-phosphorous variation within months and no significant variation between the sampling stations in Makwaye Lake Zaria. The water depth of the dam fluctuated within seasons; the water depth increased during the rainy season, while it decreased in the dry season. Ibrahim *et al.* (2009), made similar observation of water depth fluctuation within seasons in Kwantagora Reservoir. As the depth of the reservoir increased, dissolved oxygen decreased. Araoye (2008) reported that the depth of the reservoir decreased light intensity, the light penetration depends on the available intensity of the incident light, which varies with geographical location of the reservoir.

#### References

- APHA. 1999. *Standard Methods for Examination of Water and Waste Water*. American Public Health Association, New York, U.S.A.
- Abubakar, I. 2009. Limnological Studies for the Assessment of Sabke lake Katsina State, Unpublished MSc. Dissertation, Ahmadu Bello University, Zaria.
- Araoye, P. A. 2008. Physical factors and their Influence Fish Species Composition in Asa Lake, Ilorin, Nigeria. *International Journal of Tropical Biology*, 57 (1-2): 167-175.
- Atobatele, O. E. and Ugwumba, O. A. 2008. Seasonal Variation in the Physico-chemistry of a Small Tropical Reservoir (Aiba Reservoir, Iwo, Osun, Nigeria). *African Journal of Biotechnology*, Vol. 7 (12): 62-171.
- Ayoade, A. A. 2009. Changes in Physico-chemical Features and Plankton of Two Regulated High Altitude Rivers, Garhwal Himalaya, India. *European Journal of Scientific Research*, 27 (1) 7792. Retrieved from <http://www.eurojournals.com/ejsr.htm> on 25/12/2011.
- Balogun, J. K., Balarabe, M. L. and Igberaese, P. M. 2005. Some Aspects of the Limnology of Makwaye (Ahmadu Bello University Farm) Lake, Samaru, Zaria. *Academic Journal* 23(12): 850-860.
- FAO. 2009. The State of World Fisheries and Aquaculture. FAO Website. [www.fao.org/sof/sofia/index\\_en.htm](http://www.fao.org/sof/sofia/index_en.htm).
- Ibrahim, B. U., Auta, J. and Balogun, J. K. 2009. An Assessment of the Physico-chemical Parameters of Kontagora Reservoir, Niger State, Nigeria. *Bayero Journal of Pure and Applied Sciences*, 2(1): 64-69.
- Indabawa, J.I. 2009. Studies on Limnological Parameters and Phytoplankton Dynamics of Nguru Lake, Yobe State, Nigeria. *Bioscience Research Communications*, 21(4), 183-188.
- Janjua, M.Y. Ahmad, T. and Gerdeau, D. 2008. Comparison of Different Predictive Models for Estimating Fish Yields in Shahpur Dam, Pakistan. *Lake Reservoir Management*, (13): 3324.

Dr S.Y Ibrahim

458/570

Mustapha, M.K. et al.:  
Assessment of some Physico-Chemical Parameters of Tagwai Reservoir  
in Minna, Niger State, Nigeria

Mustapha, M.K. 2008. Assessment of the Water Quality of Oyun Reservoir, Offa, Nigeria, using Selected Physico-chemical Parameters. *Turkish Journal of Fisheries and Aquatic Sciences*, 8, 309-319.

Nathanson, Jerry A. 2003. *Basic Environmental Technology: Water Supply, Waste Management and Pollution Control*. Upper Saddle River, New Jersey: Prentice Hall. Retrieved from <http://www.academicjournals.org> /AJB on 23/12/2011

Tiscer, F. A., Tanimu, Y. and Chia, A. M. 2008. Seasonal Occurrence of Algae and Physico chemical Parameters of Samaru Stream, Zaria, Nigeria. *Asian Journal of Earth Sciences*, 1(1), 31-37.

Ude, E. F., Ugwu, L. L. C., Mgbenka, B. O. and Nwani, C.D. 2011. Trends in Nitrate- Nitrogen, Nitrite-Nitrogen and Phosphorus Concentration in Ebonyi River, Nigeria. *Continental Journal of Fisheries and Aquatic Science*, 5(1): 1-7, 2011. Retrieved from <http://www.wiloludjournal.com>



# ANIFS2019

Proceedings of the 2nd Conference and Annual General Meeting  
Capture Fisheries and Aquaculture Research in the 21st Century Nigeria's Economy  
July 9-11, 2019 held at the Main Auditorium,  
Usmanu Danfodiyo University, Sokoto.

Access online at:  
[www.anifs.org.ng](http://www.anifs.org.ng)

Dr. S. Ibrahim

459/570