



(KEJAANS)

KEBBI JOURNAL OF AGRICULTURE AND NATURAL SCIENCES

May, 2026, Vol. 2, Issue 2



KEJAANS

CONTACT:

The Editor-in-Chief,
Kebbi Journal of Agriculture and Natural Sciences,
Faculty of Agriculture,
Abdullahi Fodjo University of Science and Technology
Aliero,
PMB 1144, Birnin Kebbi, Nigeria.
Email: kejaans.foa@gmail.com.
Phone: +2348039370546
Website: www.kejaansafusta.com

ISSN: 3122-0584



KEBBI JOURNAL OF AGRICULTURE AND NATURAL SCIENCES
(KEJAANS)

May, 2026; Volume 2, Issue 2

OFFICIAL JOURNAL OF THE
FACULTY OF AGRICULTURE
ABDULLAHI FODIO UNIVERSITY OF SCIENCE AND TECHNOLOGY,
ALIERO

Editors

I.S. Jega
M.I. Ribah
I. Sani
M. Atiku
N.M. Kwaifa

KEJAANS



About the Journal

This official scientific publication of the Faculty of Agriculture, Abdullahi Fodio University of Science and Technology Aliero, is a non-profit, open access, double-blind peer-reviewed Journal publishing four issues (January, April, July and October) per annum. The Journal is a platform open to collaborations with researchers, authors, institutions, research agencies and private companies related to Agriculture. The Mission of the Journal is to disseminate scientific knowledge through the publication of original research articles, research notes, book reviews, letters to the editor and reviews of Literature, representing a contribution to scientific and technological knowledge in respective areas covered by the Journal. The Kebbi Journal of Agriculture and Natural Sciences seeks to validate and disseminate new knowledge, making it public in order to strengthen the human capacity, constitute a link in the scientific community to the society and encouraging the expansion of University and academic researches.

Scope of Kebbi Journal of Agriculture and Natural Sciences (KEJAANS)

The Kebbi Journal of Agriculture and Natural Sciences has the sole aim of providing an intellectual platform and ideas for scholars, by promoting interdisciplinary studies related to agriculture and natural science through publishing the latest scientific research findings that are of direct policy implications and beneficial to the research community. Consequently, the journal covers all aspects of Crop Science, Animal Science, Agricultural Economics, Agricultural Extension and Rural Development, Food Science, Fisheries and Aquaculture, Biotechnology, Soil Science and Agricultural Engineering, Forestry and Environment, Wildlife, Agricultural Education, Agro-allied Industries as well as all Natural Science researches related to Agriculture.

KEJAANS

TABLE OF CONTENT

S/N	Article and Author(s)	Page
01	Profitability And Constraints Of Broiler Chicken Production: Insights From The Agricultural Teaching And Research Farm, Federal College Of Education (Technical), Akoka, Nigeria Olowa O.W, Umoru, J.I.A and Olowa, O.A	1-10
02	<i>Carica Papaya</i> Leaf-Mediated Nanoparticle Synthesis: A Practical Approach In The Control Of Field Insects Of Soybeans. Akanji, Kayode Ayanwusi., Alao, Fatai Olaitan., Olaniran, Oladele Abiodun., Akanbi, Wasiu Babatunde and Adebayo, Timothy Abiodun.	11-23
03	Assessment Of Phytoplankton Composition And Abundance Of The Upper Niger River Basin Reservoir Ibrahim, S. U., Kolo, R. J., Peter, J. S. and Gana, A. B.	24-31
04	The Role Of Precision Agriculture On Food Security In Borno State: Barriers, Adoption, And Outcomes For Smallholder Farmers Modibbo, Hussaina1, And Chindo, I. Y1, And Muritala, I. K	32-42
05	Assessment Of Processing Characteristics And Sensory Quality Of Smoked Nile Tilapia Treated With Field Mint (<i>Mentha Arvensis</i>) Solution Kurfi, A.R., Rangaza F.A	43-49
06	Evaluation Of Suitable Land For Upland Rice (<i>Oryza Sativa</i>) Production At Toposequence Of Ncri, Badeggi, Niger State, Nigeria Daniel. M., Ibrahim. P. A., Shema. A. M., Lawal. B.A., Mohammed. T., Apuyor A. Aliyu, E.J., Dalhat, S. M., Okachi M.O Kabaraini, M. A., Shaba. Y.A. Fadieye. E.O, Usman. B., Olagoke. O.	49-59
07	Economics Of Post-Harvest Management Of Selected Vegetables In Jos East Local Government Area, Plateau State Adubok, Abraham Itsegok And Jabil, Irimiya Yarima	60-68
08	Effect Of Scent Leaf (<i>Ocimum Gratissimum</i>) Curing On <i>Clarias Gariepinus</i> Smoked With Soybean (<i>Glycine Max</i>) Shell Briquettes As A Substitute For Firewood *Ibrahim, S. U., Ibrahim, A., Yusuf, J., Umar, F., Gana, A. B., and Ayeboan, B. R.	69-79

09	Impact Of Population Growth On Food Security In Nigeria: Time Series Analysis And Vector Error Correction Approach Ibrahim Sani Shamaki, Emmanuel Ada Ojoko and Adamu Hamisu Danmusa	80-92
10	Effects Of Harvest Interval On Growth, Leaf Area Index, And Biomass Allocation Of <i>Moringa Oleifera</i> (Periyakulam-1) Under Semi-Arid Conditions Samaila Abdullahi and Zubairu Yakubu Gada	93-100
11	Effects Of Species Identity And Seed Weight On Germination And Early Seedling Growth Of <i>Acacia Senegal</i> (L.) Willd. And <i>Azadirachta Indica</i> A. Juss. In The Sudan Savanna Zone Of Nigeria Samaila Abdullahi and Abdurashid Inuwa	101-110



ASSESSMENT OF PHYTOPLANKTON COMPOSITION AND ABUNDANCE OF THE UPPER NIGER RIVER BASIN RESERVOIR

*Ibrahim, S. U., Kolo, R. J., Peter, J. S. and Gana, A. B.

Department of Water Resources, Aquaculture and Fisheries Technology, Federal University of Technology, P. M. B. 65, Minna, Niger State Nigeria

Contacts: suibrahim@futminna.edu.ng +234 7039025846

ABSTRACT

The study evaluated the composition, abundance, and spatial distribution of phytoplankton in the Upper Niger River Basin Reservoir, Nigeria, to assess ecological status and anthropogenic impacts. Samples were collected monthly from February to July 2022 at four sites (landing site, vegetated area, open water and dam site) representing different environmental conditions. Phytoplankton were sampled using a 75 mm plankton net, preserved in 4% formalin and analyzed microscopically for identification and enumeration. Four major phytoplankton classes were identified: Bacillariophyceae, Euglenophyceae, Dinophyceae, and Chlorophyceae. Bacillariophyceae (diatoms) showed the highest diversity across all stations, with the genus *Nitzschia* dominating at the landing site, indicating eutrophic to hypereutrophic conditions linked to organic pollution. Data analysis revealed significant spatial variation ($p < 0.05$) among phytoplankton groups. Euglenophyceae recorded the highest mean abundance at the landing site (151.50 ± 2.12), further suggesting nutrient enrichment. The dominance of pollution tolerant and eutrophic indicator species, especially at the landing site, reflects substantial nutrient input from human activities. This indicates advanced eutrophication, which can threaten ecological integrity through oxygen depletion, biodiversity loss, and declining water quality. These findings provide baseline data essential for developing management strategies to control nutrient pollution and sustain the reservoir's ecological health. Therefore, based on these findings, it is recommended that a continuous phytoplankton water quality program be established and also implement watershed nutrient control actions to reduced nitrogen and phosphorus loading.

Keywords: Phytoplankton, Eutrophication; Upper Niger River Basin Reservoir, Nutrient pollution; Anthropogenic activities.

Introduction

Assessing ecological quality is fundamental to the management and conservation of aquatic ecosystems (Tampo *et al.*, 2023). While chemical analyses provide temporary data, they often fail to capture long-term environmental stressors over time (Schuijt *et al.*, 2021). Consequently, biological assessment, particularly plankton, offers a more integrated approach since these

organisms reflect cumulative environmental conditions. Among them, phytoplankton are especially valuable indicators due to their foundational position in the food web, Sensitivity to physicochemical changes and direct exposure to the water column (Celia *et al.*, 2021) Phytoplankton are also highly suitable for determining the impact of toxic substances on the environment (Gojkovic *et al.*, 2023).

Phytoplankton require light and nutrients for growth, through photosynthesis, they convert solar energy into chemical energy, driving nutrient cycling and supporting higher trophic levels, including zooplankton and fish. Their abundance and composition therefore determine the productivity and carrying capacity of aquatic systems. Thus, the abundance and composition of the phytoplankton community are fundamental determinants of an aquatic system's overall productivity and carrying capacity (Celia *et al.*, 2021).

However, environmental conditions such as nutrient enrichment and rising temperatures can trigger excessive phytoplankton growth, leading to blooms. These blooms increase water turbidity, reduce light penetration, and inhibit submerged vegetation. Upon decomposition, phytoplankton consume dissolved oxygen, causing hypoxic conditions that threaten aquatic life, including fish and benthic organisms. Nevertheless, the same factors that promote their growth can also lead to severe ecological imbalance (Brenckman *et al.*, 2025).

The Upper Niger River Basin Reservoir serves as a vital resource for local communities, supporting their livelihoods through various activities, including domestic use and agriculture (Liersch *et al.*, 2019). However, these anthropogenic activities have led to increased pollution in the reservoir (Gavrilaş *et al.*, 2025). This pollution can significantly impact the abundance and diversity of phytoplankton, which play a crucial ecological role as primary producers, fuelling food webs both directly and indirectly (Naselli-Flores and Padisák, 2023).

Advances in satellite technology has significantly transformed the mapping and monitoring of phytoplankton in aquatic environments. Traditionally, phytoplankton studies relied on labour-intensive field

sampling, which was often limited in spatial and temporal coverage. However, allow scientists to observe large water bodies continuously, providing a synoptic view of phytoplankton distribution across aquatic environment, helping researchers understand bloom dynamics and ecological processes, which is crucial for monitoring water quality. Overall, phytoplankton serve as indispensable tools for evaluating aquatic health and guiding sustainable water resource management.

Materials and Methods

Description of the study area

The research was performed at the Upper Niger River Basin Reservoir, a significant water body in North Central Nigeria. The reservoir is located within the administrative boundaries of the Bosso Local Government Area of Niger State. Geographically, the sampling area lies at coordinates 9°40'40.4" N and 6°31'14.5" E. Field sampling for the determination of phytoplankton composition and abundance was carried out over a six-month duration, encompassing both dry and early rainy season conditions, from February 2022 to July 2022.

Experimental Materials

Field Sampling and Site Characterization

Phytoplankton samples were collected from the reservoir using a plankton net (75 mm) and transferred into pre-labelled plastic jar. The precise geographic coordinates of each sampling station were recorded using a handheld Global Positioning System (GPS) unit (Model: GPSMAP 76 garmin).

Sample collection, Preservation and Laboratory Processing

The samples were collected through a sweep made by moving the net within the water over 0.5 m and turns back and repeated same

process back through the same distance vertically. Immediately after collection, samples were preserved in the field by adding 4% buffered formalin to prevent biotic degradation. In the laboratory, the preserved samples were concentrated for analysis. A fixed volume of each sample was centrifuged using a benchtop centrifuge (Model: 800).

Phytoplankton Identification and Enumeration

Following concentration, a subsample of a known volume was transferred using a calibrated pipette onto a counting chamber. Phytoplankton were identified to the lowest feasible taxonomic level and counted using a high-power compound microscope (Olympus: Model), equipped with a digital camera for

imaging. Taxonomic identification was confirmed using standard authoritative monographs and regional algal flora guides.

Geospatial Analysis

A satellite image of the Upper Niger River Basin Reservoir and its catchment was acquired. The imagery was used for mapping sampling locations, assessing land use and analysing surface water characteristics.

Data Collection

The data was obtained by field survey, satellite images, GIS software, Microsoft word and Microsoft excel (Figure 1), and four sites (Figure 2) were selected based on the activities conducted on the reservoir (Table 1).




Figure 1: Nigeria Map indicating Niger State and the Study Area



Figure 2a: Quick bird Image of the Reservoir Indicating Sampling Stations




Figure 2b: Database map of study area

Table 1: Attribute of the Selected Sampling Site

Sampling	Location	Description
Station 1	Landing site	First point of contact
Station 2	Site close to vegetation	This is the site close to vegetation cover
Station 3	Open water	Site close to the middle of water
Station 4	Dam site	The site close to dam

Data Analysis

Data were analysed using one-way analysis of variance (ANOVA) using Minitab, version 14.0. Differences between treatments were compared using Tukey's multiple comparison test. The level of significance was set at $p < 0.05$.

Results and Discussions

The phytoplankton species identified across the stations in Upper Niger Reservoir (Table 2) offers a qualitative snapshot of the phytoplankton community structure at four distinct stations. Phytoplankton, as primary producers, form the base of most aquatic food webs and are highly sensitive to changes in their environment (Jahan and Singh, 2023).

Four different families of phytoplankton were identified in this study. Similarly, Etim (2019) identified 5 taxonomic families (Bacillariophyceae, Chlorophyceae, Cyanophyceae, Euglenophyceae and Chrysophyceae) in the lower Cross River system, Nigeria. However, this does not corroborate with the results reported by Moruff *et al.* (2016). These variations could be attributed to different location and environment. Consequently, the composition, abundance, and distribution of phytoplankton serve as excellent biological indicators for assessing the ecological status and water quality of aquatic ecosystems like reservoirs (Essa *et al.*, 2024).

Table 2: Phytoplankton Species Identified Across the Stations in Upper Niger Reservoir

	STATION 1	STATION 2	STATION 3	STATION 4
Bacillariophyceae				
<i>Melosira spp</i>	++	—	+	+
<i>Nitzschia spp</i>	+++	++	++	—
<i>Navicula spp</i>	+	—	—	+
<i>Peridinium spp</i>	—	+	++	—
<i>Thalassiora spp</i>	—	++	—	++
<i>Odontella spp</i>	—	+	++	—
<i>Leptocyclindrus spp</i>	+	+	—	+
<i>Rhizosolentia spp</i>	++	—	—	+
Euglenophyceaea				
<i>Prorocentrum spp</i>	++	—	—	++
Dinophyceae				
<i>Gymnodium spp</i>	—	—	++	+
Chlorophyceae				
<i>Pediastrum spp</i>	—	+	—	—
<i>Spirogyra spp</i>	++	—	+	—

— = absent, + = sparse, ++ = abundant and +++ = dominant.

The dominance of Bacillariophyceae (Diatoms) in this study, the most striking feature of the dataset is the pronounced diversity and its abundance. Flori *et al.* (2025) reported that diatoms are ubiquitous in freshwater environments, and their dominance often points to specific environmental conditions, which they require silica for the formation of their frustules. Therefore, their abundance suggests that dissolved silica is not a limiting nutrient in this system. Also, the presence of both centric (*Melosira*, *Thalassiosira*) and pennate (*Nitzschia*, *Navicula*) diatoms indicates a varied habitat structure within the reservoir. This genus (*Nitzschia*) is dominant (+++) at Station 1 and abundant (++) at Stations 2 and 3. However, many species within the genus *Nitzschia* are known to be tolerant of high levels of organic

pollution and nutrient enrichment, particularly nitrogen (Trobajo *et al.*, 2004). Their

dominance is a classic indicator of eutrophic to hypereutrophic conditions (Trobajo *et al.*, 2009). The prevalence of *Nitzschia* across the first three stations, especially its dominance at Station 1, strongly suggests significant nutrient loading in these parts of the reservoir. These motile dinoflagellates (*Peridinium* and *Gymnodinium*) are abundant at Station 3. Hall and Paerl (2011) observed that dinoflagellates can perform vertical migrations in the water column, allowing them to access nutrients in deeper waters and light at the surface, giving them a competitive advantage in stratified or stable water columns. Therefore, their abundance at Station 3 suggests nutrient rich conditions, possibly with more water column stability compared to Station 1.

This study further revealed (Table 3 and Figure1) the quantitative data on the distribution of four major phytoplankton classes (Bacillariophyceae, Euglenophyceae, Dinophyceae, and Chlorophyceae), across four

sampling stations in the Upper Niger Reservoir. This highlights significant spatial variations in the phytoplankton community structure, which are powerful indicators of

differing environmental conditions, particularly water quality and hydrology (Zhang *et al.*, 2021).

Table 3: Station Variation of Phytoplankton Distribution

SPECIES	STATION 1	STATION 2	STATION 3	STATION 4
<i>Bacillariophyceae</i>	71.25±63.56 ^a	51.25±43.48 ^a	47.50±51.47 ^a	47.75±41.06 ^a
<i>Euglenophyceaea</i>	151.50±2.12 ^a	0.00±0.00 ^c	0.00±0.00 ^c	121.50±2.12 ^b
<i>Dinophyceae</i>	0.00±0.00 ^c	0.00±0.00 ^c	92.50±3.53 ^a	81.00±1.41 ^b
<i>Chlorophyceae</i>	40.00±56.56 ^a	55.00±77.78 ^a	30.00±42.42 ^a	0.00±0.00 ^b
<i>Chlorophyceae</i>	40.00±56.56 ^a	55.00±77.78 ^a	30.00±42.42 ^a	0.00±0.00 ^b

Values on the same row with different superscripts are significantly different ($p < 0.05$)



Figure 3a: Shonghai reservoir indicating the distribution of *Bacillariophyceae*



Figure 3b: Shonghai reservoir indicating the distribution of *Euglenophyceae*



Figure 3c: Shonghai reservoir indicating the distribution of *Dinophyceae*



Figure 3d: Shonghai reservoir indicating the distribution of *Chlorophyceae*

Figure 3: Shonghai Reservoir Showing the Distribution of Phytoplankton

In station 1, *Euglenophyceaea* recorded the highest mean values and differ significantly ($p < 0.05$) from other stations. This high variability suggests that the distribution of these phytoplankton is highly heterogeneous within the stations. Such heterogeneous is a fundamental characteristic of plankton ecology and can be caused by micro-scale nutrient hotspots, physical aggregation, and boom-and-bust dynamics (Tang *et al.*, 2024). The mean abundance of *Bacillariophyceae* ranges from 47.50 to 71.25 across the stations. Statistically, there is no significant difference ($p < 0.05$) in their abundance among the four stations. This result suggests that *Bacillariophyceae* form a ubiquitous and relatively stable component of the phytoplankton community throughout the sampled areas of the reservoir. The factors controlling their overall abundance, such as the availability of dissolved silica and general light conditions, appear to be relatively uniform across these stations.

References

- Brenckman, C. M., Parameswarappa, J., William, H., Ashraf, F., & Borgaonkar, A. D. (2025). A review of harmful algal blooms: Causes, effects, monitoring, and prevention methods. *Water*, 17(13), 1980.
- Celia, G. V. S., Norberto, M. A. P. M., Enriquez-Ocaña, L. F., Ramon, H., Holguin, J. E. V., & Martinez-Cordova, L. R. (2021). Phytoplankton composition and abundance as indicators of aquaculture effluents impact in coastal environments of mid Gulf of California. *Heliyon*, 7(2).
- Essa, D. I., Elshobary, M. E., Attiah, A. M., Salem, Z. E., Keshta, A. E., & Edokpayi, J. N. (2024). Assessing phytoplankton populations and their relation to water parameters as early alerts and biological indicators of the aquatic pollution. *Ecological Indicators*, 159, 111721.
- Etim, J. B. (2019). Assessment of phytoplankton species composition, abundance and diversity within the it bridge-end area of the lower Rross River system, Nigeria. *International Journal of Fisheries and Aquatic Studies*, 7(4), 398-405.
- Flori, S., Mikus, F., Flaum, E., Moog, K., Guessoum, S., Beavis, T., Zwahlen, S. M., & Olivetta, M. (2025). Diatom ultrastructural diversity across controlled and natural environments. *Current Biology*, 35, 1 - 12.
- Gavrilaş, S., Burescu, F., Chereji, B., & Munteanu, F. (2025). The impact of anthropogenic activities on the

Conclusion

The qualitative analysis of the phytoplankton community in the Upper Niger Reservoir indicates a significant gradient of water quality across the four stations. The assemblage at Station 1 strongly suggests hypereutrophic conditions, likely driven by high inputs of organic and inorganic nutrients. Stations 2 and 3 also show clear signs of eutrophication, though the community composition differs, pointing to spatial variations in environmental drivers. Station 4 appears to be comparatively less impacted. The dominance of eutrophic indicator species is a cause for concern, as advanced eutrophication can lead to oxygen depletion, loss of biodiversity, fish deaths, and a general degradation of water quality, impacting its use for drinking, fishing, and recreation. Hence, this information is crucial for targeted management strategies aimed at mitigating nutrient pollution and preserving the ecological health of the reservoir.

- catchment's water quality parameters. *Water*, 17(12), 1791.
- Gojkovic, Z., Simansky, S., Sanabria, A., Márová, I., Garbayo, I., & Vílchez, C. (2023). Interaction of naturally occurring phytoplankton with the biogeochemical cycling of mercury in aquatic environments and its effects on global Hg pollution and public health. *Microorganisms*, 11(8), 2034.
- Hall, N. S., & Paerl, H. W. (2011). Vertical migration patterns of phytoflagellates in relation to light and nutrient availability in a shallow microtidal estuary. *Marine Ecology Progress Series*, 425, 1-19.
- Jahan, S., & Singh, A. (2023). The role of phytoplanktons in the environment and in human life: A review. *Basrah Journal of Sciences*, 41(2), 392-411.
- Liersch, S., Fournet, S., Koch, H., Djibo, A. G., Reinhardt, J., Kortlandt, J., & Baker, C. (2019). Water resources planning in the Upper Niger River basin: Are there gaps between water demand and supply? *Journal of Hydrology: Regional Studies*, 21, 176-194.
- Moruff, A. K., Mu'azu, M. M., Patric, A. R., & Lawrence, O. O. (2016). Diversity of phytoplankton communities in a Tropical River Basin, Nigeria. *Journal of Water, Energy and Environment*, 19, 52-56.
- Naselli-Flores, L., & Padisák, J. (2023). Ecosystem services provided by marine and freshwater phytoplankton. *Hydrobiologia*, 850(12), 2691-2706.
- Schuijt, L. M., Peng, F., Sanne, J. P., Dingemans, M. M. L., & Paul, J. (2021). (Eco) toxicological tests for assessing impacts of chemical stress to aquatic ecosystems: facts, challenges, and future. *Science of the Total Environment*, 795, 148776.
- Tampo, L., Kaboré, I., Mande, S. A. L. S., & Bawa, L. M. (2023). Water quality and Aquatic ecosystem assessment using water quality Indices in West Africa: Challenge and perspectives. In *Water quality-new perspectives*. IntechOpen.
- Tang, X., Liu, L., Wang, Y., Xu, J., Huang, Y., & Zhang, F. (2024). Characteristics of plankton community structure and applicability of biomanipulation in midstream of the Jialing River the biggest tributary of the Yangtze River. *Journal of Freshwater Ecology*, 39(1), 2378835.
- Trobajo, R., Clavero, E., Chepurnov, V. A., Sabbe, K., Mann, D. G., Ishihara, S., & Cox, E. J. (2009). Morphological, genetic and mating diversity within the widespread bioindicator *Nitzschia palea* (Bacillariophyceae). *Phycologia*, 48(6), 443-459.
- Trobajo, R., Cox, E. J., & Quintana, X. D. (2004). The effects of some environmental variables on the morphology of *Nitzschia frustulum* (Bacillariophyta), in relation its use as a bioindicator. *Nova Hedwigia*, 79(3), 433-445.
- Zhang, Y., Gao, W., Li, Y., Jiang, Y., Chen, X., Yao, Y., & Chen, Y. (2021). Characteristics of the phytoplankton community structure and water quality evaluation in autumn in the Huaihe River (China). *International Journal of Environmental Research and Public Health*, 18(22), 12092.