



# UMYU SCIENTIFICA

(Formerly Katsina Journal of Natural and Applied Sciences, KAJONAS)

*An official publication of the Faculty of  
Natural and Applied Sciences  
Umaru Musa Yar'adua University,  
P.M.B 2218 Katsina, Nigeria*

**Volume 3, No. 4  
December, 2024**

ISSN 2955-1145 (print)  
ISSN 2955-1153 (online)

Adamu, K. M., Mohammed, Y. M., Muhammad, H., Achebe, A. C. & Jeremiah, J. (2021a). Morphometric and biomarker indices of fishes in Dangana Lake, Lapai, Niger State. *Direct Research Journal of Biology and Biotechnology*, 7, 16-27. <https://doi.org/10.26765/4772408>.

Adamu, K.M., Kuyembo, S.M., Abbas, A.H., Mohammed, Y.M., Djadjiti, N. & Mangse, G. (2021b). Growth performance of African Catfish, Spinach and Lettuce in an aquaponic system. *Dutse Journal of Agriculture and Food Security*, 8(2), 96-106.

Adelakun, K.M., Mustapha, M.K., Amali, R.P. & Mohammed, N. (2017). Seasonal variation in nutritional quality of Catfish (*Clarias gariepinus*) from Upper Jebba Basin, Nigeria. *Journal of Nutrition & Food Sciences*, 7(5), 1-4.

Agbugui, M. O., Abhulimen, E. F., Inobeme, A. & Olori, E. (2019). Biodiversity of Fish Fauna in River Niger at Agenebode, Edo State, Nigeria *Egyptian Journal of Aquatic Biology & Fisheries*, 23(4), 159-166. <https://doi.org/10.21608/ejabf.2019.52847> DOI: <https://doi.org/10.21608/ejabf.2019.52847>

Aliko, N.G.G., Da costa, K.S., Dietoa, Y.M., Ouattara, A. & Gourene, G. (2010). Caractéristiques de la population de *Distichodus rostratus* Günther, 1864 (Pisces: Distichodontidae) du lac de barrage de Taabo (bassin du Bandama, Côte d'Ivoire). *Implications pour une gestion rationnelle du stock. Tropicultura*, 28, 50-56.

Atile, J.I., Shima, J.N. & Akombo, P.M., (2016). Food and feeding, length-weight and condition factor of the catfish *Synodontis membranaceus* (Etiene Geoffroy Saint Hilaire, 1809) (Osteichthyes: Mochokidae) from Lower Benue River, Makurdi, Nigeria. *Agriculture, Forestry and Fisheries*, 5(4), 87-96. <https://doi.org/10.11648/j.aff.20160504.11> DOI: <https://doi.org/10.11648/j.aff.20160504.11>

Benedict, O., Offem, Yemi, A. S. & Omoniyi, I. T., (2009). Length-weight relationship condition factor and sex ratio of forty six important fishes in a Tropical Flood River. *Research Journal of Fisheries and Hydrobiology*, 4(2), 65-72, 2009.

Buwono, N.R., Fariedah, F. & Anestyningrum, R.E. (2017). Komunitas Ikan di Sungai Jerowan Kabupaten Madiun. *Journal of Aquaculture and Fish Health*, 6(2), 81-88. <https://doi.org/10.20473/jafh.v6i2.11284> DOI: <https://doi.org/10.20473/jafh.v6i2.11284>

Dienye H.E, Olopade, O.A. & Toby, S.A., (2018) Species Composition and Diversity of Cast net fisheries in new Calabar, Niger Delta, Nigeria. *Journal of Biodiversity, Conservation & Bioresources Management*, 4(1), 19-26. <https://doi.org/10.3329/jbcbm.v4i1.37873> DOI: <https://doi.org/10.3329/jbcbm.v4i1.37873>

Essien-Ibok, M. A. (2020). Fish fauna of Akwa Ibom State inland waters. *Biodiversity International Journal*, 4(2), 82-88.

Ezekiel, E. N. & Abowei, J. F. N., (2014). A study of length-weight relationship and condition factor of *Hepsetus odoe* (Bloch, 1794) from Amassoma flood plains. *Annals of Biological Sciences*, 2(2), 10-17.

Froese, R. & Pauly, (2018). Diversity of Fish fauna. Retrieved 4/6/2018.

Ibrahim, M. I., Ibrahim, Y., Abdullahi, A. M., Najibullah, B. A, Obi, P. U. & Mohammed, Y. M. (2022).

Bacteria associated with smoked catfish sold at Bida Modern market, Northcentral Nigeria.

*International Journal of Fisheries and Aquatic Studies*, 10(2), 38-40.

<https://doi.org/10.22271/fish.2022.v10.i2a.2654> DOI: <https://doi.org/10.22271/fish.2022.v10.i2a.2654>

Kumar, U. & Asija, M. J. (2009). Biodiversity: Principle and Conservation. Agrobios (India) Leonard, T., Gouli, G. B. and Kouassi, S. Da Costa (2011). Length-weight relationships for 36 freshwater fish species from two tropical reservoirs: Ayamé I and Buyo, Côte d'Ivoire. *Rev. Biol. Trop. (Int. J. Trop. Biol., 60 (4), 1847-1856.* <https://doi.org/10.15517/rbt.v60i4.2185> DOI: <https://doi.org/10.15517/rbt.v60i4.2185>

Lawal, N., Abdullahi, S.A. & Abolude, D.S. (2023). Physicochemical Characteristics and Fish Abundance and Diversity of Mairua Reservoir Water, Funtua, Katsina State, North-Western Nigeria. *Journal of Applied Science and Environmental Management* 27(1), 125-132.

<https://doi.org/10.4314/jasem.v27i1.18> DOI: <https://doi.org/10.4314/jasem.v27i1.18>

Leonard, T., Gouli, G.B. & Kouassi, S.D. (2011). Length-weight relationships for 36 freshwater fish species from two tropical reservoirs: Ayamé I and Buyo, Côte d'Ivoire. *International Journal of Tropical Biology*, 60(4), 1847-1856. <https://doi.org/10.15517/rbt.v60i4.2185> DOI: <https://doi.org/10.15517/rbt.v60i4.2185>

Mo, Y., Wan, H., Cai, Y., Zhang, X., Li, R. & Wang, Y. (2023). Re-establishing fish migration channel of large reservoirs in Jinsha River Basin of China by using an eco-friendly reservoir operation method, *Journal of Hydrology: Regional Studies*, 47, 101412, <https://doi.org/10.1016/j.ejrh.2023.101412> DOI: <https://doi.org/10.1016/j.ejrh.2023.101412>

Mohammed, Y. M., Abubakar, M. D., Muhammad, A. M., Muhammad, A. S., Umar, B. L., Hadizat, M. & Achebe, A. C. (2021a). Stomach Content of *Tillapia zilli* and *Oreochromis niloticus* from Wanzun River, Lavun Local Government, Niger State Nigeria. *Asian Journal of Fisheries and Aquatic Research*, 13(5), 15-20. Article no.AJFAR.70239. <https://doi.org/10.9734/ajfar/2021/v13i530276> DOI: <https://doi.org/10.9734/ajfar/2021/v13i530276>

Mohammed, Y.M., Adamu, K.M., Ibrahim, B.U., Danjuma, S. & Amuzat, A.O. (2023). Exploring the Impact of Human Activities on Water Quality Characteristics of a Rural River in North-Central Nigeria using Macroinvertebrates Structural Assemblage. *West African Journal of Life Sciences*, 1, 023-034. <https://doi.org/10.4314/dujopas.v10i2c.1> DOI: <https://doi.org/10.4314/dujopas.v10i2c.1>

Mohammed, Y.M., Arimoro, F. O., Ayanwale, A.V., Adama, B. S., Keke, U.N., Auta, Y. I. & Umar, B.L. (2020). Seasonal changes in the abundance of benthic macroinvertebrates & physico-chemical condition of Moussa stream Bida, Nigeria. *Tropical Freshwater Biology*, 29(1), 57-70.

<https://doi.org/10.4314/tfb.v29i1.4> DOI: <https://doi.org/10.4314/tfb.v29i1.4>

Mohammed, Y.M., Arimoro, F.O., Ayanwale, A.V. Adamu, K.M., Keke, U.N., Abubakar, M.D. & Achebe, A.C. (2021b). The current state of water quality and benthic invertebrate fauna in Chikke Stream (North-Central Nigeria). *Ukrainian Journal of Ecology*, 11(3), 26-34. [https://doi.org/10.15421/2021\\_136](https://doi.org/10.15421/2021_136)

Mustapha, M.K. (2008). Assessment of the Water Quality of Oyun Reservoir, Offa, Nigeria, Using Selected Physico-Chemical Parameters. *Turkish J Fish Aqua Sci.* 8, 309-319.

Mustapha, M.K. (2010). Fish fauna of Oyun reservoir, Offa, Nigeria. *Journal of Aquatic Sciences*, 25 (1), 106-114.

Nazeef, S., Ja'afar A., Abubakar K.A., and Kabiru M., (2021). Fish species biodiversity of Dadin-Kowa reservoir. *World Journal of Advanced Research and Reviews*, 11(02), 100-111.

<https://doi.org/10.30574/wjarr.2021.11.2.0370> DOI: <https://doi.org/10.30574/wjarr.2021.11.2.0370>

Okomoda, V.T., Solomon, S.G. & Aladi, S.L. (2012). Fish Fauna in Lower River Niger at Idah in Kogi State. *Journal of Agricultural & Veterinary science*, 4, 34-40.

Oladipo, S.O., Nneji, L.M., Iyiola, O.A., Nnejid, I.C., Ayoola, A.O., Adalakune, K.M., Anifowoshe, A.T. Adeola, A.C. & Mustapha, M.K. (2021). Patterns of ichthyofaunal diversity and distribution across Jebba Hydro-Electric Power (HEP) dam, Jebba, north-central Nigeria. *Brazilian Journal of Biology*, 81(2), 258-2. <https://doi.org/10.22271/fish.2022.v10.i2a.2654> DOI: <https://doi.org/10.1590/1519-6984.222952>

Olopade O.A., Dienye H.E., Bamidele N. & Oderhohwo, O. (2020). Catchability and diversity of fish species captured by gill net in new Calabar River, Nigeria. *Indonesia Fishery Research Journal*, 26(2), 107-117. <https://doi.org/10.15578/ifrj.26.2.2020.107-117> DOI: <https://doi.org/10.15578/ifrj.26.2.2020.107-117>

Olowo, U.C., Egun, N. K., Ehiglatior, A.P. & Oboh, I.P. (2022). Growth analysis and sex ratio of fish species from the Ovia River, Edo State, Nigeria. *BIOLOGIJA*. 2022. Vol. 68. No. 2. P. 110–120. <https://doi.org/10.6001/biologija.v68i2.4731> DOI: <https://doi.org/10.6001/biologija.v68i2.4731>

Omoike, A. (2021) The Trend in Fish Species Diversity and Abundance at Asejire Reservoir South Western Nigeria. *HSOA Journal of Aquaculture & Fishery*, 5(040), 1-9. <https://doi.org/10.24966/AAF-5523/100040> DOI: <https://doi.org/10.24966/AAF-5523/100040>

Pius, D.E., Ja'afaru, A., Abubakar K.A., Zira J.D., Kefas, M. & Barau B.W. (2020). Fish biodiversity and abundance in river Taraba, Taraba State, Nigeria. *Ethiopian Journal of Environment Studies & Management*, 13(5), 535-544.

Rosa, R.S., Charvet-Almeida, P. & Quijada, C.C.D. (2010). Biology of the South American Potamotrygonid stingrays, p. 241-285. <https://doi.org/10.1201/9781420080483-c5> DOI: <https://doi.org/10.1201/9781420080483-c5>

Rusmilyansari, A. A., Wahab, W. & Reika, C. (2021). Fish species composition and diversity in a river, a swamp, and a reservoir in Banjar District, South Kalimantan Province. *AAAL Bioflux*, 14(1), 412-423.

Seehausen, O. (2015). Process and pattern in cichlid radiations – inferences for understanding unusually high rates of evolutionary diversification. *New Phytology*, 207: 304-312. <https://doi.org/10.1111/nph.13450> DOI: <https://doi.org/10.1111/nph.13450>

Sikoki, F.D., Zabbey, N. & Anyanwu, N. (2008). Fish Assemblage of Onu -Iyi-Ukwu Stream in Southern Tropical Nigeria. *Fish Biology*, 17(2), 13-23. <https://doi.org/10.4314/tfb.v17i2.45813> DOI: <https://doi.org/10.4314/tfb.v17i2.45813>

Solomon, S.G, Ayuba V.O, Tahir, M.A. & Okomoda, V.T. (2017). Abundance composition of fish in lake kalgwai Jigawa State, Nigeria. *Jordan Journal of Agricultural Sciences*, 13, 45-54. <https://doi.org/10.12816/0039652> DOI: <https://doi.org/10.12816/0039652>

Solomon, S.G. Okomoda, V.T. & Aladi. S.L. (2012). Fish Fauna in Lower River Niger at Idah in Kogi State. *Journal of Agricultural and Veterinary Science*, 4, 34-40.

Sulaiman, Z., Hu T. H. & Lim K.K.P. (2018). Annotated checklist of freshwater fishes from Brunei Darussalam, Borneo. *Zootaxa*, 4379(1), 24-46. <https://doi.org/10.11646/zootaxa.4379.1.2> DOI: <https://doi.org/10.11646/zootaxa.4379.1.2>

Uduma, A.U. (2018). Physico-chemical analysis of the quality of sachet water consumed in Kano metropolis. *American Journal of Environment, Energy and Power Research* 2, 1-10.

Wu, Z.Q. & Huang, L.L. (2017). *Fish Biodiversity and its Relationship with Environment in Lijiang River*. Science Press, Beijing.

Wubie, A., Tadesse, Z., Tesfay, G. & Lakew, A. (2017). Fish distribution and abundance in the Upper Awash River, West Shewa. *Ethiopian Journal of Science and Sustainable Development*, 5(2), 109-118.

Xing, Y. C., Zhang, C. G., Fan, E. Y. & Zhao, Y. H. (2016). Freshwater fishes of China: species richness, endemism, threatened species and conservation. *Diversity and Distribution*, 22, 358-370.

Published  
2024-12-30

Issue  
Vol. 3 No. 4 (2024)

Section  
Articles

License



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/).

UMYU Scientifica recognizes the importance of protecting authors' intellectual property while promoting the free exchange of scientific knowledge. The journal adopts a copyright-retention model that empowers authors to maintain ownership of their work while granting the journal rights necessary for publication and dissemination.

## 1. Copyright Ownership

Authors publishing with *UMYU Scientifica* retain full copyright and publishing rights to their work. By submitting a manuscript, authors agree to grant the journal a non-exclusive license to publish, reproduce, distribute, and archive the article in all forms and media for the purpose of scholarly communication.

## 2. Licensing Terms

All articles are published under the Creative Commons Attribution-NonCommercial (CC BY-NC) license.

This license permits others to:

- **Share** - copy and redistribute the material in any medium or format.
- **Adapt** - remix, transform, and build upon the material.
- **For non-commercial purposes only**, provided that proper credit is given to the original author(s) and *UMYU Scientifica* as the source, a link to the license is provided, and any modifications are clearly indicated.

Commercial reuse or distribution of the content requires **written permission** from both the author and the editorial office.

### 3. Author Rights

Authors are free to:

- Deposit all versions of their manuscript (preprint, accepted version, and published version) in institutional, disciplinary, or public repositories without embargo.
- Use and distribute their published article for **non-commercial scholarly purposes**, including teaching, conference presentations, and research sharing.
- Include their work in future books, theses, or compilations, provided proper citation to the journal is made.

### 4. Publisher's Rights

Upon publication, *UMYU Scientifica* retains the right to:

- Host, index, and disseminate the article through the journal's website and partner databases.
- Archive the content in long-term preservation systems such as the PKP Preservation Network (PKP-PN) and the Umaru Musa Yar'adua University Institutional Repository.

### 5. Attribution and Citation

Users must give appropriate credit to the author(s), include a link to the article's DOI or the journal webpage, and indicate if changes were made. Proper citation is required whenever the work is reused or referenced.

### 6. License Reference

For detailed terms of use, please refer to the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0):

<https://creativecommons.org/licenses/by-nc/4.0/>

#### How to Cite

Adama, S. B., Zakari, K., Lemu, M. M., Auta, Y. I., Saba, J. J., Samuel, P. O., & Mohammed, Y. M. (2024). Ichthyofauna Diversity and Morphometric Analysis in an Anthropogenically Stressed Water Body Downstream of Jebba Dam, Nigeria. *UMYU Scientifica*, 3(4), 337-344. <https://doi.org/10.56919/usci.2434.028>

More Citation Formats

### Similar Articles

- Akeem Babatunde Dauda, Ibraheem Kabir, Mansur Abdullahi Dambatta, Abdulsalam Muhammad Abdullahi, Kamisu Nuhu Muhammad, Hauwa Ohunene Sadiq, Morphometric Analysis and Growth Patterns of *Oreochromis niloticus* (Linnaeus, 1758) in Tiqa Damand River Wudil, Nigeria, UMYU Scientifica: Vol. 4 No. 3 (2025)
- Panda Ahmad Anas, Timothy Auta, Armaya'u Bichi Hamisu, Sadauki Amadu Mustapha, Comparative Assessment of Growth, Nutrient Utilization and Profitability of *Clarias gariepinus* (Burchell, 1822) Fed Locally Manufactured Feeds and a Foreign Feed, UMYU Scientifica: Vol. 4 No. 4 (2025)
- Abdulbari Dandashire Saleh, Baha'uddeen Salisu, Idris Isma'il Kariya, Nuru Abdullahi Nabage, Muhammad Sanusi Idris, Umar Sambo Umar, Abdulmajid Isa Jibrin, Alamin Ahmad Muhammad, Abubakar Sadiq Maigari, Ahmed Isah Haruna, Abdullahi Shehu Ibrahim, Geology and Economic Potential of Manawaji and Environs, Gulani Area, Upper Benue Trough, Nigeria, UMYU Scientifica: Vol. 4 No. 3 (2025)
- Yusuf Ibrahim, Abubakar Sani Bawale, Trend and Variability Analysis of Rainfall and Temperature in the Dagona Wetland, Nigeria (1993-2023), UMYU Scientifica: Vol. 5 No. 1 (2026)
- Fiddausi Dayyabu, Mustapha Zakariyya Karkarna, Nura Umar Kura, Surayya Farouk Durumin iya, Multivariate Hydrogeochemical Assessment of Urban Groundwater Quality in Kano Metropolis, Nigeria, UMYU Scientifica: Vol. 4 No. 3 (2025)
- Francis M Obiaju, Auta Timothy, Bemshima O James, Sergius T Ugwueke, Ehime Itama, Ecological and Human Risk Assessment of Heavy Metal Level in Water, Sediment and *Clarias gariepinus* from Zobe Dam, Katsina State, Northwestern Nigeria, UMYU Scientifica: Vol. 5 No. 1 (2026)
- Dini Sabo, Shehu Habibu, Magaji Ladan, Shuaibu Abdullahi, Daniel Eric, Development of Starch-based Biodegradable Plastics reinforced with Natural Fiber for enhanced mechanical and biodegradation properties, UMYU Scientifica: Vol. 4 No. 4 (2025)
- Yusuf Muhammad Bawa, Saghir Kalimullah, Nasir Hassan Wagini, Mohammed Adamu Garga, Mohammad Ahmad, Nuraddeen Muhammad Hassan, Nasiru Ibrahim, Abdulganiyu A Alabi, Ahmad Muhammad Daniyan, Sulaiman Muhammad, Bello Salisu, Ibrahim Ladan Garba, Richard Agyo-Likita, Concentration-Dependent Effects of Dimethoate, Lambda-Cyhalothrin and their Mixture on Germination, Vegetative Growth and Yield of Cowpea (*Vigna unguiculata*), UMYU Scientifica: Vol. 4 No. 4 (2025)
- Aliyu Abdulhamid Omar, Auwal Barde Alhassan, Alhaji Mohammed Usman, Umar Aliyu, Molecular identification of sibling species in the *Anopheles gambiae* complex using PCR in Bauchi State, Nigeria, UMYU Scientifica: Vol. 5 No. 1 (2026)
- Zaharaddeen Usman, Fatima Mukhtar, Baha'uddeen Salisu, Abdulbari Dandashire Saleh, Integrated Phytochemical Profiling (GC-MS/FTIR), Molecular Docking, and Bioevaluation of *Vernonia amygdalina* and *Psidium guajava* Against Multidrug-Resistant *Salmonella typhi*, UMYU Scientifica: Vol. 4 No. 4 (2025)

You may also [start an advanced similarity search](#) for this article.



[Submit Article](#)

[Aims & Scope](#)

[Author Guidelines](#)

[Processing Charge\(s\)](#)

[Open Access Policy](#)

Copyright Policy

Digital Preservation

Ethics & Malpractice

Dealing of Misconduct

Platform &  
workflow by  
OJS / PKP

ORIGINAL RESEARCH ARTICLE

## Ichthyofauna Diversity and Morphometric Analysis in an Anthropogenically Stressed Water Body Downstream of Jebba Dam, Nigeria

Adama Solomon Bake<sup>1</sup>, Zakari Kasim<sup>1</sup>, Lemu Mohammed Mohammed<sup>2</sup>, Auta Yohannah Iliya<sup>1</sup>, Saba Jonathan Jiya<sup>1</sup>, Samuel Patrick Ozovehe<sup>1</sup> and Mohammed Yakubu Manbe<sup>1\*</sup>

<sup>1</sup>Department of Animal Biology, Federal University of Technology, Minna, Niger State, Nigeria

<sup>2</sup>Department of Zoology, Usmanu Danfodiyo University, Sokoto, Sokoto State, Nigeria

<sup>3</sup>Niger State College of Agriculture Mokwa, Niger State, Nigeria

<sup>4</sup>Department of Biology, Ibrahim Badamasi Babangida University, Lapai, Niger State, Nigeria

### ABSTRACT

Fishery resources are on the decline due to incessant and unregulated fishery activities practiced in most parts of the globe. This study investigated the fish composition of an anthropogenically stressed water body in Nigeria (Downstream Jebba Dam). Samples were collected using standard methods and procedures. A total of 6,850 fish belonging to 17 families and 25 species were collected from the sampled stations. Fish composition shows the Mochokidae family with the most dominant species at 29.33% and represented *Synodontis batensoda*, which was well distributed in all the study stations but highest in the station with 934 individuals, while the Gymnarchidae family was the least recorded with 0.16% and represented *Gymnarchus niloticus*. Length-Weight relationship (LWR) value ranged between 0.0018 and 0.215 ( $b < 3$ ) for the dominant fish species, indicating negative allometric growth as the fish species grows in length faster when compared to the weight. In terms of mean condition factor (K of the dominant fish species of Downstream Jebba Dam, *Cynothrissa mento* had the highest with 10.09, followed by *Oreochromis niloticus* (9.82), *Mugil cephalus* (9.02), *Labeo coubie* (3.72), *Bagrus bayad* (0.90), *Alestes macrolepidotus* (0.52), *Clarias laticeps* (0.31), *Momyrus rume* (0.26), *Clarias gariepinus* (0.19) and *Citharus citharus* (0.16). The high condition factor ( $> 1$ ) observed in some fish indicates that these fish species were physiologically stable and successful well, while those with less than ( $< 1$ ) mean value implied that it was physiologically unstable, as showed in some species. Overall, the medium abundance of fish species in this study is an indication of an impaired water system.

### INTRODUCTION

Fish populations or abundance respond to various factors such as unregulated fishing, point and non-point source pollution, eutrophication, and land use type, which can result in variation in the sizes, species assemblage, distribution, and abundance of fish in aquatic habitat environments (Mohammed *et al.*, 2021a; Ibrahim *et al.*, 2022). Favourable environmental conditions are another factor that is responsible for fish species abundance in water bodies globally (Froese and Pauly 2018). The freshwater ecosystem of Nigeria is on the decline due to numerous anthropogenic activities around water bodies coupled with the land used type (Mohammed *et al.*, 2020; Mohammed *et al.*, 2021b; Oladipo *et al.*, 2021). Due to overexploitation and poor management of inland water bodies, Nigeria's fisheries resources are currently declining (Adamu *et al.*, 2021a,b). To ensure the sustainability of fishery resources, it is necessary to have a sufficient

understanding of the species composition and their diversity with abundance in major water bodies in the country (Adadu *et al.*, 2019; Adamu *et al.*, 2021a,b). One of the most important factors that contribute to changes in fish diversity and composition in freshwater bodies of Nigeria is indiscriminate and unregulated fishing, which exerts significant pressure on the fishery composition of Nigeria (Adadu *et al.*, 2019; Adamu *et al.*, 2021a,b; Oladipo *et al.*, 2021). Due to anthropogenic such as industrialization, urbanisation, and farming activities coupled with pressure from artisanal fishermen in our water bodies, fishery resources are declining, and fishes are migrating downstream in most cases in need of food and good habitat for spawning (Kumar and Asiji, 2009).

The development of the Jebba can be attributed to the building of hydro-electric power plants (Dam) in 1975.

**Correspondence:** Yakubu Manbe Mohammed. Department of Biology, Ibrahim Badamasi Babangida University, Lapai, Niger State, Nigeria. ✉ [yakubmohammedmanbe@yahoo.com](mailto:yakubmohammedmanbe@yahoo.com)


**How to cite:** Adama, S. B., Zakari, K., Lemu, M. M., Auta, Y. I., Saba, J. J., Samuel, P. O., & Mohammed, Y. M. (2024). Ichthyofauna Diversity and Morphometric Analysis in an Anthropogenically Stressed Water Body Downstream of Jebba Dam, Nigeria. *UMYU Scientifica*, 3(4), 337 – 344. <https://doi.org/10.56919/usci.2434.028>

### ARTICLE HISTORY

Received February 24, 2024  
Accepted November 23, 2024  
Published December 13, 2024

### KEYWORDS

Fish composition, Jebba dam, Length-weight relationship



© The authors. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>)

Even though the Jebba Dam was built to raise the living standard of the public, the aquatic ecosystem and the biotic community are at risk due to the rate at which anthropogenic activities are increasing within the water body (Oladipo *et al.*, 2021). This has resulted in the degradation of habitat and, eventually, in the disturbance of the water discharge's spatiotemporal pattern. The region's fisheries resources are frequently threatened by overexploitation, ongoing, indiscriminate fish harvesting, and pollution (Oladipo *et al.*, 2021). These dangers may have the unintended consequence of reducing the size of the population, trawling fish, and gene pool depletion (Aliko *et al.*, 2010).

In Nigeria and throughout Africa, fish are among the greatest and most important sources of animal protein for diet (Adadu *et al.*, 2019; Adamu *et al.*, 2021a,b; Mohammed *et al.*, 2021a). Different biotic and abiotic condition influence fish species assemblages and their diversity in streams and rivers. As the human population keeps increasing, fish species in the River Niger are being lost due to numerous anthropogenic factors such as using of chemicals for fishing, overfishing, using undersized marshes, spraying herbicides by rice farmers, and disposing of waste materials (Adadu *et al.*, 2019; Adamu *et al.*, 2021a,b; Mohammed *et al.*, 2021a). Different studies

on fish assemblage have been carried out around the Jebba Hydro-electric Dam (Oladipo *et al.*, 2021) using both taxonomic and molecular methods, but no recent data on the morphometric composition of fishes downstream of Jebba Dam. Thus, this study is designed to assess the composition, diversity, distribution, and Morphometric of fish in the River Niger Downstream Jebba dam.

**MATERIALS AND METHODS**

**Description of the study area**

The study was carried out downstream of Jebba dam, Niger State; River Niger surrounds Jebba town. The northern part is called Gungun, in Mokwa local Government Niger state, and south in Kwara State Moro local Government area. Jebba Dam falls within the Guinea savannah of the Northcentral part of Nigeria, characterized by two (Dry and wet) seasons (Adelakun *et al.*, 2017). The river is located between longitude 4°49'34.11"E and latitude 9°7' 9.31"N or 4°82'61.42 and 9°11'92.52 respectively. The majority of the people in the studied are artisanal fishermen and traders while few of them engage in agricultural activities around the river bank.




Figure 1: Map of River Niger showing the four stations

**Table 1: Fish species composition and distribution in Downstream Jebba Dam (November 2021-September 2022).**

Family	Taxa	Stations				Total	%
		1	2	3	4		
Centropomidae	<i>Lates niloticus</i> (Linnaeus, 1758)	30	16	19	0	65	0.95
Cyprinidae	<i>Labeo coubie</i> (uppel, 1832)	431	538	472	30	1471	21.47
Characidae	<i>Hydrocynus forskali</i> (Cuvier, 1819)	52	80	73	0	205	2.99
	<i>Alestes macrolepidotus</i> (Valenciene,1846)	53	49	46	0	148	2.16
	<i>Alestes baremose</i> (Leach, 1825)	59	73	98	0	230	3.36
Citharinidae	<i>Citharinus citharus</i> (Geoffrey, 1809)	51	60	38	2	151	2.20
Gymnarchidae	<i>Gymnarchus niloticus</i> (Cuvier, 1829)	2	7	2	0	11	0.16
Malapteruridae	<i>Malapterurus electricus</i> (Gmel, 1789)	7	8	2	0	17	0.25
Schilbeidae	<i>Eutropius niloticus</i> (Gunther, 1864)	17	11	21	0	49	0.72
Bagridae	<i>Bagrus bayad</i> (Forska, 1775)	65	60	51	2	178	2.60
	<i>Bagrus doermai</i> (forskal, 1775)	0	13	19	0	32	0.47
Claroteidae	<i>Clarotes laticeps</i> (Ruppel,1829)	27	48	48	1	124	1.81
	<i>Auchenoglanis biscutatus</i> (Ruppel, 1829)	0	27	0	0	27	0.39
Clariidae	<i>Clarias gariepinus</i> (Burchell, 1822)	76	84	44	27	231	3.37
Cichlidae	<i>Tilapia zillii</i> (Gervil,1848)	12	24	0	0	36	0.53
	<i>Oreochromis niloticus</i> (Linnaeus, 1758)	264	275	196	11	746	10.89
	<i>Sarotherodon</i> (Linnaeus, 1758)	34	15	23	0	72	1.05
Mormyridae	<i>Mormyrus rume</i> (Cuvies, 1846 )	38	69	57	7	171	2.50
	<i>Hyperopisus bebe</i> (Gunther, 1886)	11	11	11	11	44	0.64
	<i>Mormyrops deliciosus</i> (Leach, 1818)	2	7	3	0	12	0.18
Osteoglossidae	<i>Heterotis niloticus</i> (Cuvier,1829)	5	3	9	3	20	0.29
Mugilidae	<i>Mugil cephalus</i> (Lacepede, 1803)	198	170	179	0	547	7.80
Potamotrygonidae	<i>Stingrays</i> (Valenciennes,1846)	7	3	8	2	20	0.29
Clupeidae	<i>Cynothrissa mento</i> (Johnels, 1954)	87	54	93	0	234	3.42
Mochokidae	<i>Synodontis batensoda</i> (Daget, 1954)	645	934	383	47	2009	29.33
Total		2173	2639	1895	143	6,850	100.0

**Diversity indices of fish species in Downstream Jebba Dam.**

Table 2 shows the diversity indices of fish species downstream of Jebba Dam. The highest species richness (25) was recorded at station 2, followed by stations 1 and 3 with 23 species, and then station 4 with 11 species. The number of individuals was highest in station 2 (2639), followed by station 1, with (2173) species, then station 3 (1895), and (143) species at station 4. Station 2 had the highest margalef index (3.05), stations 1 and 3 were almost the same in species composition (2.86) and (2.92),

respectively, while station 4 had the least margalef index (2.02). The Shannon-Weiner index was highest in station 3 (4.01), followed by station 1 (3.39), while station 2 was (3.13), and the lowest was in station 4 (2.50). Simpson’s dominance was recorded highest in station 2 (0.19), than station 1 with (0.16), while station 3 was (0.13), and the lowest was recorded at station 4 with (0.12). The Evenness index indicated that no significant difference between stations 1 and 2 when (p>0.05) (0.38 and 0.36). The highest evenness was in station 3 (4.01), while station 4 was the least even (2.50).

**Table 2: Diversity indices of fishes in downstream Jebba Dam.**

Indices	Station 1	Station 2	Station 3	Station 4
Taxa (Species)	23	25	23	11
Number of individual fish	2173	2639	1895	143
Simpson dominance	0.16	0.19	0.13	0.12
Shannon index (H)	3.39	3.13	4.01	2.5
Evenness (E)	0.38	0.36	0.45	0.28
Margalef’s index (D)	2.86	3.05	2.92	2.02

**Morphometrics of selected fish species in downstream Jebba dam.**

Table 3 shows the length-weight relationship and condition factor of dominant fish species in the sampled stations of Downstream Jebba Dam. The length-weight

relationship (r) of all the sampled fish from Downstream Jebba dam ranged from 0.018 in *Clarotes laticeps* to 0.215 in *Labeo coubie*. The Length-Weight relationship (LWR) of the dominant fish species ranged between 0.0018-0.215, and the observed “b” value was <3; thus the allometric growth was negative. In term of mean condition factor (K



### Sampling stations

Station 1 is popularly called Inda doku, and it is located below Jebba Bridge, close to the Jebba water board. This station has a less human settlement with high anthropogenicity like farming, washing of cars, swimming, domestic uses, washing of clothes, and plates, among others. Station 2 is called Kpata Sarki by the locals, and it is located 200 meters from the Jebba Paper mill industry. This station is characterized by some natural vegetation with scattered human settlements, and several anthropogenic activities such as farming, grazing of livestock, and laundry activities are common in this site. Station 3 is called Tsamanagi, and it is located 200 meters away from Jebba Paper Mill. Agricultural activities and other anthropogenic activities at the river bank characterize this station. The river bed is dominated by projected rocks, alluvia deposits, sand, and mud at some parts of the site. Station 4 is called Panga, and it is approximately 200 meters away from station 3. This station is characterized by low human settlement with more agricultural activities such as farming and fishing

### Collection of fish sample

The fish data and samples were collected from the fishermen at different major landing sites between 8 am to 10 am, and the sites were sampled weekly. For the periods of seven months (November 2021 to January 2022 and June 2022 to September 2022), making up three months in the dry season and four months in the wet season. Fish sampling was by direct observation of the fish at the landing sites, and collected fish were sorted into taxonomic groups, counted, and identified from family to species level by use of an identification guide and each species of the fishes collected was preserved and transported in a cool box to the laboratory of Animal Biology department of Federal university of technology Minna.

### Determination of the morphometrics of fishes

Standard Length (S.L): It is the measurement of the distance between the tail fin to the tip of the fish's mouth using tape or thread (Wu and Huang, 2017). Total Length (T.L) is the measurement from the tip of the fish's mouth to the tail tip; thread can be used and later measured on a ruler or tape on a flat surface (Wu and Huang, 2017). The body weight of the dominant fish species was measured to the nearest (0.1g) with a portable weighing balance (Adelaku *et al.*, 2017)

### Data analysis

All data collected were typed in Microsoft Excel 2016 edition. The fish assemblage and distribution were subjected to diversity indexes such as Shannon-Weiner index (H), Simpson dominance index (C), and Margalef species richness index (D), while the length-weight relationship was subjected to descriptive statistics. Also, the Regression coefficient (b), correlation coefficient (r), and the condition factor of the dominant fishes were all

## RESULTS

### Fish species composition and distribution in downstream Jebba Dam.

Table 1 shows a total of 6,850 fish belonging to 17 families and 25 taxa encountered during the study period in downstream Jebba Dam, Kwara state. Fish species composition and distribution in this study shows Mochokidae family was the dominant species with 29.33% and represented *Synodontis batensoda*, which was distributed across all the studied stations but highest in station with 934 individuals, while the Gymnarchidae family was the least recorded with 0.16% and represented *Gymnarchus niloticus*. The family Cyprinidae recorded 21.47% of the total catch and was represented by *Labeo conbie* which was well distributed in all stations. The Characidae family recorded about 8.51% of all the total catch and was represented by 3 species, namely, *Aleste baremose*, *Aleste macrolepidotus*, and *Hydrocynus forskali*. *Aleste baremose* was the most dominant species in this family, and none of them was recorded in station 4. The Mugilidae family was representative of *Mugil cephalus* only and accounted for 7.8% of the total catch, which was distributed in all stations. The family Centropomidae was represented with one species (*Lates niloticus*), and it account for 0.95% of the total catch. Malapteruridae was represented by *Malapterurus electrical*, which accounts for 0.25% of the fish catch and was only found in stations 1,2 and 3. Osteoglossidae was represented by *Heterotis niloticus* only, which accounted for 0.29% of the total catch, and it was distributed sparsely in all stations. Clariidae family also recorded one species which is *Clarias gariepinus*, which accounts for 3.37% of the total catch and is evenly distributed in all stations. Family Gymniridae was represented by just a species (*Sting rays*), which accounted for 0.29% of the total catch and were slightly distributed in all stations. The family Claroteidae was represented by two species (*Anchenoglanis biscutatus* and *Clarotes laticeps*), which account for 2.2% of the total catch. The family Schilbeidae was represented by *Eutropius niloticus* and accounted for 0.72% of the total catch recorded. The family Cichlidae accounts for 12.47% of the total catch and was represented by 3 species, namely *Tilapia zilli*, *Oreochromis niloticus*, and *Sarotherodon* sp. The family Clupeidae accounted for about 3.42% of the total catch and was represented by *Cynothrissa mento*. The family Mormyridae accounted for about 3.32% of the total catch, and it was represented by 3 species, namely, *Mormyrus rume*, *Mormyrus deliciosus*, and *Hyperopisus bebe*. The family Bagridae accounted for about 3.07% of the total catch and was represented by 2 species, namely, *Bagrus bayad* and *Bagrus doemac*. The family Citharinidae was represented by *Citharinus citharus* and accounted for 2.20% of the total catch. The family Osteoglossidae was represented by *Heterotis niloticus* only, which account for 0.29% of the total catch.

of the dominant fish species of Downstream Jebba Dam, *Cynothrissa mento*, had the highest with 10.09, followed by *Oreochromis niloticus* (9.82), *Mugil cephalus* (9.02) *Labeo coubie* (3.72), *Bagrus bayad* (0.90), *Alestes macrolepidotus* (0.52), *Clarote laticeps* (0.31), *Momyrus rume* (0.26), *Clarias gariepinus*

(0.19) and *Citharus citharus* (0.16). The high condition factor (>1) observed in some fish indicates that these fish species were physiologically stable and successful well, while those with less than (< 1) mean value implied that it was physiologically unstable, as showed in some species.

**Table 3: Mean Condition factors (K) of some Dominant fish species in Downstream Jebba dam.**

Fish Species	Mean condition (K) factors	Range	Length-Weight Relationship
<i>Momyrus rume</i>	0.26	0.18-0.46	0.025
<i>Citharus citharus</i>	0.16	0.25-0.37	0.022
<i>Labeo coubie</i>	3.72	1.25-10.94	0.215
<i>Bagrus bayad</i>	0.90	1.67-1.92	0.026
<i>Clarias gariepinus</i>	0.19	0.17-0.4	0.034
<i>Alestes macrolepidotus</i>	0.5	0.52-0.90	0.021
<i>Cynothrissa mento</i>	10.09	7.87-16.80	0.034
<i>Clarotes laticeps</i>	0.31	0.0-1.25	0.018
<i>Oreochromis niloticus</i>	9.82	0.58-16.56	0.109
<i>Mugil cephalus</i>	9.02	6.40-18.45	0.080

**DISCUSSION**

The results of this study show the abundance of fish in downstream Jebba Dam, which could be due to good water quality, which enhances good breeding as most fish species prefer less disturbed water for spawning. A total of 6,850 fish specimens from 25 species and 17 families were identified from the four sampled stations of Downstream Jebba Dam. In this study, the fish population decreases downward the sampling station 4, which has high anthropogenic activities. Additionally, more fish were captured in this study when compared to the finding of Agbugui *et al.* (2019), who reported 706 fish from 18 families and 35 species at the three sites at River Niger in Agenebode, Edo state. Okomoda *et al.* (2012) also report the presence of 13 families and 28 species in the Lower River Niger near Idah in Kogi State.

Additionally, Also Lawal *et al.* (2023) reported a total of 8273 fish from Mairua Reservoir Water, Funtua, Katsina State, North-Western Nigeria. Oladipo *et al.* (2021) also reported the patterns of ichthyofaunal diversity and distribution across the Jebba Hydro-Electric Power dam and reported a total of 9605 freshwater fishes were recorded during the sampling period. Abiodun and Odunze (2011) noted that Jebba Lake, Nigeria, is home to 51 species from 12 families in their findings. The variation in the number of families and species found in a certain river, as well as in other rivers, can be ascribed to seasonal variations, environmental shifts, and human activities that lead to fish migration and relocation from one place to another (Sikoki *et al.*, 2008). At every sampling station, *Synodontis batensoda* was the most dominant fish, and this shows that the species is tolerant of environmental conditions such as low temperature, conductivity, and pH, as well as water pollution (Wubie *et al.*, (2017). Mochokidae and Cichlidae were the most abundant family in this study. Oladipo *et al.* (2021) earlier reported the abundance of these families in the Jebba Hydro-Electric Power dam. Also, Abiodun and John (2017) reported the abundance of Mochokidae and Cichlidae lower River

Niger basin. These families' supremacy may be related to their great degree of environmental adaptability and prodigious breeding skills (Mustapha, 2010).

The abundance of the family Cyprinidae in this study is consistent with the findings of Rusmilyausari *et al.* (2021), who observed a similar abundance in the Martapura River in South Kalimantan, Indonesia. According to Buwono *et al.* (2017), there are comparatively many species in the freshwater Cyprinidae family. In line with the assertion made by Sulaiman *et al.* (2018), 40 of the 104 freshwater species discovered in Brunei Darussalam are members of the Cyprinidae family. Comparably, Ning *et al.* (2016) also reported the abundance of the Cyprinidae family in the Ganjiang River. *Tilapia zilli*, *Oreochromis niloticus*, and *Sarotherodon* sp are the representatives of the Cichlidae family in this study. The results of this study, which focused on the dominance of cichlids, in particular *Oreochromis niloticus*, are consistent with many other findings from Nigeria. Nazceef *et al.* (2021) studied the biodiversity of fish species in the Dadin-Kowa reservoir in Gombe State and reported the presence of the chichlidae family. Also, chichlidae abundance has been reported by many studies in Nigeria (Solomon *et al.*, 2017; Dienne *et al.*, 2018; Pius *et al.*, 2020; Abdulkarim *et al.*, 2020; Olopade, 2020; Oladipo *et al.*, 2021; Omoike, 2021; Lawal *et al.*, 2023) who all documented the dominance of this species. The dominance of cichlid fish can be attributed to several noteworthy features, such as their rapid rate of proliferation and their status as the second most varied family of freshwater fish (Seehausen, 2015). In the freshwater ecology of Nigeria, cichlids have been observed to be the most abundant group (Mustapha, 2010). Comparably, it has been reported that the lower River Niger basin is home to large populations of Cichlidae and Mochokidae (Atile *et al.*, 2016; Abiodun and John, 2017). These families' supremacy may be related to their great degree of environmental adaptability and prodigious breeding skills (Mustapha, 2010). The Characidae family represented prominent species like *Alestes baremose*, *Hydrocynus forskali*, and *Alestes macrolepidotus*



were also abundant in this study. This exceptional abundance is explained by the species' adaptability to a variety of feeding habits, their body size in relation to speed of growth, fecundity level, and availability of food sources (Pius *et al.*, 2020).

Mormonidae abundance was low when compared to the report of Abiodun and Isiba (2017), who reported the abundance of this group in Lower River Niger, Idah, Kogi State. *Malapterurus electricus* is the only representative of the family Malapteruridae. The presence of this family conforms to the report of Mo *et al.* (2019) and in contrast with the reports of Essien Iboke (2020), who reported their absence in their findings. *Potamotrygon motoro* was the only taxon that represented the family Potamotrygonidae which account for 0.29% of the composition. *Entropius niloticus*, belonging to the family Schilbiidae was also documented in this study. *Clarias gariepinus* is the only representative of the Clariidae family found in all of the studied stations. Nazeef *et al.* (2021) also reported their presence in all the sampled stations of the Dadin-Kowa reservoir. *Gymnarchus niloticus*, the only taxa represented in the family Gymnarchidae, was the least abundant group in this study. This is in contrast to the findings of Agbugui *et al.* (2019), who reported that the family Gymnarchidae was the most abundant, with 8.49% of only specie *Gymnarchus niloticus* in River Niger at Agenebode, Edo State, Nigeria.

The chosen fish species' length-weight correlation coefficient ( $r=0.71$ ), which was recorded at the downstream Jabba Dam, revealed a substantial positive connection between length and weight, indicating that weight increases as length increases. Positive correlations were also reported by Solomon *et al.* (2012) and Benedict *et al.* (2009) in Cross River wetlands. Similarly, Leonard *et al.* (2011) reported a good correlation ( $r = 0.874; 0.973$ ) between Buyo and Ayame Reservoirs and Ezekiel and Abowei (2014) in the Niger Delta's Amassoma flood plain. The variances in the values could be caused by differences in the environment, location, time of year, and the sizes and numbers of species sampled. The positive association demonstrated that the growth pattern did not override the significant link between fish weight and length. These observations may also be the result of variations in the quantity and quality of fish present, food availability, and water quality. The Length-Weight relationship (LWR) 0.022-0.035 data for fish species indicate that "b" is less than three for every species. A negative allometric growth pattern was seen in this study. Abowei and Hart (2014) also reported negative growth patterns of some fish species in the lower River Nun in the Niger Delta. Negative allometric growth was also reported by Olowo *et al.* (2022) from fish species from the Ovia River, Edo State, Nigeria. The study's condition factors varied according to the species: *Citharinus citharus* had the lowest condition factor (0.16), while *Cynothrissa mento* had the greatest condition factor (10.09). Olowo *et al.* (2022) reported positive condition factors from fish species from the Ovia River, Edo State, Nigeria. The variations may result from variations in the physical and chemical

characteristics of the environments, the plant and animal populations, the weight of the individual species studied, the time of the sampling, and the availability of food.

## CONCLUSION

The fish species found downstream of the Jabba Dam show a total of 6,850 fish from 17 families, and 25 species were seen throughout the study period, indicating a less perturbed water body, and the river continues to provide good habitat for the development and preservation of fish populations. Also, the length-weight relationship of a few dominating species, such as *Alestes macrolepilotus*, *Mugil cephalus*, *Cynothrissa mento*, *Labeo conbie*, and *Synodontis batensoda*, had a strong positive link, as indicated by the correlation coefficient ( $r$ ) value, but the regression coefficient ( $b$ ) value indicated a negative allometric growth rate. The study also found that the condition factor of a few dominant fish species was physiologically stable and successful in the aquatic environment, with the exception of the species that showed values below 1, which indicates that they are not physiologically stable. Overall, the studied river shows a medium fish production potential in the studied river.

## REFERENCES

- Abdulkarim, M., Magaji, I.M., Yusuf, Z.A. & Magaji, N.S. (2020). diversity and spatio-temporal distribution of fish species in Tiga Dam, Kano state. *Nigerian Journal of Fisheries*, 17(1), 1969-1972.
- Abiodun, J.A. & John P. (2017). Biodiversity and abundance of Fish and some processing methods in Lower Niger River, Idah, Kogi State Nigeria. *Nigeria Journal of Fisheries and Aquaculture*, 5(2), 20-25.
- Abiodun, J.A. & Odunze, F.C. (2011). Fish diversity and fisheries of Jebba Lake Nigeria. *Nigeria journal of fisheries*, 8(2), 284-290.
- Abowei, F. N. & Hart, A. I. (2014). Some morphometric parameters of 10 finfish species from the lower Nun River, Niger Delta, Nigeria. *Research Journal of Biological Sciences*, 4(3), 282-288.
- Adadu, M.O., Garba, A.A. & Mshelia, J.P. (2019). Diversity and abundance of fish species in river Okpokwu, Benue State, Nigeria. *International Journal of Fisheries and Aquatic Studies*, 7 55-60
- Adamu, K. M., Mohammed, Y. M., Muhammad, H., Achebe, A. C. & Jeremiah, J. (2021a). Morphometric and biomarker indices of fishes in Dangana Lake, Lapai, Niger State. *Direct Research Journal of Biology and Biotechnology*, 7, 16-27. [Crossref].
- Adamu, K.M., Kuyembo, S.M., Abbas, A.H., Mohammed, Y.M., Djadjiti, N. & Mangse, G. (2021b). Growth performance of African Catfish, Spinach and Lettuce in an aquaponic system. *Dutse Journal of Agriculture and Food Security*, 8(2), 96-106.
- Adelakun, K.M., Mustapha, M.K., Amali, R.P. & Mohammed, N. (2017). Seasonal variation in nutritional quality of Catfish (*Clarias gariepinus*)

- from Upper Jebba Basin, Nigeria. *Journal of Nutrition & Food Sciences*, 7(5), 1-4.
- Agbugui, M. O., Abbulimen, E. F., Inobeme, A. & Olori, E. (2019). Biodiversity of Fish Fauna in River Niger at Agenebode, Edo State, Nigeria. *Egyptian Journal of Aquatic Biology & Fisheries*, 23(4), 159-166. [Crossref]
- Aliko, N.G.G., Da costa, K.S., Dietoa, Y.M., Ouattara, A. & Gourene, G. (2010). Caractéristiques de la population de *Distichodus mustratus* Günther, 1864 (Pisces: Distichodontidae) du lac de barrage de Taabo (bassin du Bandama, Côte d'Ivoire). Implications pour une gestion rationnelle du stock. *Tropicalicultura*, 28, 50-56.
- Atile, J.I., Shima, J.N. & Akombo, P.M., (2016). Food and feeding, length-weight and condition factor of the catfish *Synodontis membranaceus* (Etiene Geoffroy Saint Hilaire, 1809) (Osteichthyes: Mochokidae) from Lower Benue River, Makurdi, Nigeria. *Agriculture, Forestry and Fisheries*, 5(4), 87-96. [Crossref]
- Benedict, O., Offem, Yemi, A. S. & Omoniyi, I. T., (2009). Length-weight relationship condition factor and sex ratio of forty six important fishes in a Tropical Flood River. *Research Journal of Fisheries and Hydrobiology*, 4(2), 65-72, 2009.
- Buwono, N.R., Fariedah, F. & Anestyningrum, R.E. (2017). Komunitas Ikan di Sungai Jerowan Kabupaten Madiun. *Journal of Aquaculture and Fish Health*, 6(2), 81-88. [Crossref]
- Dienye H.E, Olopade, O.A. & Toby, S.A., (2018) Species Composition and Diversity of Cast net fisheries in new Calabar, Niger Delta, Nigeria. *Journal of Biodiversity, Conservation & Bioresources Management*, 4(1), 19-26. [Crossref]
- Essien-Ibok, M. A. (2020). Fish fauna of Akwa Ibom State inland waters. *Biodiversity International Journal*, 4(2), 82-88.
- Ezekiel, E. N. & Abowei, J. F. N., (2014). A study of length-weight relationship and condition factor of *Hepsetus odoe* (Bloch, 1794) from Amassoma flood plains. *Annals of Biological Sciences*, 2(2), 10-17.
- Froese, R. & Pauly, (2018). *Diversity of Fish fauna*. Retrieved 4/6/2018.
- Ibrahim, M. I., Ibrahim, Y., Abdullahi, A. M., Najibullah, B. A, Obi, P. U. & Mohammed, Y. M. (2022). Bacteria associated with smoked catfish sold at Bida Modern market, Northcentral Nigeria. *International Journal of Fisheries and Aquatic Studies*, 10(2), 38-40. [Crossref]
- Kumar, U. & Asija, M. J. (2009). *Biodiversity: Principle and Conservation*. Agrobios (India)
- Leonard, T., Gouli, G. B. and Kouassi, S. Da Costa (2011). Length-weight relationships for 36 freshwater fish species from two tropical reservoirs: Ayamé I and Buyo, Côte d'Ivoire. *Rev. Biol. Trop. (Int. J. Trop. Biol.*, 60 (4), 1847-1856. [Crossref]
- Lawal, N., Abdullahi, S.A. & Abolude, D.S. (2023). Physicochemical Characteristics and Fish Abundance and Diversity of Mairua Reservoir Water, Funtua, Katsina State, North-Western Nigeria. *Journal of Applied Science and Environmental Management* 27(1), 125-132. [Crossref]
- Leonard, T., Gouli, G.B. & Kouassi, S.D. (2011). Length-weight relationships for 36 freshwater fish species from two tropical reservoirs: Ayamé I and Buyo, Côte d'Ivoire. *International Journal of Tropical Biology*, 60(4), 1847-1856. [Crossref]
- Mo, Y., Wan, H., Cai, Y., Zhang, X., Li, R. & Wang, Y. (2023). Re-establishing fish migration channel of large reservoirs in Jinsha River Basin of China by using an eco-friendly reservoir operation method, *Journal of Hydrology: Regional Studies*, 47, 101412, [Crossref]
- Mohammed, Y. M., Abubakar, M. D., Muhammad, A. M., Muhammad, A. S., Umar, B. L., Hadizat, M. & Achebe, A. C. (2021a). Stomach Content of *Tilapia zilli* and *Oreochromis niloticus* from Wanzun River, Lavun Local Government, Niger State Nigeria. *Asian Journal of Fisheries and Aquatic Research*, 13(5), 15-20. Article no. AJFAR.70239. [Crossref]
- Mohammed, Y.M., Adamu, K.M., Ibrahim, B.U., Danjuma, S. & Amuzat, A.O. (2023). Exploring the Impact of Human Activities on Water Quality Characteristics of a Rural River in North-Central Nigeria using Macroinvertebrates Structural Assemblage. *West African Journal of Life Sciences*, 1, 023-034. [Crossref]
- Mohammed, Y.M., Arimoro, F. O., Ayanwale, A.V., Adama, B. S., Keke, U.N., Auta, Y. I. & Umar, B.L. (2020). Seasonal changes in the abundance of benthic macroinvertebrates & physico-chemical condition of Moussa stream Bida, Nigeria. *Tropical Freshwater Biology*, 29(1), 57-70. [Crossref]
- Mohammed, Y.M., Arimoro, F.O., Ayanwale, A.V. Adamu, K.M., Keke, U.N., Abubakar, M.D. & Achebe, A.C. (2021b). The current state of water quality and benthic invertebrate fauna in Chikke Stream (North-Central Nigeria). *Ukrainian Journal of Ecology*, 11(3), 26-34. [Crossref]
- Mustapha, M.K. (2008). Assessment of the Water Quality of Oyun Reservoir, Offa, Nigeria, Using Selected Physico-Chemical Parameters. *Turkish J Fish Aqua Sci*. 8, 309-319.
- Mustapha, M.K. (2010). Fish fauna of Oyun reservoir, Offa, Nigeria. *Journal of Aquatic Sciences*, 25 (1), 106-114.
- Nazeef, S., Ja'afar A., Abubakar K.A., and Kabiru M., (2021). Fish species biodiversity of Dadin-Kowa reservoir. *World Journal of Advanced Research and Reviews*, 11(02), 100-111. [Crossref]
- Okomoda, V.T., Solomon, S.G. & Aladi, S.L. (2012). Fish Fauna in Lower River Niger at Idah in Kogi State. *Journal of Agricultural & Veterinary science*, 4, 34-40.
- Oladipo, S.O., Nneji, L.M., Iyiola, O.A., Nnejid, I.C., Ayoola, A.O., Adedokun, K.M., Anifowoshe, A.T. Adeola, A.C. & Mustapha, M.K. (2021).

- Patterns of ichthyofaunal diversity and distribution across Jebba Hydro-Electric Power (HEP) dam, Jebba, north-central Nigeria. *Brazilian Journal of Biology*, 81(2), 258-2. [Crossref]
- Olopade O.A., Dinye H.E., Bamidele N. & Oderhohwo, O. (2020). Catchability and diversity of fish species captured by gill net in new Calabar River, Nigeria. *Indonesia Fishery Research Journal*, 26(2), 107-117. [Crossref]
- Olowo, U.C., Egun, N. K., Ehigiator, A.P. & Oboh, I.P. (2022). Growth analysis and sex ratio of fish species from the Ovia River, Edo State, Nigeria. *BIOLOGIJA*. 2022. Vol. 68. No. 2. P. 110–120. [Crossref]
- Omoike, A. (2021) The Trend in Fish Species Diversity and Abundance at Asejire Reservoir South Western Nigeria. *HSOA Journal of Aquaculture & Fishery*, 5(040), 1-9. [Crossref]
- Pius, D.E., Ja'afaru, A., Abubakar K.A., Zira J.D., Kefas, M. & Barau B.W. (2020). Fish biodiversity and abundance in river Taraba, Taraba State, Nigeria. *Ethiopian Journal of Environment Studies & Management*, 13(5), 535-544.
- Rosa, R.S., Charvet-Almeida, P. & Quijada, C.C.D. (2010). Biology of the South American Potamotrygonid Stingrays, p. 241-285. [Crossref]
- Rusmilyansari, A. A., Wahab, W. & Reika, C. (2021). Fish species composition and diversity in a river, a swamp, and a reservoir in Banjar District, South Kalimantan Province. *AAACL Bioflux*, 14(1), 412-423.
- Sechhausen, O. (2015). Process and pattern in cichlid radiations – inferences for understanding unusually high rates of evolutionary diversification. *New Phytology*. 207: 304-312. [Crossref]
- Sikoki, F.D., Zabbey, N. & Anyanwu, N. (2008). Fish Assemblage of Onu –Iyi-Ukwu Stream in Southern Tropical Nigeria. *Fish Biology*, 17(2), 13-23. [Crossref]
- Solomon, S.G, Ayuba V.O, Tahir, M.A. & Okomoda, V.T. (2017). Abundance composition of fish in lake kalgwai Jigawa State, Nigeria. *Jordan Journal of Agricultural Sciences*, 13, 45-54. [Crossref]
- Solomon, S.G, Okomoda, V.T. & Aladi. S.L. (2012). Fish Fauna in Lower River Niger at Idah in Kogi State. *Journal of Agricultural and Veterinary Science*, 4, 34-40.
- Sulaiman, Z., Hu T. H. & Lim K.K.P. (2018). Annotated checklist of freshwater fishes from Brunei Darussalam, Borneo. *Zootaxa*, 4379(1), 24–46. [Crossref]
- Uduma, A.U. (2018). Physico-chemical analysis of the quality of sachet water consumed in Kano metropolis. *American Journal of Environment, Energy and Power Research* 2, 1-10.
- Wu, Z.Q. & Huang, L.L. (2017). *Fish Biodiversity and its Relationship with Environment in Lijiang River*. Science Press, Beijing.
- Wubie, A., Tadesse, Z., Tesfay, G. & Lakew, A. (2017). Fish distribution and abundance in the Upper Awash River, West Shewa. *Ethiopian Journal of Science and Sustainable Development*, 5(2), 109–118.
- Xing, Y. C., Zhang, C. G., Fan, E. Y. & Zhao, Y. H. (2016). Freshwater fishes of China: species richness, endemism, threatened species and conservation. *Diversity and Distribution*, 22, 358–370. [Crossref]

# Ichthyofauna Diversity and Morphometric Analysis in an Anthropogenically Stressed Water Body Downstream of Jebba Dam, Nigeria

**Solomon Bake Adama**

Department of Animal Biology, Federal University of Technology, Minna, Niger State, Nigeria  
Author

**Kasim Zakari**

Department of Animal Biology, Federal University of Technology, Minna, Niger State, Nigeria  
Author

**Mohammed Mohammed Lemu**

Department of Zoology, Usmanu Danfodiyo University, Sokoto, Sokoto State, Nigeria  
Author

**Yohannah Iliya Auta**

Department of Animal Biology, Federal University of Technology, Minna, Niger State, Nigeria  
Author

**Jonathan Jiya Saba**

Niger State College of Agriculture Mokwa, Niger State, Nigeria  
Author

**Patrick Ozovehe Samuel**

Department of Animal Biology, Federal University of Technology, Minna, Niger State, Nigeria  
Author

**Yakubu Manbe Mohammed**

Department of Biology, Ibrahim Badamasi Babangida University, Lapai, Niger State, Nigeria  
Author

DOI: <https://doi.org/10.56919/usci.2434.028>

**Keywords:** Fish composition, Jebba dam, Length-weight relationship

## Abstract

Fishery resources are on the decline due to incessant and unregulated fishery activities practiced in most parts of the globe. This study investigated the fish composition of an anthropogenically stressed water body in Nigeria (Downstream Jebba Dam). Samples were collected using standard methods and procedures. A total of 6,850 fish belonging to 17 families and 25 species were collected from the sampled stations. Fish composition shows the Mochokidae family with the most dominant species at 29.33% and represented *Synodontis batensoda*, which was well distributed in all the study stations but highest in the station with 934 individuals, while the Gymnarchidae family was the least recorded with 0.16% and represented *Gymnarchus niloticus*. Length-Weight relationship (LWR) value ranged between 0.0018 and 0.215 ( $b < 3$ ) for the dominant fish species, indicating negative allometric growth as the fish species grows in length faster when compared to the weight. In terms of mean condition factor (K of the dominant fish species of Downstream Jebba Dam, *Cynothrissa mento* had the highest with 10.09, followed by *Oreochromis niloticus* (9.82), *Mugil cephalus* (9.02), *Labeo coubie* (3.72), *Bagrus bayad* (0.90), *Alestes macrolepidotus* (0.52), *Clarote laticeps* (0.31), *Momyrus rume* (0.26), *Clarias gariepinus* (0.19) and *Citharinus citharus* (0.16). The high condition factor ( $>1$ ) observed in some fish indicates that these fish species were physiologically stable and successful well, while those with less than ( $<1$ ) mean value implied that it was physiologically unstable, as showed in some species. Overall, the medium abundance of fish species in this study is an indication of an impaired water system.

## References

- Abdulkarim, M., Magaji, I.M., Yusuf, Z.A. & Magaji, N.S. (2020). diversity and spatio-temporal distribution of fish species in Tiga Dam, Kano state. Nigerian Journal of fisheries, 17(1), 1969-1972.
- Abiodun, J.A. & John P. (2017). Biodiversity and abundance of Fish and some processing methods in Lower Niger River, Idah, Kogi State Nigeria. Nigeria Journal of Fisheries and Aquaculture, 5(2), 20-25.
- Abiodun, J.A. & Odunze, F.C. (2011). Fish diversity and fisheries of Jebba Lake Nigeria. Nigeria journal of fisheries, 8(2). 284-290.
- Abowei, F. N. & Hart, A. I. (2014). Some morphometric parameters of 10 finfish species from the lower Nun River, Niger Delta, Nigeria. Research Journal of Biological Sciences. 4(3), 282-288.
- Adadu, M.O., Garba, A.A. & Mshelia, J.P. (2019). Diversity and abundance of fish species in river Okpokwu, Benue State, Nigeria. International Journal of Fisheries and Aquatic Studies, 7 55.60